

Preliminary studies on the phylogeny of *Orthotrichum* (Bryophyta) inferred from nuclear ITS sequences

Jakub Sawicki^{1,*}, Vítězslav Plášek² & Monika Szczecińska¹

¹⁾ Department of Botany and Nature Protection, University of Warmia and Mazury in Olsztyn, Plac Łódzki 1, PL-10-727 Olsztyn, Poland (*corresponding author's e-mail: jakub.sawicki@uwm.edu.pl)

²⁾ Department of Biology and Ecology, University of Ostrava, Slívova 32, CZ-710 00 Slezská Ostrava, Czech Republic

Received 8 July 2008, revised version received 1 Dec. 2008, accepted 3 Dec. 2008

Sawicki, J., Plášek, V. & Szczecińska, M. 2009: Preliminary studies on the phylogeny of *Orthotrichum* (Bryophyta) inferred from nuclear ITS sequences. — *Ann. Bot. Fennici* 46: 507–515.

The study presents a phylogenetic analysis of species of the moss genus *Orthotrichum*. ITS1 and ITS2 for 30 species were sequenced. The results do not fully reflect the current division of *Orthotrichum* into subgenera and sections. Molecular data divide the genus into two groups of species with superficial and immersed stomata, and indicate a clear distinctness of dioecious species. This suggests that a previous concept postulating that the subgenus *Orthophyllum* should be excluded from the genus *Orthotrichum* might be justified.

Key words: *Orthotrichum*, ITS, molecular systematics, mosses, phylogeny

Introduction

The genus *Orthotrichum* is a widespread moss group, which includes approximately 155 species (Goffinet *et al.* 2007) distributed throughout the world from the Arctic to the Antarctic, except in deserts and wet tropical forests. Species of *Orthotrichum* grow on trees and rocks to a height of ca. 5000 m a.s.l. (Lewinsky 1993). In the most recent revision, Orthotrichaceae was divided into two subfamilies, each comprising two tribes: the Schlotheimieae and Macromitriaceae (Macromitrioideae), and the Zygodontaeae and Orthotricheae (Orthotrichoideae), and *Orthotrichum* was placed in the latter group (Goffinet & Vitt 1998, Goffinet *et al.* 2004). The subdivision of *Orthotrichum* has been a moot point since the end of the 19th century. Certain taxa have been alternately

included in and excluded from the genus in the attempts to divide it into lower taxonomic units, subgenera and sections. The basis for the classification of *Orthotrichum* in a historical perspective was described in detail by Lewinsky (1993) and Lewinsky-Haapasaari and Hedenäs (1998).

According to the latest revision, the genus *Orthotrichum* is divided into seven subgenera (Lewinsky 1993): *Callistoma*, *Exiguifolium*, *Gymnopus*, *Orthotrichum*, *Orthophyllum*, *Phaneropus* and *Pulchella*. Based on morphological and molecular data, Goffinet *et al.* (2004) excluded the subgenus *Exiguifolium* from *Orthotrichum* and transferred it to the genus *Letaria*. The subgenera are distinguished based on the following criteria: stoma type (superficial vs. immersed), details of peristome teeth, presence or absence of connecting membrane, cell division of the inner

peristome layer, and ecology. Two of the subgenera, *Gymnopus* and *Pulchella*, are further divided into sections, *Affinia* and *Leiocarpa*, and *Pulchella*, *Diaphana* and *Rivularia*, respectively. The features determining species affiliation to particular sections are usually the details of the structure of the endostome and leaves.

Despite numerous controversies and ambiguities regarding its division (Vitt 1971, Lewinsky 1993, Lewinsky-Haapasaari & Hedenäs 1998), *Orthotrichum* has never been subject to a phylogenetic analysis. The only available information was provided by an analysis of the family Orthotrichaceae, which comprised only several species of the genus *Orthotrichum* (Goffinet et al. 1998, 2004).

The internal transcribed spacer (ITS) region is commonly used in phylogenetic and population genetic studies on bryophytes (Fiedorow et al. 1998, Shaw 2000, Shaw & Allen 2000, Shaw et al. 2005, Juslén 2006, Goryunov et al. 2007, Sawicki & Zielinski 2008, Plášek et al. 2009). In plants, the ITS region is grouped into arrays consisting of hundreds to thousands of tandem repeats. This region includes two spacers, ITS1 and ITS2, that separate the 18S, 5.8S and 26S genes of nuclear ribosomes (Baldwin et al. 1995). A review of the applications of the ITS region in bryophyte systematics is given in Vanderpoorten et al. (2006). Although ITS sequences have proved effective in phylogenetic studies on bryophytes, it should be noted that single gene phylogeny often disagrees with species phylogeny (Miyamoto & Fitch 1995, Maddison 1997). Those differences may stem not only from differing rates of evolution of particular genome regions (Graur & Li 2000), but also from such phenomena as hybridization, common in bryophytes (Natcheva & Cronberg 2004), or horizontal gene transfer (Gustavsson et al. 2005). This concerns primarily chloroplast genes (Shaw et al. 2005, Sawicki et al. 2008), but cases of such incompatibility among nuclear genes have been also reported (Gustavsson et al. 2005, Ghatnekar et al. 2006).

Current morphological revisions of various moss taxa are often supported by molecular data (Virtanen 2003, Hyvönen et al. 2004, Pedersen & Hedenäs 2005, Cano et al. 2005). As compared with morphological data, DNA sequences are not

influenced by the changing environmental conditions in which the plants have grown. Hence, molecular data can be used as a powerful tool in resolving taxonomic and systematic problems. Using nuclear ITS sequences from 28 *Orthotrichum* species, we investigated the phylogeny of *Orthotrichum*. The aim of the study was to find out if the genus *Orthotrichum* and its subgenera and sections are monophyletic based on this data.

Material and methods

Material

Our analyses included 30 species representing three genera of the family Orthotrichaceae. The genus *Orthotrichum* was represented by 28 species representing five subgenera. *Zygodon rupertis* was used as outgroup, based on a previous higher-level analysis (Goffinet et al. 1998, Goffinet et al. 2004).

The list of species used in a molecular analysis, the details concerning voucher data and the GenBank accession numbers are given in Table 1. Since *O. diaphanum* has two substantially different ITS sequences (J. Sawicki unpubl. data), only the sequence showing greater similarity to other species of the genus *Orthotrichum* was used for this analysis. The recently described species *O. moravicum* (Plášek et al. 2009) and the doubtful species *O. fastigiatum*, often treated as *O. affine*, were also analyzed.

DNA extraction

Total genomic DNA was extracted from herbarium material. Single stem was ground with silica beads in a FastPrep tissue disruptor for 20 seconds and subsequently treated processed using the DNAEasy® Plant Mini Kit (Qiagen) following the manufacturer's protocol. Extraced DNA samples were stored at -20°C .

ITS amplification and sequencing

For amplification and sequencing of ITS we used the primers of Fiedorow et al. (1998). The

Table 1. Locality, GenBank accession numbers and ITS sequence length of *Orthotrichum* specimens used in analysis. Herbaria: NYBG = New York Botanical Garden, USA; OP = Opava, Czech Republic; OLS = Olsztyn, Poland.

Species	Provenance, herbarium	Accession number ITS1/ITS2	Sequence length ITS1/ITS2
Subgenus <i>Gymnoporos</i> section <i>Leiocarpa</i>			
<i>O. lyellii</i>	Slovakia, Poloniny Mts., Ruský potok village, herb. OP	EU8663206/EU072689	447/480
<i>O. striatum</i>	Czech Rep., Moravia, Jiřňavské vrchy hills, herb. OP	EU443993/EU072697	428/452
<i>O. vladikavkanum</i>	Tadžikistan, Dushanbe, in Valle Ramit, herb. NYBG	EU8663214/EU871640	429/453
Subgenus <i>Gymnoporos</i> section <i>Affinia</i>			
<i>O. affine</i>	Czech Rep., Prov. Bohemia, Pivoň village, herb. OP	EU8660400/EU072690	429/452
<i>O. fastigiatum</i>	Czech Rep., Prov. Moravia, Lázně Jeseník town, herb. OP	EU8660401/EU072692	428/452
<i>O. pylaisii</i>	Grenlandia, Godthab distr. Kilearsarfik, herb. NYBG	EU8663210/EU871637	420/452
<i>O. sordidum</i>	Armenia, Yerevan, Pambakskij Khrebet, herb. NYBG	EU8663212/EU871639	427/452
<i>O. speciosum</i>	Czech Rep., Prov. Bohemia, Ostrůvek, herb. OP	EU8663213/EU072695	416/452
Subgenus <i>Orthophyllum</i>			
<i>O. gymnostomum</i>	Canada, Newfoundland, Notre Dame Bay, herb. NYBG	EU8663204/EU072687	462/484
<i>O. obtusifolium</i>	Slovakia, Nizké Tatry Mts., Liptovský dvůr, herb. OP	EU8663208/EU072693	643/484
Subgenus <i>Pulchella</i> section <i>Pulchella</i>			
<i>O. consimile</i>	USA, WA, Grays Harbor Co., Lake Quinault, herb. NYBG	EU443997/EU484066	398/435
<i>O. pulchellum</i>	USA, Ciallam Co., Little River Road, herb. NYBG	EU443996/EU484065	405/441
<i>O. scanicum</i>	Czech Rep., Prov. Bohemia, Český les Mts., herb. OP	EU8663211/EU871638	404/434
Subgenus <i>Pulchella</i> section <i>Rivularia</i>			
<i>O. rivulare</i>	Great Britain, S. Devon, R. Bovey, herb. NYBG	EU4840663/EU484070	396/435
Subgenus <i>Pulchella</i> section <i>Diaphana</i>			
<i>O. alpestre</i>	USA, CA, Mono Co. Inyo National Forest, herb. NYBG	EU443998/EU484067	396/435
<i>O. diaphanum</i>	Poland, Młock near Ciechanów, on Salix sp., herb. OLS	EU484077/EU484073	408/431
<i>O. moravicum</i>	Czech Rep., Prov. Moravia, 3 km NE of Bílá, herb. OP	EU8663207/EU072688	403/432
<i>O. pallens</i>	Poland, Góry Bialskie, Bielice village, herb. OP	EU490618/EU072694	396/435
<i>O. patens</i>	Czech Rep., Prov. Moravia, Zábřeh na Moravě, herb. OP	EU8663209/EU871636	397/436
<i>O. pumilum</i>	Poland, Prov. Mazowieckie, Ciechanów, herb. OLS	EU443994/EU035537	402/435
<i>O. stellatum</i>	USA, NY, Putnam Co, Fields Farmstead herb. NYBG	EU484081/EU484068	396/437
<i>O. stramineum</i>	Czech Rep., Prov. Bohemia, Rybník, herb. OP	EU443999/EU072696	397/436
<i>O. tenellum</i>	Canary Islands, Cumbre Neuva, Cumbre, herb. NYBG	EU443995/EU484064	398/436
Subgenus <i>Phaneroporium</i>			
<i>O. laevigatum</i>	Canada, Alberta, 3 km E of Rock Lake, herb. NYBG	EU8663205/EU871635	429/452
<i>O. rupestre</i>	Czech Rep., Prov. Bohemia, Branov, Malá Pleš, herb. OP	EU443991/EU072686	431/454
Subgenus <i>Orthotrichum</i>			
<i>O. anomalum</i>	Poland, Góry Bialskie Mts., Stary Gieraltów, herb. OP	EU443992/EU072691	396/435
<i>O. cupulatum</i>	Poland, Pieniny Mts., Kornajowska skała, herb. NYBG	EU484072/EU484071	396/435
<i>O. pellucidum</i>	USA, Gunnison Co., Cement Creek, herb. NYBG	EU484062/EU484069	396/435
Genus <i>Uloata</i>			
<i>Uloata crispa</i>	Czech Rep., Prov. Moravia, Moravský kras karst, herb. OP	EU8663215/EU871641	469/441
Genus <i>Zygodon</i>			
<i>Zygodon rupestris</i>	Czech Rep., Prov. Silesia, Salajka reserve, herb. OP	EU8663216/EU871642	304/466

sequences of the applied primers were as follow: ITS1-forward 5' CAAGGTTTCCGTAGGTG-AAC 3'; ITS1-reverse 5' CAAGAGCCAAGATATCCG 3'; ITS2-forward 5' CGGATATCTTGGCTCTTG 3'; ITS2-reverse 5' CCGCTTAGTGATATGCTTA 3'. The ITS region was amplified in a volume of 25 μ l containing 20 mM (NH₄)SO₄, 50 mM Tris-HCl (pH 9.0 at 25 °C), 1.5 mM MgCl₂, 1 μ l BSA, 200 μ M each dATP, dGTP, dCTP, dTTP, 1.0 μ M of each primer, one unit of Taq polymerase (Qiagen) and 1 μ l of the DNA solution. The reaction was processed at 94 °C for 1 min. followed by 30 cycles at 94 °C for 1 min., 59 °C for 1 min., and 72 °C for 1.5 min., with a final extension step of 72 °C for 5 min. Finally 5 μ l of the amplification products were visualized on 1.5% agarose gel with ethidium bromide staining. Purified PCR products were sequenced in both directions using ABI BigDye 3.1 Terminator Cycle Kit (Applied Biosystems) and then visualized using an ABI Prism 3130 Automated DNA Sequencer (Applied Biosystems).

Data analyses

Electropherograms were edited and assembled using Sequencher 4.5 (Genecodes Inc.). The assembled sequences were aligned using Muscle 3.6 (Edgar 2004) and manually adjusted with BioEdit 7 (Hall 1999). Phylogenetic analyses were conducted using maximum parsimony (MP), minimum evolution (ME) and Bayesian inference. Gaps were excluded from all phylogenetic analyses. MEGA 4 (Tamura et al. 2007) was used for the Minimum Evolution (ME) analysis and Maximum Parsimony (MP) analysis. The pairwise distances were estimated with the Maximum Composite Likelihood method (Tamura et al. 2004) and initial trees generated using a neighbour-joining (NJ) method. The ME tree was searched using the Close Neighbor Interchange (CNI) algorithm (Nei & Kumar 2000) at the search level of 2, and the maximum number of trees retained at each step was set to 100. For parsimony analyses, we applied branch and bound search as implemented in MEGA 4. Statistical significance of clades within inferred trees was evaluated using

the bootstrap method (Felsenstein 1985) with 1000 replicates.

Bayesian inference was performed using MrBayes 3.12 (Huelsenbeck & Ronquist 2001). The parameters of the likelihood model were those of the general time reversible model (nst = 6) with the proportion of invariable sites in accordance with the best fitted nucleotide evolution model selected on the basis of the Akaike Information Criterion (Akaike 1974) scores in the Modeltest 3.7 (Posada and Crandall 1998). The MCMC algorithm was run for 1 000 000 generations with four incrementally heated chains, starting from random trees and sampling one out of every 100 generations. Trees were checked for stability, which appeared at around 40 000–50 000 chains, therefore, the first 1000 trees were discarded as burn-ins. Remaining trees were used to construct the Bayesian consensus tree. We consider good bootstrap support > 70% and weak support < 70%. In case of the Bayesian clade, credibility values, significant support was estimated as $\geq 95\%$.

Results

The length of the ITS1 spacer ranged from 304 bp in *Zygodon rupestris* to 643 bp in *O. obtusifolium*. The shortest ITS1 sequence among *Orthotrichum* species, 396 bp in length, was found in several taxa with immersed stomata: *O. alpestre*, *O. anomalum*, *O. cupulatum*, *O. pallens*, *O. pelucidum* and *O. rivulare*. The length of the ITS2 spacer ranged from 432 bp in *O. moravicum* to 484 bp in *O. gymnostomum* and *O. obtusifolium*. ITS2 was ca. 20 bp longer in species with superficial stomata than in taxa with immersed stomata.

The alignment had a total length of 1266 bases. The ITS dataset contained 414 variable sites, of which 169 were parsimony informative. A maximum parsimony (MP) analysis resulted in 19 most parsimonious trees of 396 steps, with a consistency index (CI) of 0.7761 and a retention index (RI) of 0.8368. The Minimum Evolution (ME) method (figure not shown) and Bayesian interference (Fig. 1) resulted in very similar trees, differing mostly in the position of *O. lyellii* and *Ulota crispa*. Three main clades were formed. Species of the subgenus *Orthophyllum* formed




Fig. 1. Phylogram based on the Bayesian approach for 28 *Orthotrichum* species with ITS sequence data. Clade credibility values above 95% are given above the branches. Bootstrap values of clades supported under parsimony and Minimum Evolution (set in italics) are given under the branches. Species with immersed stomata are underlined and names of dioecious species are set in bold face.

a distinct, well-supported clade (MP 99% and ME 100% bootstrap support and Bayesian inference 100% clade credibility). The second clade was formed by species with superficial stomata, including *Ulotia crisper* (in the case of MP and ME analysis) which seems to be more closely related to the monoecious species than *O. lyellii*. Within this clade, monoecious species formed a distinct, well-supported group (MP 97% and ME 99% bootstrap support and Bayesian inference 97% clade credibility), however showing no distinct division into subgenera. Only in the case of the minimum evolution analysis, members of the subgenus *Phaneroporos*, *O. laevigatum* and *O. rupestre*, formed their own, but poorly supported clade (ME 44% bootstrap support). Among species of the subgenus *Gymnoporos*, a well-supported clade (MP 92% and ME 93% bootstrap support and Bayesian inference 100% clade credibility) was formed by *O. affine*, *O. fastigiatum*, *O. sordidum* and *O. striatum*, which

does not reflect the current division of this subgenus into sections. The third main clade was formed by species with immersed stomata. Three species of the subgenus *Orthotrichum*, *O. anomalum*, *O. cupulatum* and *O. pellucidum*, formed a distinct, well-supported (ME 84% bootstrap support, Bayesian inference 99% clade credibility) to poorly-supported (MP 62% bootstrap support) clade. Among sections of the subgenus *Pulchella*, only the section *Diaphana* seems to be polyphyletic. Three representatives of the section *Pulchella*, *O. consimile*, *O. pulchellum* and *O. scanicum*, formed their own but poorly supported clade (ME 68% and MP 30% bootstrap support).

Discussion

Molecular data do not support the current division of the genus *Orthotrichum* into subgenera. It seems that the only distinguishable subgenus is

Orthophyllum, whose taxonomic distinctness has been frequently postulated before (Hagen 1908, Damsholt *et al.* 1969, Goffinet *et al.* 2004). Its two species *O. gymnostomum* and *O. obtusifolium* were indeed placed in a separate genus, *Stroemia*, by Hagen (1908). They were separated by obtuse leaves with incurved or plane leaf margins and incrassate leaf cells with a stout, central papilla on each side. Since *Stroemia* was an illegitimate name, it was later replaced by *Nyholmiella* (Damsholt *et al.* 1969). A later revision of the genus *Orthotrichum* resulted in the inclusion of *O. gymnostomum* and *O. obtusifolium* into *Orthotrichum* (Vitt 1973), as the above features were also observed in other representatives of this genus. The affiliation of these species to the genus *Orthotrichum* was tested by Lewinsky-Haapasaari and Hedenäs (1998) with the use of cladistic methods. However, an analysis of the selected morphological characters of the above taxa did not confirm their distinctness sufficiently to place them into a separate genus. On the other hand, an analysis of a nuclear ITS sequence conducted in this study suggested that the members of the subgenus *Orthophyllum* are genetically distinct from the other species of *Orthotrichum*. In addition, the investigated sequence revealed a closer relationship between *Ulotia crispata* and other *Orthotrichum* species than between these species and *O. gymnostomum* and *O. obtusifolium*, which strongly supports separation of the latter two species from *Orthotrichum*. The distinctness of *O. obtusifolium* from the other species of *Orthotrichum* was also revealed by an analysis of sequences from four loci (26S, *nad5*, *rps4* and *trnL-trnF* region), which provided a basis for excluding species of the subgenus *Orthophyllum* from *Orthotrichum* (Goffinet *et al.* 2004). Certainly, such a decision should be based on an analysis of a greater number of genes and populations of these species.

Our results also support considerable distinctness of other species with superficial stomata from those with immersed stomata. Stoma type is important in the taxonomy of *Orthotrichum*. They are either level, with surrounding exothecial cells (superficial, phaneropore), or immersed and almost covered by the surround-

ing cells (immersed, cryptopore). The occurrence of both stomata types on all continents suggests that both types have a long history. However, their historical relationships are still unclear (Lewinsky 1977). Vitt (1971) tried to explain the fact as a genetic change, because the species with superficial stomata have a haploid chromosome number, $n = 6$, whereas cryptopore species have $n = 11$. However, autopolyploidy plus the loss of one chromosome seem to be unlikely to create the immersed type of stomata, because it would mean that this feature is controlled by only one chromosome. The separation of the two types of stomata is probably due to a much more complex mechanism, dependent on the interaction between genes in different chromosomes (Lewinsky 1977). According to Paton and Pearce (1957), the development of cryptopore stomata as a result of adaptation to xeric conditions seems unlikely because many species with immersed stomata are found in moist habitats (*O. pulchellum*, *O. consimile*). The position of stomata provided a basis for the division of this genus by Lindberg (1879), who divided it into two subgroups, *Gymnoporos* and *Calypatoros*, comprising species with superficial and immersed stomata respectively. Within the first of those subgroups, taxa belonging to two subgenera, *Gymnoporos* and *Phaneroporos*, did not form clades in our study. A clear division of the genus *Orthotrichum* into groups of species with immersed and superficial stomata was also confirmed by chloroplast and mitochondrial sequences (Goffinet *et al.* 2004). An analysis of the nuclear locus 26S, mitochondrial locus *nad5* and chloroplast loci *rps4* and *trnL-trnF* revealed the distinctness of *O. affine*, *O. laevigatum* and *O. lyellii* from *O. alpestre*, *O. anomalum*, *O. assimile* and *O. macrocephalum*, placing these species in two separate clades.

Molecular data do not support the division of the subgenus *Gymnoporos* into the sections *Leiocarpa* and *Affinia*, as proposed by Lewinsky (1993) and Lewinsky-Haapasaari and Hedenäs (1998). The type taxon of the section *Leiocarpa*, *O. striatum*, was included in a moderately supported clade together with *O. affine*, *O. fastigiatum* and *O. sordidum* of the section *Affinia*. Another member of the section *Leiocarpa*, the

dioecious *O. lyellii*, was clearly distinct from the other species of the subgenus *Gymnopus*. A large number of fixed differences in relation to the other species in the subgenus suggests that this taxon should be placed into a separate infrageneric taxon. Its exclusion from the section *Leiocarpa* and the formation of the section *Lyelliana* by Schimper (1876, followed Lewinsky 1993), seems justified. The distinctness of *O. lyellii* from members of the section *Leiocarpa* was also noted by Vitt (1971), who however did not classify this taxon into a separate section or subgenus. Based on the ITS sequences *Ulotia crispera* is closely related to *Orthotrichum* species of the subgenera *Gymnopus* and *Phaneroporium*. *Gymnopus* species were found to be more genetically similar to *U. crispera* than to the dioecious *O. lyellii* belonging to the same subgenus. Similar results were obtained while analyzing the chloroplast *rbcl* sequence which showed a close relationship between *Ulotia* and *Orthotrichum* species of the subgenus *Gymnopus* (Goffinet *et al.* 1998). An analysis of four loci from the nuclear (26S), mitochondrial (*nad5*) and chloroplast (*rps4*, *trnL-trnF*) genome also showed that *Orthotrichum* species with superficial stomata are closer to species of *Ulotia* than to *Orthotrichum* species with immersed stomata of the subgenera *Orthotrichum* and *Pulchella* (Goffinet *et al.* 2004).

A further group was formed by species of the subgenera *Orthotrichum* and *Pulchella*. Unlike species with superficial stomata, representatives of the subgenus *Orthotrichum* formed a clade. As regards morphological characteristics, subgenus *Orthotrichum* differs from subgenus *Pulchella* mostly in having straighter peristome teeth and no connecting membrane (Lewinsky 1993). The placing of species of this subgenus within the 'Pulchella' group is consistent with the results of Vitt (1971) and with one of the three relationship ideas proposed by Lewinsky-Haapasaari and Hedenäs (1998). In both cases, the authors placed the subgenus *Orthotrichum* within the *Pulchella* clade, based on morphological characters. However, our results did not confirm a close relationship between members of the subgenera *Orthotrichum* and *Phaneroporium*, suggested by Lewinsky (1993). Taxa belonging

to these groups are genetically different, despite certain similarities such as the preferred habitats and the characteristics of peristome and calyptra.

For the subgenus *Pulchella*, in contrast to the subgenus *Gymnopus*, our study resulted in a moderately supported division into sections. The only clearly distinct section was *Pulchella*, whose species were grouped within a single clade in all analyses. The section *Pulchella* could be considered monophyletic, while the largest section, *Diaphana*, is rather paraphyletic. The position of *O. rivulare* may also indicate a genetic distinctness of the section *Rivularia*.

The number and ornamentation of exostome teeth and endostome segments have a central role in the taxonomy of *Orthotrichum*. The ancestors of the genus probably had a very well developed double peristome (with 16 exostome teeth and endostome segments) covered with papillae (Lewinsky 1977). Vitt (1973) suggests that a reduction in peristome characters (mainly in the number of endostome segments and exostome teeth, from 16 to 8 or 0) may result from adaptation to different ecological conditions. According to that author, specialization to xeric habitats, such as the trunks of trees and dry rock surfaces, is the driving force. This view, however, cannot be regarded as generally valid, as a reduction of endostome segments has not been observed in many xeric species growing in dry places (*O. diaphanum*, *O. striatum*), whereas a species (*O. affine*) of shaded and moist places has a reduced peristome of eight teeth and segments. The presence of a preperistome, considered by Vitt (1973) as an advanced character, was subsequently regarded as a primitive state, because it is present in many species with a well-developed, not reduced peristome (Lewinsky 1989). We found no correlation between particular peristome characters and adaptive evolution in *Orthotrichum*.

Although our results do not fully reflect the current taxonomic division, it points to the fact that the former taxonomic concepts were often correct. This refers in particular to the division proposed by Lindberg (1879) and to the separation of the genus *Stroemia* (Hagen 1908), which seems justified. However, this remains a hypothesis with some support and it should be confirmed by further studies.

Acknowledgements

DNA analysis was made possible through financial support from The Lewis B. and Dorothy Cullman Program for Molecular Systematic Studies. We also thank W. R. Buck for lending us herbarium specimens of several *Orthotrichum* species, and two anonymous referees for helpful comments on the manuscript.

References

- Akaike, H. 1974: A new look at the statistical model identification. — *IEEE Transactions on Automatic Control* 19: 716–723.
- Baldwin, B. G., Sanderson, M. J. & Porter, J. M. 1995: The ITS region of nuclear ribosomal DNA: a valuable source of evidence on angiosperm phylogeny. — *Ann. Missouri Bot. Garden* 82: 247–277.
- Cano, M. J., Werner, O. & Guerra, J. 2005: A morphometric and molecular study in *Tortula subulata* complex (Pottiaceae, Bryophyta). — *Bot. J. Linn. Soc.* 149: 333–350.
- Damsholt, K., Holmen, K. & Warncke, E. 1969: A list of the bryophytes of Denmark. — *Bot. Tidsskr.* 65: 163–183.
- Edgar, R. C. 2004: Muscle: multiple sequence alignment with high accuracy and high throughput. — *Nucleic Acid Res.* 32: 1792–1797.
- Felsenstein, J. 1985: Confidence limits on phylogenies: An approach using the bootstrap. — *Evolution* 39: 783–791.
- Fiedorow, P., Odrzykoski, I. & Szweykowska-Kulińska, Z. 1998: Phylogenetic studies of liverworts using molecular biology techniques. — In: Małuszyńska, J. (ed.), *Plant cytogenetics*: 244–249. Silesian University Press, Katowice.
- Ghatnekar, L., Jaarola, M. & Bengtsson, B. O. 2006: The introgression of a functional nuclear gene from *Poa* to *Festuca ovina*. — *Proc. R. Soc. London B* 273: 395–399.
- Goffinet, B. & Vitt, D. H. 1998: A revised classification of the Orthotrichales (Bryopsida) based on comparative morphology and a *rbcL* based phylogeny. — In: Bates, J. W., Ashton, N. W. & Duckett, J. G. (eds.), *Bryology for the twenty-first century*: 143–159. Maney Publ. & Brit. Bryol. Soc., London.
- Goffinet, B., Bayer, R. J. & Vitt, D. H. 1998: Circumscription and phylogeny of the Orthotrichales (Bryopsida) inferred from *rbcL* sequence analyses. — *Am. J. Bot.* 85: 1324–1337.
- Goffinet, B., Shaw, A. J., Cox, C. J., Wickett, N. J. & Boles, S. 2004: Phylogenetic inferences in the Orthotrichoideae (Orthotrichaceae: Bryophyta) based on variation in four loci from all genomes. — *Monogr. Syst. Bot. Missouri Bot. Garden* 98: 270–289.
- Goffinet, B., Buck, W. R. & Wall, M. A. 2007: *Orthotrichum freyanum* (Orthotrichaceae, Bryophyta), a new epiphytic species from Chile. — *Nova Hedwigia* 131: 1–11.
- Goryunov, D. V., Ignatova, E. A., Ignatov, M. S., Milyutina, I. A. & Troitsky, A. V. 2007: Support from DNA data for narrow species concept in *Schistidium* (Grimmiaceae, Musci). — *J. Bryol.* 29: 98–103.
- Graur, D. & Li, W.-H. 2000: *Fundamentals of molecular evolution*. — Sinauer Assoc., Sunderland, MA.
- Gustavsson, S., Lebrun, A.-S., Norden, K., Chaumont, F. & Johanson, U. 2005: A novel plant major intrinsic protein in *Physcomitrella patens* most similar to bacterial glycerol channels. — *Pl. Phys.* 139: 287–295.
- Hagen, I. 1908: Forarbejder til en norsk lovmosflora. I. Orthotrichaceae. — *Det Kongelige Norske Videnskabers Selskabs Skrifter* 1907, 13: 1–100.
- Hall, T. A. 1999: BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. — *Nucl. Acids. Symp. Ser.* 41: 95–98.
- Huelsensbeck, J. P. & Ronquist, F. R. 2001: MRBAYES: Bayesian inference of phylogeny. — *Bioinformatics* 17: 754–755.
- Hyvönen, J., Koskinen, S., Merrill, G. L. S., Hedderson, T. A. & Stenroos, S. 2004: Phylogeny of Polytrichales (Bryophyta) based on simultaneous analysis of molecular and morphological data. — *Mol. Phyl. Evol.* 31: 915–928.
- Juslén, A. 2006: Phylogeny of Vetaformaceae, Lepicoleaceae and Herbertaceae (including Mastigophoraceae) inferred from chloroplast *trnL-F*, nuclear ITS2, and morphology. — *Ann. Bot. Fennici* 43: 349–362.
- Lewinsky, J. 1977: The genus *Orthotrichum*. Morphological studies and evolutionary remarks. — *J. Hattori Bot. Lab.* 43: 31–61.
- Lewinsky, J. 1989: Does the orthotrichaceous type of peristome exist? — *J. Hattori Bot. Lab.* 67: 335–363.
- Lewinsky, J. 1993: A synopsis of the genus *Orthotrichum* Hedw. (Musci, Orthotrichaceae). — *Bryobrothera* 2: 1–59.
- Lewinsky-Haapasaari, J. & Hedenäs, L. 1998: A cladistic analysis of the moss genus *Orthotrichum*. — *Bryologist* 101: 519–555.
- Lindberg, S. O. 1879: *Musci Scandinavici in systemate novo naturali dispositi*. — Iesaae Edquist, Uppsala.
- Maddison, W. P. 1997: Gene trees in species trees. — *Syst. Biol.* 46: 523–536.
- Miyamoto, M. M. & Fitch, W. M. 1995: Testing species phylogenies and the phylogenetic methods with congruence. — *Syst. Biol.* 44: 64–76.
- Natcheva, R. & Cronberg, N. 2004: What do we know about hybridization among bryophytes in nature? — *Can. J. Bot.* 82: 1687–1704.
- Nei, M. & Kumar, S. 2000: *Molecular evolution and phylogenetics*. — Oxford Univ. Press, New York.
- Paton, J. A. & Pearce, J. V. 1957: The occurrence, structure and functions of stomata in British bryophytes. — *Trans. Brit. Bryol. Soc.* 3: 228–259.
- Pedersen, N. & Hedenäs, L. 2005: Taxonomic and nomenclatural implications of phylogenetic studies of the Bryaceae based on molecular data and morphology. — *Bryologist* 108: 123–128.
- Plášek, V., Sawicki, J., Trávníčková, V. & Pasečná, M. 2009: *Orthotrichum moravicum* (Orthotrichaceae), a new moss species from the Czech Republic. — *Bryologist* 112: 329–336.
- Posada, D. & Crandall, K. A. 1998: Modeltest: testing

- the model of DNA substitution. — *Bioinformatics* 14: 817–818.
- Sawicki, J. & Zieliński, R. 2008: Phylogenetic relationships between five *Sphagnum* species of the section *Acutifolia* Based on DNA markers. — *Cas. Sleż. Muż. Opava* 57: 63–80.
- Sawicki, J., Plášek, V. & Szczecińska, M. 2008: Testing candidate *trnH-psbA* noncoding plant barcode region in the *Orthotrichum* genus. — *Scripta Fac Rerum Natur. Univ. Ostraviensis. Environmental Changes and Biological Assessment IV*: 170–177.
- Shaw, A. J. 2000: Molecular phylogeography and cryptic speciation in the mosses *Mielichhoferia elongata* and *M. mielichhoferiana* (Bryaceae). — *Mol. Ecol.* 9: 595–608.
- Shaw, A. J. & Allen, B. H. 2000: Phylogenetic relationships, morphological incongruence and geographic speciation in the Fontinalaceae (Bryophyta). — *Mol. Phyl. Evol.* 16: 225–237.
- Shaw, A. J., Cox, C. J. & Boles, S. B. 2005: Phylogeny, species delimitation, and recombination in *Sphagnum* section *Acutifolia*. — *Syst. Bot.* 30: 16–33.
- Schimper, W. P. 1876: *Synopsis Muscorum Europaeorum praemissa introductione de elementis bryologicis tractante*. — Stuttgartia.
- Tamura, K., Nei, M. & Kumar, S. 2004: Prospects for inferring very large phylogenies by using the neighbor-joining method. — *Proc. Nat. Acad. Sci. USA* 101: 11030–11035.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. 2007: MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. — *Mol. Biol. Evol.* 24: 1596–1599.
- Vanderpoorten, A., Goffinet, B. & Quandt, D. 2006: Utility of the internal transcribed spacers of the 18S-5.8S-26S nuclear ribosomal DNA in land plant systematics with special emphasis on Bryophytes. — In: Sharma, A. K. & Sharma, A. (eds.), *Plant genome: biodiversity & evolution*, vol. 2B: Lower plants: 385–407. Sci. Publ., Enfield.
- Virtanen, V. 2003: Phylogeny of Bartramiaceae based on morphology and on *rbcL*, *rps4* and *trnL-F* sequence data. — *Bryologist* 106: 280–296.
- Vitt, D. H. 1971: The infrageneric evolution, phylogeny, and taxonomy of the genus *Orthotrichum* (Musci) in North America. — *Nova Hedwigia* 21: 683–711.
- Vitt, D. H. 1973: A revision of genus *Orthotrichum* in North America, north of Mexico. — *Bryophyt. Biblioth.* 1: 1–208.