

Aspidistra jingxiensis (Asparagaceae), a new species from Guangxi, China

Chun-Rui Lin, Wei-Bin Xu, Yu-Song Huang & Yan Liu*

Guangxi Institute of Botany, Guangxi Zhuangzu Autonomous Region and the Chinese Academy of Sciences, Guilin, 541006, China (*corresponding author's, e-mail: gxibly@163.com)

Received 3 Aug. 2011, final version received 25 Sep. 2011, accepted 27 Sep. 2011

Lin, C. R., Xu, W. B., Huang, Y. S. & Liu, Y. 2012: *Aspidistra jingxiensis* (Asparagaceae), a new species from Guangxi, China. — *Ann. Bot. Fennici* 49: 193–196.

A new species *Aspidistra jingxiensis* Yan Liu & C.R. Lin (Asparagaceae) is described and illustrated from the limestone areas in the Guangxi Zhuangzu Autonomous Region, China. It is similar to *A. punctatoides* in the shape and size of the flowers, but differs in its perianth lobes 8 (occasionally 10), mushroom-shaped and white pistil, and the glabrous stigma upper surface with four radial, bifurcate deep grooves.

More than 100 species of the genus *Aspidistra* (Asparagaceae) — whose main centre of diversity is in China and Vietnam — have formally been described. In recent years, we had carried out field surveys in limestone areas, and several new species of *Aspidistra* have been described (Hou *et al.* 2009, Lin *et al.* 2009, 2010, Lin & Liu 2011, Liu *et al.* 2011).

During a field trip in the year 2000 to Jingxi County, Guangxi, near the border with northern Vietnam, we collected non-flowering specimens of *Aspidistra*. The plants were transferred to the Guilin Botanical Garden where they were cultivated for further study. The plants produced purple flowers in 2003, and in the following years, they flowered regularly. During the course of investigating limestone plants in June 2011, we collected a flowering plant near Jingxi County again. After consulting the relevant literature (Lang *et al.* 1999, Liang & Tamura 2000, Li 2004, Tang & Liu 2003, Tillich 2005, 2008, Xu *et al.* 2010) as well as herbarium specimens, we concluded that our specimens represented a new species, which is described here.

Aspidistra jingxiensis Yan Liu & C.R. Lin, *sp. nova* (Figs. 1 and 2)

TYPE: China. Guangxi Zhuangzu Autonomous Region, Jingxi County, alt. 635 m, limestone mountains, 7 June 2011, *Chun-Rui Lin & Yan Liu* 1102 (holotype IBK; isotype IBK).

ETYMOLOGY: The specific epithet is derived from the name of the type locality, Jingxi County, Guangxi Zhuangzu Autonomous Region, China.

Herbs perennial, evergreen, rhizomatous. Rhizome creeping, subterete, 8–10 mm thick, covered with scales, nodes dense. Vaginal leaves 3–4, purple-red, 2–12 cm long, enveloping base of petiole, becoming black-brown when dry. Leaves solitary, ca. 5 mm apart; petiole stiff upright, 6–24 cm long, 3–4 mm thick, adaxially sulcate; leaf blade usually narrow elliptic to oblong-lanceolate, 28–35 cm long, 6–8 cm wide, dark green with small yellow-white spots on both surfaces, base cuneate to broadly cuneate, inequilateral, apex acuminate, margin entire. Peduncle decumbent or declining, 3–5 cm long, with 4–5 bracts, bracts gradually wider from base to top of peduncle, two most distal bracts




Fig. 1. *Aspidistra jingxiensis* (drawn from the holotype by Xi-Yun Zhu). — **A:** Flowering plant. — **B:** Flower with half of perianth removed showing stamens and pistil. — **C:** Perianth, dissected open to show stamens. — **D:** Stigma apical view. — **E:** Pistil.

adjacent to perianth broadly ovate-cucullate, purplish red, 6–8 mm long, 5–6 mm wide, apex subobtusate. Flowers solitary; perianth broadly campanulate, 10–12 mm long, fleshy, pale green-yellow with purple spots outside; lobes 8 (occasionally 10), usually suberect, ovate-triangular, 5–6 mm long, 3–4 mm wide at base, adaxially pale green-yellow with purple spots, and with 2–3 keels at base; tube 5–7 mm long, 8–10 mm in diameter, blackish purple inside. Stamens 8(10), opposite to lobes, inserted in middle of perianth tube, positioned lower than stigma, filaments horizontal, ca. 0.6 mm long, anthers

oblong, pale yellow, 2–3 mm long and 1–1.5 mm wide. Pistil mushroom-shaped, white, ca. 6 mm long, ovary inconspicuous, style cylindrical, ca 4 mm long and 2 mm in diameter, stigma peltate, glabrous, 5–6 mm in diameter, upper surface nearly flat and with 4 radial, bifurcate deep grooves, 4-lobed at margin, lobes emarginate at apex. Flowering from May to June.

In the wild, *Aspidistra jingxiensis* is known only from the vicinity of the Jingxi county seat, in southwestern Guangxi Zhuang Autonomous Region, China, that borders northern Vietnam. It grows on shaded limestone slopes in evergreen




Fig. 2. A–E: *Aspidistra jingxiensis*. — A: Habit. — B: Flower from above. — C: Perianth, dissected open to show stamens. — D: Pistil. — E: Flower side view. — F and G: *A. punctatoides*. — F: Flower dissected, showing stamens. — G: Flower from above.

broad-leaved forests at an altitude of 600–700 m.

Aspidistra jingxiensis is similar to *A. punctatoides* (Fig. 2F–G; see also Lin & Liu 2011) in the shape and size of the flowers, but differs in several characters (see Appendix).

ADDITIONAL SPECIMEN EXAMINED: China. Guangxi Zhuangzu Autonomous Region, Guilin City, Yanshan Township, introduced by Yan Liu from the type locality, cultivated, 12 June 2010, *Chun-Rui Lin 039* (IBK).

Acknowledgements

We are grateful to Mr. Xi-Yun Zhu (IBK) for the drawing. This study was supported by the Special Fund for Basic Sci-

entific Research of the Guangxi Institute of Botany (09015) to Chun-Rui Lin (IBK); the Western Program for Fostering Personal Ability, CAS (2007) and the Knowledge Innovation Project of the Chinese Academy of Sciences, grant no. KSCX2-YW-Z-0912 to Yan Liu (IBK).

References

- Hou, M. F., Liu, Y., Kono, Y. & Peng, C. I. 2009: *Aspidistra daxinensis* (Ruscaceae), a new species from limestone areas in Guangxi, China. — *Botanical Studies* 50: 371–378.
- Lang, K. Y., Li, G. Z., Liu, Y., Wei, Y. G. & Wang, R. X. 1999: [Taxonomic and phylogeographic studies on the genus *Aspidistra* Ker-Gawl., (Liliaceae) in China]. — *Acta Phytotaxonomica Sinica* 37: 468–508. [In Chinese].

- Li, G. Z. 2004: [*The genus Aspidistra*]. — Guangxi Sci. & Technol. Publ. House, Nanning. [In Chinese].
- Liang, S. Y. & Tamura, M. N. 2000: *Aspidistra*. — In: Wu, Z. Y. & Raven, P. H. (eds.), *Flora of China*, vol. 24: 240–250. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis.
- Lin, C. R., Liang, Y. Y. & Liu, Y. 2009: *Aspidistra bamaensis* (Ruscaceae), a new species from Guangxi, China. — *Annales Botanici Fennici* 46: 416–418.
- Lin, C. R. & Liu, Y. 2011: *Aspidistra punctatoides* sp. nov. (Ruscaceae) from limestone areas in Guangxi, China. — *Nordic Journal of Botany* 29: 189–193.
- Lin, C. R., Peng, C. I., Kono, Y. & Liu, Y. 2010: *Aspidistra obconica* (Ruscaceae), a new species from limestone areas in Guangxi, China. — *Botanical Studies* 51: 263–268.
- Liu, Y., Kono, Y., Lin, C. R., Xu, W. B. & Peng, C. I. 2011: *Aspidistra erecta* (Asparagaceae), a new species from limestone areas in Guangxi, China. — *Botanical Studies* 52: 363–369.
- Tang, S. C. & Liu, Y. 2003: *Aspidistra guangxiensis* (Con-vallariaceae), a new species from China. — *Novon* 13: 480–482.
- Tillich, H. J. 2005: A key for *Aspidistra* (Ruscaceae), including fifteen new species from Vietnam. — *Feddes Repertorium* 116: 313–338.
- Tillich, H. J. 2008: An updated and improved determination key for *Aspidistra* Ker-Gawl. (Ruscaceae, Monocotyledons). — *Feddes Repertorium* 119: 449–462.
- Xu, W. F., He, H. Z. & Yang, L. 2010: *Aspidistra chishuiensis* (Ruscaceae), a new species from Guizhou, China. — *Annales Botanici Fennici* 47: 118–120.

Appendix. Morphological comparison between *Aspidistra jingxiensis* and *A. punctatoides*.

	<i>A. jingxiensis</i>	<i>A. punctatoides</i>
Leaf blade	28–35 × 6–8 cm, dark green with small yellow-white spots on both surfaces	15–27 × 4.5–7.5 cm, dark green
Petiole	6–24 cm long	11–26 cm long
Peduncle	3–5 cm long	2–3 cm long
Perianth	lobes 8(10), 5–6 × 3–4 mm, adaxially with 2–3 keels at base; tube 5–7 mm long, diam. 8–10 mm	lobes 8(6), 5–6 × 3–4 mm, adaxially with two prominent keels ca. 1 mm high, each keel basally fusing with a keel of the adjoining lobe and forming a protruding lip at fusion point; tube 5–7 mm long, diam. 10–14 mm
Pistil	mushroom-shaped, white, ca. 6 mm long stigma diam. 5–6 mm, upper surface glabrous and with 4 radial, bifurcate deep grooves	turbinate, white adaxially, purple abaxially, stigma ca. 8 mm long, diam. 10 mm, upper surface densely papillate and with 3–4 inconspicuous radial, bifurcate lines