

Genetic evidence of a progenitor–derivative species pair in East Asian *Lilium*

Mi Yoon Chung^{1,3}, Jordi López-Pujol² & Myong Gi Chung^{3,*}

¹ Department of Biology Education, Seowon University, Cheongju 361-742, Republic of Korea

² BioC, GReB, Botanic Institute of Barcelona (IBB-CSIC-ICUB), Passeig del Migdia s/n, ES-08038 Barcelona, Spain

³ Department of Biology and the Research Institute of Natural Science, Gyeongsang National University, Jinju 660-701, Republic of Korea (*corresponding author's e-mail: mgchung@gnu.ac.kr)

Received 31 Jan. 2014, final version received 27 Jan. 2015, accepted 12 Feb. 2015

Chung M.Y., López-Pujol J. & Chung M.G. 2015: Genetic evidence of a progenitor–derivative species pair in East Asian *Lilium*. — *Ann. Bot. Fennici* 52: 211–218.

Boreal *Lilium distichum* and temperate *L. tsingtauense* are morphologically very similar, thus they have been placed in the section *Martagon*. Recent molecular phylogenetic analyses revealed that *L. distichum* and *L. tsingtauense* are indeed the most closely related species within that section. *Lilium distichum* has a wider geographic range and a broader niche than *L. tsingtauense*. We hypothesized that *L. distichum*–*L. tsingtauense* might be a classical “progenitor–derivative” (P–D) species pair and examined the levels of allozyme diversity in the two species in South Korea. Whereas the allelic composition of *L. tsingtauense* represented a subset of *L. distichum*, the former had significantly lower allozyme variability at both the population and the species levels than the latter. Except for the locus *Fe* (fluorescent esterase), allele frequencies of *L. distichum* were very similar to those of *L. tsingtauense*. Accordingly, pairwise genetic identities between populations of *L. distichum* and *L. tsingtauense* were very high, with a mean of 0.919. Our allozyme results support the hypothesis that *L. tsingtauense* is a derivative species of the progenitor *L. distichum*.

Introduction

Lilium consists of approximately 100 species that are widely distributed throughout boreal and temperate regions of the northern hemisphere (MacRae 1998). Among them, *L. distichum* is a boreal species usually occurring on forested slopes, forest margins, and hillsides along streams (200–1800 m), distributed in Russian Far East (Primorsky Krai), northeastern China (Heilongjiang, Jilin, and Liaoning provinces) and the Korean Peninsula (Liang & Tamura 2000;

Fig. 1). In Korea, *L. distichum* occurs in grasslands or gaps of *Quercus mongolica*-dominated temperate deciduous forests, at altitudes of 900–1600 m a.s.l. along the Baekdudaegan (the main mountain system of Korea, which runs through most of the length of the Korean Peninsula) and in one of its main branches, the Nakdongjeongmaek (Fig. 1). The species is relatively common where it occurs, with the number of individuals per population on the order of hundreds. Its congener *L. tsingtauense* is a temperate species with a narrower range, distributed in eastern




Fig. 1. Locations of sampled populations of *Lilium distichum* (LD-1 to LD-7) and *L. tsingtauense* (LT-1 to LT-11) in South Korea. Sample sizes: LD-1 = 47, LD-2 = 24, LD-3 = 44, LD-4 = 21, LD-5 = 27, LD-6 = 45, LD-7 = 37; LT-1 = 18, LT-2 = 48, LT-3 = 41, LT-4 = 29, LT-5 = 37, LT-6 = 22, LT-7 = 36, LT-8 = 38, LT-9 = 42, LT-10 = 58, LT-11 = 74. The solid line indicates the location and shape of the main mountain range of the country, the Baekdu-daegan, which runs from north to south along the Korean Peninsula (spanning over 1600 km). The dotted line represents the Nakdongjeongmaek, one of the 13 mountainous branches of the Baekdu-daegan. The approximate geographic ranges of both *Lilium* species are shown in the inset map (dotted line, *L. distichum*; dashed line, *L. tsingtauense*).

China (Shandong and Anhui provinces) and in the Korean Peninsula (Liang & Tamura 2000; Fig. 1). In China, it occurs on sunny forested slopes or in bushy or grassy places, at a wide range of elevations (100–1000 m a.s.l.; Liang & Tamura 2000, Guo *et al.* 2011). In Korea, *L. tsingtauense* is mainly found in relatively open habitats in low-altitude mountains (0–500 m; hereafter we define low elevation as < 500 m, mid elevation as 500–1000 m, and high elevation as > 1000 m) across the Peninsula, including peripheral regions of the Baekdudaegan and the islands. Isolated individuals, however, can be found at mid to high elevations in montane habitats on the main ridge of the Baekdudaegan. Like

L. distichum, *L. tsingtauense* is locally common, with populations of hundreds of individuals.

Because of their similar morphology, *L. distichum* and *L. tsingtauense* were placed in the same section *Martagon*, one of the seven sections in *Lilium* proposed by Comber (1949). A close relationship has been further corroborated by molecular phylogenetic analyses, which clearly point out that *L. distichum* and *L. tsingtauense* are the most closely related species within this section (Sultana *et al.* 2010, 2011, Lee *et al.* 2011). They are sympatric in some locations in mid-elevation mountains in Korea (e.g. LD-2, 902 m, and LT-1, 889 m, at Mt. Deokhang; Fig. 1). However, *L. distichum* has a much wider geographic range, with

a broader niche (as estimated from ecological niche modelling analyses in Chung *et al.* 2014a) than *L. tsingtauense*. *Lilium distichum* thrives in places with temperatures ranging from -21.5°C to 24.8°C (mean temperatures in winter and summer, respectively) and precipitation ranging from 550 mm to 1500 mm, whereas *L. tsingtauense* has narrower temperature and precipitation ranges (from -7°C to 24.6°C , and from 700 mm to 1500 mm, respectively). Based on phylogenetic information, ecological traits, and distribution data, we hypothesize that *L. distichum*–*L. tsingtauense* might constitute a “progenitor–derivative” (hereafter P–D) species pair; that is, instead of true sister species pair, a D or “derived” species budded off and acquired new traits while the P or “parental” species remained largely unchanged (Gottlieb 1984, 2003, Crawford 1985, 1990, 2010). Under this scenario, we can expect three genetic outcomes. First, the spectrum of alleles observed in *L. tsingtauense* should be a subset of those found in *L. distichum*, with few, if any, unique alleles. Second, levels of within-population genetic variation observed in *L. tsingtauense* should be lower than in *L. distichum*. Third, interspecific genetic identities would be similar to or slightly lower than intraspecific identities, as a result of the recent divergence of the D species (Gottlieb 1973, 2003, Crawford 1983, 2010).

Material and methods

Study plants

Lilium distichum is a herb 60–120 cm tall, with a

papillose stem and leaves in a whorl of 7–9 near the middle of stem. Two to seven orange to pale vermilion flowers with purple-red spots (tepals are ca. 5.0 cm long) are arranged in racemes, nodding to horizontal, and open from July to August. Fruits (capsules) are obovoid, 1.5 cm long. *Lilium tsingtauense* is a somewhat shorter plant (40–85 cm tall), with leaves in one or two whorls of 5–14. Orange to vermilion flowers with purple-red spots are solitary or in racemes of up to seven, of similar size to those of *L. distichum*, and open from June to July. Capsules are also ca. 1.5 cm long. The breeding systems and pollinators for the two *Lilium* species are unknown, although many lilies are known to be insect-pollinated and self-compatible (Yang & Sun 2005, Arzate-Fernández *et al.* 2007, Rodger *et al.* 2010) with few exceptions (e.g. *L. longiflorum* is generally regarded as self-incompatible; Miller 1993).

Population sampling

For *L. distichum*, we collected 245 individuals from seven populations (Table 1 and Fig. 1), five from the main ridge of the Baekdudaegan (LD-1 to LD-3, LD-6, and LD-7) and two from the Nakdongjeongmaek (LD-4 and LD-5). For *L. tsingtauense*, we sampled a total of 443 individuals from 11 populations (Table 1 and Fig. 1), mostly located on the low- to mid-elevation peripheral areas of the Baekdudaegan, although three populations (LT-9, LT-10, and LT-11) were located on Oenaro Island, on the southern coast of the Korean Peninsula (Fig. 1). As populations

Table 1. Levels of genetic diversity of *Lilium distichum* and *L. tsingtauense* in South Korea estimated from 7 and 11 populations, respectively. Statistics for levels of within-population genetic variation are provided in Chung *et al.* (2014a: table 1). Abbreviations: n = number of individuals sampled, %P = percentage of polymorphic loci, AR = mean allelic richness, A = mean number of alleles per locus, H_o = observed heterozygosity, H_e = H-W expected heterozygosity or genetic diversity, SE = standard error.

Species	n	%P	AR	A	H_o (SE)	H_e (SE)
<i>Lilium distichum</i>						
population average	35	65.3	1.90	1.94	0.151 (0.012)	0.190 (0.012)
pooled samples	245	85.7		2.57		0.204 (0.046)
<i>Lilium tsingtauense</i>						
population average	40	56.5	1.68	1.81	0.090 (0.009)	0.113 (0.007)
pooled samples	443	78.6		2.57		0.114 (0.037)

of both species are of large size, we collected samples (one leaf per individual) as random as possible, from well-separated individuals. Samples were the same as those used by Chung *et al.* (2014a), in a study that was aimed to infer past history and palaeodistribution (at the Last Glacial Maximum, LGM) of the two *Lilium* species on the Korean Peninsula. Leaf samples were wrapped in damp paper towels, placed in plastic bags, returned to the laboratory, and then stored at 4 °C until protein extraction.

Enzyme electrophoresis

For enzyme extraction, leaf samples were crushed using chilled mortars and pestles by adding a crushing buffer (Mitton *et al.* 1979), and enzyme extracts were absorbed onto paper wicks (Whatman 3MM chromatography paper). We conducted electrophoresis on 13% starch gels, with two buffer systems. We used a modification (Hauffer 1985) of the system 6 of Soltis *et al.* (1983) to resolve alcohol dehydrogenase (*Adh*), diaphorase (*Dia-1*, *Dia-2*), fluorescent esterase (*Fe*), phosphoglucosomerase (*Pgi-1*, *Pgi-2*), phosphoglucumutase (*Pgm*), and triosephosphate isomerase (*Tpi-1*, *Tpi-2*). We also used the morpholine–citrate buffer system (pH 6.1) of Clayton and Tretiak (1972) to resolve isocitrate dehydrogenase (*Idh*), malate dehydrogenase (*Mdh-1*, *Mdh-2*), and 6-phosphogluconate dehydrogenase (*6Pgd-1*, *6Pgd-2*). We followed stain recipes from Soltis *et al.* (1983) except for diaphorase (Cheliak & Pitel 1984). We designated putative loci sequentially, with the most anodally migrating isozyme designated as *I*, the next 2, and so on. We also designated different alleles within each locus sequentially by alphabetical order. The observed enzyme banding patterns were consistent with their typical subunit structure and subcellular compartmentalization in diploid plants (Weeden & Wendel 1990).

Data analysis

To estimate genetic diversity and structure of the two *Lilium* species, we considered that a locus was polymorphic when two or more alleles were

observed, regardless of their frequencies. We estimated the following genetic diversity parameters using the programs POPGENE (Yeh *et al.* 1999) and FSTAT (Goudet 1995): percent polymorphic loci (%P), mean number of alleles per locus (*A*), allelic richness (AR) using a rarefaction method that compensates uneven population sample sizes (Hurlbert 1971), observed heterozygosity (H_o), and Nei's (1978) unbiased gene diversity or Hardy-Weinberg (H-W) expected heterozygosity (H_e).

To determine the degree of genetic divergence among populations of each species and between the two species, we calculated Nei's (1978) unbiased genetic identity (*I*) between all pairs of studied populations.

Results

For *L. distichum*, 12 (*Adh*, *Dia-1*, *Dia-2*, *Fe*, *Idh*, *Mdh-1*, *Mdh-2*, *6Pgd-1*, *6Pgd-2*, *Pgi-2*, *Pgm*, and *Tpi-1*) out of 14 putative loci were polymorphic across the seven populations. High levels of genetic variation were found within populations: %P = 65.3, AR = 1.90, *A* = 1.94, and H_e = 0.190 (Table 1). For *L. tsingtauense*, 11 (*Adh*, *Dia-1*, *Dia-2*, *Fe*, *Idh*, *Mdh-1*, *Mdh-2*, *6Pgd-2*, *Pgi-2*, *Pgm*, and *Tpi-1*) of the 14 surveyed loci were polymorphic across the 11 populations. All within-population genetic parameters showed moderate values: %P = 56.5, AR = 1.68, *A* = 1.81, and H_e = 0.113 (Table 1). For the two *Lilium* species, slightly higher levels of genetic variation were observed for total samples (Table 1). On average, populations of *L. distichum* harboured significantly higher levels of genetic variation than those of *L. tsingtauense* (Wilcoxon rank sum test: $p = 0.058$ for %P, $p = 0.008$ for AR, and $p = 0.001$ for H_e). Details on genetic estimates at the population level are available in Chung *et al.* (2014a: Table 1).

Except for *Fe*, allele frequencies of *L. distichum* were very similar to those of *L. tsingtauense* (Table 2). In *L. distichum*, the most common allele at the *Fe* locus was *Fe^b* (occurring at a frequency of 0.700; Table 2), which was absent from *L. tsingtauense* (that is, it can be regarded as a diagnostic allele for *L. distichum*). In *L. tsingtauense*, the most common allele at

the *Fe* locus was, instead, *Fe^e* (occurring at a frequency as high as 0.897; Table 2), absent from *L. distichum* (i.e. a diagnostic allele for *L. tsingtauense*).

Discarding the alleles *Fe^b* and *Fe^e*, all the alleles that were species-specific (five in each species) were present at very low frequencies (0.001–0.190; Table 2). Accordingly, pairwise Nei's (1978) *I* values between congeneric populations of *L. distichum* and *L. tsingtauense* were high, ranging from 0.876 (LD-4 vs. LT-4) to 0.943 (LD-2 vs. LT-7), and with a mean \pm SD of 0.919 ± 0.017 (Table 3). Pairwise Nei's (1978) *I* values between conspecific populations of *L. distichum* were high, ranging from 0.946 (LD-1 vs. LD-5) to 0.998 (LD-6 vs. LD-7), and with a mean \pm SD of 0.971 ± 0.014 (Table 3). High *I* values were also found for *L. tsingtauense*, ranging from 0.977 (LT-4 vs. LT-6) to 1.000 (LT-10 vs. LT-11), and with a mean \pm SD of 0.990 ± 0.005 (Table 3).

Discussion

Information on allele distributions and levels of genetic variation between closely related species has provided valuable insights into the identification of P–D species pairs (Gottlieb 1973, 2003, Crawford & Smith 1982, Crawford 1983, 2010, Loveless & Hamrick 1988, Pleasants & Wendel 1989, Chung *et al.* 1991, Maki *et al.* 1999, Hiramatsu *et al.* 2001, López-Pujol *et al.* 2001). In accordance with those previous studies, our predictions concerning the origin of *L. tsingtauense* as a D species from *L. distichum* are supported by genetic data.

First, *L. tsingtauense* exhibits significantly lower allozyme variability at both the population and the species levels than *L. distichum* (Table 1). Second, to some extent, the allelic composition of *L. tsingtauense* represents a subset of *L. distichum*. Across the 12 polymorphic loci, 34 alleles were detected in *L. distichum*; 28 (82%) of these were also detected in *L. tsingtauense* (Table 2). Except for *Fe^e* (allele frequency = 0.897), all the species-specific alleles of *L. tsingtauense* (*Dia-1^d*, *Dia-2^a*, *Mdh-2^e*, *Pgi-2^e*, and *Tpi-1^a*) were detected at very low frequencies (0.002–0.032; Table 2). Thus, we

could speculate that these rare alleles may have remained undetected in *L. distichum*. The differences in allele composition and frequency (e.g. the two species have different major alleles at the *Fe* locus, and five additional alleles in each species are taxon-specific) might be, instead, attributable to interspecific divergence between

Table 2. Allele frequencies of 12 and 11 polymorphic loci for *Lilium distichum* and *L. tsingtauense* in Korea, respectively. Frequencies indicating most common alleles at each locus are set in boldface, diagnostic alleles are set in italics, and species-specific alleles are underlined.

Locus	Allele	<i>L. distichum</i>	<i>L. tsingtauense</i>
<i>Adh</i>	<i>b</i>	0.986	0.995
	<i>c</i>	0.014	0.005
<i>Dia-1</i>	<i>a</i>	0.012	0.001
	<i>b</i>	0.982	0.988
	<i>c</i>	0.006	0.007
	<i>d</i>	0	<u>0.005</u>
<i>Dia-2</i>	<i>a</i>	0	<u>0.002</u>
	<i>b</i>	0.949	0.976
	<i>c</i>	0.051	0.021
<i>Fe</i>	<i>a</i>	<u>0.035</u>	0
	<i>b</i>	<i>0.700</i>	0
	<i>c</i>	0.249	0.077
	<i>d</i>	0.016	0.026
	<i>e</i>	0	<i>0.897</i>
<i>ldh</i>	<i>a</i>	0.218	0.035
	<i>b</i>	0.761	0.938
	<i>d</i>	0.020	0.027
<i>Mdh-1</i>	<i>a</i>	0.055	0.026
	<i>b</i>	0.945	0.974
<i>Mdh-2</i>	<i>b</i>	0.941	0.936
	<i>c</i>	0	<u>0.032</u>
	<i>d</i>	0.059	0.033
<i>6Pgd-1</i>	<i>a</i>	<u>0.002</u>	0
	<i>b</i>	<u>0.190</u>	0
	<i>c</i>	0.808	1
<i>6Pgd-2</i>	<i>a</i>	0.067	0.019
	<i>b</i>	0.869	0.881
	<i>c</i>	0.063	0.099
<i>Pgi-2</i>	<i>b</i>	0.163	0.067
	<i>c</i>	0.818	0.827
	<i>d</i>	0.018	0.096
<i>Pgm</i>	<i>e</i>	0	<u>0.010</u>
	<i>a</i>	<u>0.059</u>	0
	<i>b</i>	0.227	0.164
	<i>c</i>	0.665	0.694
	<i>d</i>	0.049	0.142
<i>Tpi-1</i>	<i>a</i>	0	<u>0.007</u>
	<i>b</i>	0.814	0.973
	<i>c</i>	0.171	0.020
	<i>d</i>	<u>0.014</u>	0

sister species (that is, some of the observed isozyme diversity might have arisen — or have been lost — since they diverged). Anyway, such numbers of unique alleles are not uncommon in recently-derived species (0–8 in the 11 P–D species pairs compiled by Pleasants and Wendel 1989). Third, mean interspecific genetic identity for the pair *L. distichum*–*L. tsingtauense* ($I = 0.919$; Table 3) is only slightly lower than the mean intraspecific (i.e. inter-population) identities ($I = 0.971$ for *L. distichum* and $I = 0.990$ for *L. tsingtauense*; Table 3), and almost the same as that ($I = 0.915$) averaged from 11 progenitor-derivative P–D species pairs compiled by Pleasants and Wendel (1989).

The value of genetic identity for the pair *L. distichum*–*L. tsingtauense* is much higher than those estimated for pairs of six species of *Lilium* [*L. distichum*, *L. tsingtauense*, *L. cernuum*, *L. amabile*, *L. callosum*, and *L. lancifolium* (= *L. tigrinum*)] studied by our research group ($n = 14$, $I = 0.436$ to 0.704 with a mean of 0.575 ; Table 3). Also notably, the mean I (0.919) between *L. distichum* and *L. tsingtauense* is slightly higher than that for *L. longiflorum*–*L. formosanum* ($I = 0.816$; Hiramatsu et al. 2001; see below), which is a well-known P–D species pair within East Asia.

The distribution patterns of *L. distichum* and *L. tsingtauense* also suggest that they may constitute a P–D species pair. Although *L. tsingtauense* occurs mainly in forests in low mountains and *L. distichum* is usually found at high

elevations, these species grow together in some locations at mid-elevation mountains in South Korea (Lee et al. 2011; M.Y. Chung and M.G. Chung pers. obs.). Also, *L. distichum* has a much wider geographic range than *L. tsingtauense*. In general, P species are more widely distributed than their D species (with D species occurring in small areas at the edges of widely distributed species), a common feature shown by many P–D species pairs (Crawford 2010). Despite the sympatric distribution of *L. distichum* and *L. tsingtauense* in the central ranges of the Baekdudaegan (Fig. 1), no hybridization events have been reported between them; thus, their high allele similarity should not be attributed to introgression and/or hybridization.

Our genetic results for the *L. distichum*–*L. tsingtauense* pair are similar to those for *L. longiflorum*–*L. formosanum*, the only P–D species pair detected to date within the genus (Hiramatsu et al. 2001). *Lilium longiflorum* (P species) and *L. formosanum* (D species) have been classified into the section *Leucolirion* (Comber 1949), and a series of molecular phylogenetic studies have revealed a very close relationship between them (Nishikawa et al. 1999, Hayashi & Kawano 2000, Lee et al. 2011). In addition, *L. longiflorum* has a much wider geographic range than *L. formosanum*. The former is distributed from the northernmost islands of the Ryukyu Archipelago to the northern tip of Taiwan (and to small islands in the eastern part of Taiwan), whereas the latter is restricted to main-

Table 3. Average Nei's (1978) genetic identity values, standard deviations and ranges (in parentheses) for all intraspecific (along the diagonal) and interspecific (above the diagonal) population pairwise comparisons.

	<i>L. distichum</i>	<i>L. tsingtauense</i>	<i>L. cernuum</i> ^a	<i>L. amabile</i> ^a	<i>L. callosum</i> ^b	<i>L. lancifolium</i> ^b
<i>L. distichum</i>	0.971 ± 0.014 (0.946–0.998)	0.919 ± 0.017 (0.876–0.943)	0.625 ± 0.031 (0.548–0.673)	0.704 ± 0.028 (0.669–0.774)	0.584 ± 0.016 (0.559–0.610)	0.595 ± 0.019 (0.571–0.622)
<i>L. tsingtauense</i>		0.990 ± 0.005 (0.977–1.000)	0.626 ± 0.027 (0.573–0.682)	0.671 ± 0.013 (0.637–0.698)	0.529 ± 0.009 (0.510–0.542)	0.534 ± 0.008 (0.533–0.541)
<i>L. cernuum</i> ^a			0.973 ± 0.014 (0.942–0.993)	0.505 ± 0.030 (0.454–0.564)	0.494 ± 0.022 (0.459–0.538)	0.647 ± 0.024 (0.613–0.671)
<i>L. amabile</i> ^a				0.999 ± 0.001 (0.997–1.000)	0.657 ± 0.005 (0.651–0.667)	0.440 ± 0.007 (0.433–0.451)
<i>L. callosum</i> ^b					1.000	0.436
<i>L. lancifolium</i> ^b						1.000

^a Values for eight populations from Chung et al. (2014b).

^b Value from one population from M.Y. Chung et al. (unpubl. data).

land Taiwan (Hiramatsu *et al.* 2001). As seen in many P–D species pairs, *L. formosanum* possesses a subset (85%) of *L. longiflorum* alleles for 11 polymorphic loci. In addition, *L. formosanum* exhibits lower allozyme variability than *L. longiflorum* (at the population level, %P = 48.2 vs. 31.7, $A = 1.72$ vs. 1.43, and $H_e = 0.187$ vs. 0.121; at the species level, %P = 100 vs. 76.9, $A = 3.46$ vs. 2.46, and $H_e = 0.312$ vs. 0.142; Hiramatsu *et al.* 2001). Similar to *L. tsingtauense*, the recently derived species *L. formosanum* harbors a small number of species-specific alleles (three).

However, the mean I value (0.816) between *L. formosanum* and *L. longiflorum* is lower than that between *L. distichum* and *L. tsingtauense* ($I = 0.919$), which can be attributed to several causes. These are (i) a longer divergence between *L. longiflorum* and *L. formosanum* compared with that of the *L. distichum*–*L. tsingtauense* pair, or (ii) the fewer opportunities for gene flow for the pair *L. longiflorum*–*L. formosanum* given that they occur on islands. The mean intraspecific genetic identity for *L. longiflorum* ($I = 0.850$; Hiramatsu *et al.* 2001), whose populations occur on isolated islands along over 1300 km, is even lower than that for the pair *L. distichum*–*L. tsingtauense* ($I = 0.919$).

The present study suggests that *L. tsingtauense* was probably (recently) derived from southern Korean populations of *L. distichum* when it migrated from more northern areas during one of the cold phases of the Pleistocene (Im 1992, Kong & Watts 1993). Based on the study of Guo *et al.* (2011) with ISSR (which revealed higher genetic variability for the South Korean populations compared to the Chinese populations of Shandong), one can hypothesize that *L. tsingtauense* expanded to China from Korea through the Yellow/East China Sea shelf, which was periodically exposed at each glacial period (and during the LGM, until ca. 16000 years BP; Chung 2007). This hypothesis could be tested by employing molecular phylogeography approaches and by sampling the whole region.

Acknowledgements

The authors thank Beom Jin Shim and Myeong Soon Park for laboratory assistance. This research was supported by the

Basic Science Program through the National Research Foundation of Korea (NRF-2013R1A1A3010892) funded by the Ministry of Education, Science and Technology, the Republic of Korea to M.Y.C.

References

- Arzate-Fernández A.-M., Miwa M., Shimada T., Yonekura T. & Ogawa K. 2007: *In vitro* propagation of Miyamaskashi-yuri (*Lilium maculatum* Thunb. var. *bukosanense*), an endangered plant species. — *Revista Fitotecnia Mexicana* 30: 373–379.
- Cheliak W.M. & Pitel J.P. 1984: *Technique for starch gel electrophoresis of enzyme from forest tree species*. — Information Report PI-X-42, Canadian Forestry Service, Petawawa National Forestry Institute, Chalk River, Ontario.
- Chung C.-H. 2007: Vegetation response to climate change on Jeju Island, South Korea, during the last deglaciation based on pollen record. — *Geoscience Journal* 11: 147–155.
- Chung M.G., Hamrick J.L., Jones S.B. & Derda G.S. 1991: Isozyme variation within and among populations of *Hosta* (Liliaceae) in Korea. — *Systematic Botany* 16: 667–684.
- Chung M.Y., López-Pujol J. & Chung M.G. 2014a: Comparative biogeography of the congener lilies *Lilium distichum* and *Lilium tsingtauense* in Korea. — *Flora* 209: 435–445.
- Chung M.Y., Chung M.G., López-Pujol J., Ren M.-X., Zhang Z.-Y. & Park S.J. 2014b: Were the main mountain ranges in the Korean Peninsula a glacial refugium for plants? Insights from the congeneric pair *Lilium cernuum*–*Lilium amabile*. — *Biochemical Systematics and Ecology* 53: 36–45.
- Clayton J.W. & Tretiak D.N. 1972: Amine citrate buffers for pH control in starch gel electrophoresis. — *Journal of the Fisheries Research Board of Canada* 29: 1169–1172.
- Comber H.F. 1949: A new classification of the genus *Lilium*. — *Lily Yearbook*, Royal Horticultural Society, London 13: 86–105.
- Crawford D.J. 1983: Phylogenetic and systematic inferences from electrophoretic studies. — In: Tanksley S.D. & Orton T.J. (eds.), *Isozymes in plant genetics and breeding, Part A*: 257–287. Elsevier Science Publishers, Amsterdam.
- Crawford D.J. 1985: Electrophoretic data and plant speciation. — *Systematic Botany* 10: 405–416.
- Crawford D.J. 1990: Genetics of plant isozymes. — In: Soltis D.E. & Soltis P.S. (eds.), *Isozymes in plant biology*: 146–164. Dioscorides Press, Portland.
- Crawford D.J. 2010: Progenitor–derivative species pairs and plant speciation. — *Taxon* 59: 1413–1423.
- Crawford D.J. & Smith E.B. 1982: Allozyme variation in *Coreopsis nuectenoides* and *C. nuectenoides* (Compositae), a progenitor–derivative species pair. — *Evolution* 36: 379–386.
- Gottlieb L.D. 1973: Genetic differentiation, sympatric spe-

- ciation and the origin of a diploid species of *Stephanomeria*. — *American Journal of Botany* 60: 545–553.
- Gottlieb L.D. 1984: Isozyme evidence and problem solving in plant systematics. — In: Grant W.F. (ed.), *Plant bio-systematics*: 343–357. Academic Press, Orlando.
- Gottlieb L.D. 2003: Rethinking classic examples of recent speciation in plants. — *New Phytologist* 161: 71–82.
- Goudet J. 1995: FSTAT (version 1.2): A computer program to calculate *F*-statistics. — *Journal of Heredity* 86: 485–486.
- Guo W., Jeong J., Kim Z., Wang R., Kim E. & Kim S. 2011: Genetic diversity of *Lilium tsingtauense* in China and Korea revealed by ISSR markers and morphological characters. — *Biochemical Systematics and Ecology* 39: 352–360.
- Haufler C.H. 1985: Enzyme variability and modes of evolution in *Bommeria* (Pteridaceae). — *Systematic Botany* 10: 92–104.
- Hayashi K. & Kawano S. 2000: Molecular systematics of *Lilium* and allied genera (Liliaceae): phylogenetic relationships among *Lilium* and related genera based on *rbcL* and *matK* gene sequence data. — *Plant Species Biology* 15: 73–93.
- Hiramatsu M., Ii K., Okubo H., Huang K.L. & Huang C.W. 2001: Biogeography and origin of *Lilium longiflorum* and *L. formosanum* (Liliaceae) endemic to the Ryukyu Archipelago and Taiwan as determined by allozyme diversity. — *American Journal of Botany* 88: 1230–1239.
- Hurlbert S.H. 1971: The nonconcept of species diversity: a critique and alternative parameters. — *Ecology* 52: 577–586.
- Im H.-T. 1992: Plant geographical study for the plant of Cheju. — *Korean Journal of Plant Taxonomy* 22: 219–234.
- Kong W.-S. & Watts D. 1993: *The plant geography of Korea with an emphasis on the alpine zones*. — Kluwer Academic Publishers, Dordrecht.
- Lee C.S., Kim S.-C., Yeau S.H. & Lee N.S. 2011: Major lineages of the genus *Lilium* (Liliaceae) based on nrDNA ITS sequences, with special emphasis on the Korean species. — *Journal of Plant Biology* 54: 159–171.
- Liang S. & Tamura, M.N. 2000: *Lilium*. — In: Wu Z.Y., Raven P.H. & Hong D.Y. (eds.), *Flora of China*, vol. 24 (Liliaceae): 135–149. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis.
- López-Pujol J., Bosch M., Simon J. & Blanché C. 2001: Allozyme diversity of two endemic *Petrocoptis* species: *P. montsicciana* and its close relative *P. pardoii* (Caryophyllaceae). — *Canadian Journal of Botany* 79: 1379–1389.
- Loveless M.D. & Hamrick J.L. 1988: Genetic organization and evolution history in two North American species of *Cirsium*. — *Evolution* 42: 254–265.
- McRae E.A. 1998: Lily species. — In: McRae E.A. (ed.), *Lilies: a guide for growers and collectors*: 105–204. Timber Press, Portland.
- Maki M., Morita H., Oiki S. & Takahashi H. 1999: The effect of geographic range and dichogamy on genetic variability and population genetic structure in *Tricyrtis* section *Flavae* (Liliaceae). — *American Journal of Botany* 86: 287–292.
- Miller W.B. 1993: *Lilium longiflorum*. — In: De Hertogh A. & Le Nard M. (eds.), *The physiology of flower bulbs*: 391–422. Elsevier, Amsterdam.
- Mittton J.B., Linhart Y.B., Sturgeon K.B. & Hamrick J.L. 1979: Allozyme polymorphisms detected in mature needle tissue of ponderosa pine. — *Journal of Heredity* 70: 86–89.
- Nei M. 1978: Estimation of average heterozygosity and genetic distance from a small number of individuals. — *Genetics* 89: 583–590.
- Nishikawa T., Okazaki K., Uchino T., Arakawa K. & Nagamine T. 1999: A molecular phylogeny of *Lilium* in the internal transcribed spacer region of nuclear ribosomal DNA. — *Journal of Molecular Evolution* 49: 238–249.
- Pleasant J.M. & Wendel J.F. 1989: Genetic diversity in a clonal narrow endemic, *Erythronium propullans*, and in its widespread progenitor, *Erythronium albidum*. — *American Journal of Botany* 76: 1136–1151.
- Rodger J.G., van Kleunen M. & Johnson S.D. 2010: Does specialized pollination impede plant invasions? — *International Journal of Plant Sciences* 171: 383–391.
- Soltis D.E., Haufler C.H., Darrow D.C. & Gastony G.J. 1983: Starch gel electrophoresis of ferns: a compilation of grinding buffers, gel and electrode buffers, and staining schedules. — *American Fern Journal* 73: 9–27.
- Sultana S., Lee S.-H., Bang J.-W. & Choi H.-W. 2010: Physical mapping of rRNA gene loci and inter-specific relationships in wild *Lilium* distributed in Korea. — *Journal of Plant Biology* 53: 433–443.
- Sultana S., Lim Y.P., Bang J.-W. & Choi H.-W. 2011: Internal transcribed spacer (ITS) and genetic variations in *Lilium* native to Korea. — *Horticulture, Environment, and Biotechnology* 52: 502–510.
- Weeden N.F. & Wendel J.F. 1990: Genetics of plant isozymes. — In: Soltis D.E. & Soltis P.S. (eds.), *Isozymes in plant biology*: 46–72. Dioscorides Press, Portland.
- Yang L. & Sun X. 2005: Characteristics of reproduction and breeding of *Lilium pumilum*. — *Acta Horticulturae Sinica* 32: 918–921.
- Yeh F.C., Yang R.C. & Boyle T.B.J. 1999: *POPGENE version 1.31, Microsoft Windows-based freeware for population genetic analysis*. Quick users' guide. — University of Alberta, Edmonton.