

Microlepia ravenii (Dennstaedtiaceae, Pteridophyta) a new species from Vietnam

Shann-Jye Moore & Jenn-Che Wang*

Department of Life Science, National Taiwan Normal University, Taipei, 116, Taiwan
(*corresponding author's e-mail: biofv017@ntnu.edu.tw)

Received 18 Jan. 2010, revised version received 8 Mar. 2010, accepted 10 Mar. 2010

Moore, S.-J. & Wang, J.-C. 2011: *Microlepia ravenii* (Dennstaedtiaceae, Pteridophyta) a new species from Vietnam. — Ann. Bot. Fennici 48: 284–287.

Microlepia ravenii S.-J. Moore (Dennstaedtiaceae, Pteridophyta), a new species from northern Vietnam, is described and illustrated. It is morphologically similar to *M. marginata* and *M. hookeriana* but differs in having cuneate and not auriculate bases of lower pinnae, cartilaginous and alternatively crenate and denticulate margins of pinnae, and a glabrous undersurface of pinnae.

The genus *Microlepia*, belonging in Dennstaedtiaceae (Smith *et al.* 2006), consists of approximately 45 species with pantropical and somewhat extratropical distributions; however, its center of distribution, and the majority of its taxa, are found in tropical to warm-temperate Asia (Kramer 1990). This genus is generally thought to be a close relative of *Dennstaedtia* (Holttum 1966, Pichi Sermolli 1977, Tryon & Tryon 1982) but differs from the latter in having pouch-like indusia and echinate spores. While preparing a monographic treatment of *Microlepia* at the herbarium of Missouri Botanical Garden in 2003, we found a peculiar specimen collected in northern Vietnam. After examining the types and all the specimens of similar taxa in B, BM, BR, E, G, GH, HAST, K, KYO, KUN, L, MICH, MO, NY, P, PE, TAIF, TI, TNS, UC, US, W and WU, and comparing all related references (e.g. Tardieu-Blot & Christensen 1939, Ching 1959, Holttum 1966, Tagawa & Iwatsuki 1979, Biswas 1988, 1991, Shieh 1994, Iwatsuki 1995, Fraser-Jenkins 1997), we came to the conclusion that the specimen represents a hitherto undescribed species.

***Microlepia ravenii* S.-J. Moore, sp. nova** (Fig. 1)

M. marginatae et M. hookeriana affinis sed basibus pinnarum inferiorum cuneatis et non auriculatis, marginibus pinnarum cartilagineis alternatim crenatis et serrulatis, paginis pinnarum inferioribus glabris.

TYPE: Vietnam. Cao Bang, Nguyen Binh, municipality Ca Thanh, vicinity of Ca Lu village, about 7–8 km to SE from Yen Lac village, about 38 km to NWW from Cao Bang town, 22°44'N, 105°50'E, 1350–1450 m, L. Averyanov *et al.* CBL 523 (holotype MO).

ETYMOLOGY: The epithet honours Dr. Peter Raven, the President of Missouri Botanical Garden (MO), to thank for his encouragement during the first author's short period research at MO.

Rhizome unknown (long-creeping?). Stipe 56 cm long, glabrous except for the very upper part, slightly sulcate on upper surface, chestnut-brown at base, brown-stramineous on upper part. Lamina pinnate, 37 cm long, 20 cm wide, acute at apex, obtuse at base, thickly papyraceous.




Fig. 1. Photograph of the holotype specimen of *Microlepia ravenii*. — **a:** Habit. — **b:** Portion enlargement of pinna showing margin, venation, and sori. Scale bars in mm.

Pinnae ca. 12–13 pairs, glabrous on both surfaces except on upper surface of costa, free, longest at base; basal pinnae 18 cm long, 1.8 cm wide, slightly falcate, cuneate at base, acuminate at apex, stalk ca. 2.5–3 mm long and with some dark and dirty hairs on undersurface; middle pinnae ca. 9–12 cm long, 1.4–1.6 cm wide, slightly falcate, obliquely and broadly cuneate at base; upper pinnae with very obscure auricle at acroscopic base; margins crenate, but with one or seldom two tiny teeth between rounded teeth, cartilaginous. Rachis grooved and glabrous on upper surface, moderately to densely hairy on

lower surface; hairs 0.22–0.38 mm long, slightly to moderately curved, dark and dirty-brown, multicellular, usually 4–7 cells, up to 8 cells, septate. Costa raised and glabrous on lower surface, slightly grooved and moderately to sparsely hairy on upper surface; hairs similar to those on rachis, but shorter, 0.15–0.28 mm long. Venation free, evident; vein-groups 2–4 times forked; veinlets usually ending at a distance ca. 0.2–0.5 mm to margin, slightly raised on both sides. Sori at tip of veinlets submarginal, at a distance ca. 0.8–1.2 mm to margin. Indusia pocket-like, glabrous.

Microlepia ravenii is similar to *M. marginata*, differing in having cuneate bases on both sides of the lower pinnae (vs. obtuse to acute bases acroscopically and cuneate bases basiscopically), lower pinnae without an auriculate lobe (vs. with a distinct enlarged auriculate lobe on acroscopic bases), pinna margins cartilaginous and alternatively crenate and denticulate (vs. non-cartilaginous and lobate to pinnately incised), pinna undersurfaces glabrous (vs. sparsely to very densely hairy), and indusia glabrous (vs. usually hairy). It is also similar to *M. hookeriana*, but differs in having an acute frond apex (vs. caudate), bases on both sides of lower pinnae cuneate (vs. truncate to obtuse, or acroscopic sides truncate and basiscopic sides obtuse, sometimes a little cordate), lower pinnae without auriculate lobe (vs. with a distinct triangular auriculate lobe on acroscopic bases, sometimes also on basiscopic bases), pinna margins cartilaginous and alternatively crenate and denticulate (vs. non-cartilaginous and dentate), upper surface of rachis glabrous (vs. hairy), and sori loosely arranged (vs. more closely arranged).

Microlepia ravenii is known only from the type locality in an old open mixed semideciduous and coniferous (with *Keteleeria fortunei*) primary forest. It grows along tops of a limestone ridge and on steep slopes near the ridge tops at an altitude of ca. 1350–1450 m.

SELECTED SPECIMENS EXAMINED. — *Microlepia hookeriana*.

Burma. Sumprabum, *F. Kingdom-Ward* 20581 (BM, GH). **China.** Kwangtung, *C. L. Tso* 21042 (K). Fukien, *J. L. Gressitt* 1699 (BM, G, GH, UC). Hainan, *F. A. McClure* 8509 (BM, E, G, K, MICH, NY, P, UC). Hongkong, *W. T. Tang* 29680 (G, MICH, NY, UC). Kwangsi, *R. C. Ching* 6556 (NY). Kwangtung, *S. K. Lau* 2396 (GH, MICH). Yunnan, *A. Henry* 10399 (B, E, K, NY, P). **India.** Arunachal Pradesh, *C. R. Fraser-Jenkins* FN66 (TAIF). Assam, *Wallich Cat. no* 2684 (BM, K). Darjeeling, *C. B. Clarke* 35072C (NY). Khasia, *C. B. Clarke* 45420A (K). **Japan.** Ryukyu, *S. Kurata & T. Nakaike* 1911 (A, K, KYO, RYU, TNS). **Malaysia.** Borneo, *J. & M. S. Clemens* 27407 (BM, GH, L, NY). **Indonesia.** Java, *W. F. Winckel* 1696B (L). Sumatra, *W. J. J. O. de Wilde and B. E. E. de Wilde-Duyfjes* 19192 (L). **Taiwan.** Bankinsing, *U. J. Faurie* 295 (BM, L, P). Nantou Hsien, *Huang, Kou & Kao* 893 (MICH, UC). Taipei Hsien, *U. J. Faurie* exsicc. n. 031 (B, GH, MICH, UC). **Thailand.** Chiangrai, *K. Iwatsuki & N. Fukuoka* T3693 (A, K, KYO, L). Loei, *T. Smitinand* 5910 (K). Nakawan Sritamarat, *M. Tagawa, K. Iwatsuki & N. Fukuoka* T4686 (A, K, KYO, L). Nakhon Si Thammarat, *C. F. van Beusekom & C. Phengkhla* 807 (B, BM, L, MICH). Phang Nga, *A. F. G. Kerr* 17210 (BM, K). Prachinburi, *E. Hennip-*

man 3933 (L). Udawn, *E. Hennipman* 3672 (L, MICH, NY). **Vietnam.** Bac-Giang, Lang Met, V.1925 *Colani s.n.* (GH, MICH, P). Indochine, *P. A. Eberhardt* 5093 (P). Kon Plong District, *D. K. Harder et al.* 4659 (UC). Tam Dao Hill Station, *H. Werff, B. Gray, N. K. Dao & C. I. Peng* 14132 (MO, UC). Tonkin, *B. Balansa* 120 (BR, G, GH, P). — *Microlepia marginata*. **China.** Anhui, *Deng & Yao* 79072 (NY). Chekiang, *E. D. Merrill* 11290 (GH, NY). Fokien (Fukien), *S. T. Dunn* 3829 (MICH, P). Guizhou, *Sino-American Guizhou Bot. Exped.* 163 (BM, MICH, NY). Hainan, *F. A. McClure* 8412 (BM, G). Hongkong, *S. Y. Hu* 8817 (A, K). Hubei, *Sino-American Exped.* 2052 (A). Hunan, *L.-B. Luo* 175 (L, MICH, NY, PE, WU). Jiangxi, *C.-E. Tan* 95543 (BM, PE). Kwangsi, *R. C. Ching* 6797 (NY). Kwangtung, *Y. Tsiang* 6125 (BM). Sichuan, *D. E. Boufford & B. Bartholomew* 24011 (BM, BR, E, G, L, MICH, NY, PE). Yunnan, *M. Kato et al.* 2718 (A, BM). **India.** Khasi Hills, *C. J. Simons s.n.* (BM). **Japan.** Hondo, *J. Ohwi & S. Okuyama* NSM. no. 195 (A, BM, BR, G, K, L, MICH, NY, TI, W). Honshu, *M. Tagawa* 7146 (BM, G, K, KYO, L, W). Kyushu, *M. Tagawa* 7868 (KYO, US). Nagasaki, VIII.1887 *O. Warburg s.n.* (B). Shikoku, *U. J. Faurie* 4608 (B, P). Yokohama, 1862 *C. Maximowicz s.n.* (B). **Laos.** Env. de Nape, 13.X.1928 *M. Delacour s.n.* (P). **Nepal.** Kathmandu Valley, *L. Fleming* 1332 (MICH). **Taiwan.** Hsinchu County, *P.-F. Lu* 7148 (TAIF). Nantou Hsien, *C. E. DeVol* 8085 (BM). Taihoku, *M. Tagawa* 282 (L, MICH, KYO). **Thailand.** Phang Nga, *A. F. G. Kerr* 17207 (BM). Klang Dong, *K. Larsen* 9348 (A). Udawn, *E. Hennipman* 3530 (B, BM, L, MICH, NY). **Vietnam.** Tonkin, *B. Balansa* 1790 (G, GH, P).

Acknowledgments

The authors thank Dr. Si He and Dr. Michele Funston for examining the holotype. We also thank the staff of the herbaria mentioned above for their kind assistance during the first author's visit.

References

- Biswas, A. 1988: The study on the fern genus *Microlepia* Presl in India. II. *Microlepia trapeziformis* (Roxb.) Kuhn Complex. — *Journal of Economic and Taxonomic Botany* 12: 313–327.
- Biswas, A. 1991: The study on the fern genus *Microlepia* Presl in India. I. *Microlepia strigosa* (Thunb.) Pr. complex. — *Journal of Economic and Taxonomic Botany* 15: 31–39.
- Ching, R. C. 1959: Pteridophyta (Ophioglossaceae-Oleandraceae). — In: Chien, S. S. & Chun, W. Y. (eds.), *Flora Reipublicae Popularis Sinicae* 2: 1–406. Typis Academicae Scientiarum Sinicae, Pekini.
- Fraser-Jenkins, C. R. 1997: *New species syndrome in Indian pteridology and the ferns of Nepal*. — International Book Distributors, Dehradun.
- Holtum, R. E. 1966: *A revised Flora of Malaya, vol. II. Ferns of Malaya*. — Government Printing Office, Sin-

- gapore.
- Iwatsuki, K. 1995: Dennstaedtiaceae. — In: Iwatsuki, K., Yamazaki, T., Boufford, D. E. & Ohba, H. (eds.), *Flora of Japan, vol. I: Pteridophyta and Gymnospermae*: 59–66. Kodansha, Tokyo.
- Kramer, K. U. 1990: Dennstaedtiaceae. — In: Kramer, K. U. & Green, P. S. (eds.), *The families and genera of vascular plants, vol. I: Pteridophytes and Gymnosperms*: 81–94. Springer-Verlag, Berlin.
- Pichi Sermolli, R. E. G. 1977: Tentamen pteridophytorum genera in taxonomicum ordinem redigendi. — *Webbia* 31: 313–512.
- Shieh, W. C. 1994: Dennstaedtiaceae. — In: Huang, T. C. (ed.), *Flora of Taiwan*, vol. 1, 2nd ed.: 150–171. Editorial Committee of the Flora of Taiwan, Taipei.
- Smith, A. R., Pryer, K. M., Schuettpelz, E., Korall, P., Schneider, H. & Wolf, P. G. 2006: A classification for extant ferns. — *Taxon* 55: 705–731.
- Tagawa, M. & Iwatsuki, K. 1979: Pteridophytes (Psilotaceae–Dennstaedtiaceae). — In: Smitinand, T. & Larsen, K. (eds.), *Flora of Thailand* 3(1): 1–128. The Tistr Press, Bangkok.
- Tardieu-Blot, M. & Christensen, C. 1939: Dennstaedtioidees. — In: Lecomte, H. (ed.), *Flore generale de l'Indo-Chine* 7–2(6): 90–102. Massion et C^e, Paris.
- Tryon, R. M. & Tryon, A. F. 1982: *Ferns and allied plants, with special reference to tropical America*. — Springer-Verlag, New York-Heidelberg-Berlin.