

# Liverworts and hornworts of Taiwan: an updated checklist and floristic accounts

Jian Wang<sup>1</sup>, Ming-Jou Lai<sup>2†</sup> & Rui-Liang Zhu<sup>1,\*</sup>

<sup>1</sup> Department of Biology, School of Life Science, East China Normal University, 3663 Zhong Shan North Road, Shanghai 200062, China (\*corresponding author's e-mail: rzhu@bio.ecnu.edu.cn)

<sup>2</sup> Department of Landscape Architecture, P.O. Box 834, Tunghai University, Taichung, Taiwan 407, China

Received 2 Mar. 2010, revised version received 21 June 2010, accepted 22 June 2010

Wang, J., Lai, M. J. & Zhu, R. L. 2011: Liverworts and hornworts of Taiwan: an updated checklist and floristic accounts. — *Ann. Bot. Fennici* 48: 369–395.

An updated checklist of the liverworts (Marchantiophyta) and hornworts (Anthocerotophyta) of Taiwan is presented. Based on published records, the present checklist includes 512 species of liverworts belonging to 116 genera in 52 families and 19 species of hornworts belonging to six genera in three families. Lejeuneaceae with 121 spp. in 24 genera is the largest family, containing ca. 23% of the total number of species. Genera with more than ten species are *Frullania* (49 spp.), *Cololejeunea* (41 spp.), *Plagiochila* (36 spp.), *Radula* (26 spp.), *Lejeunea* (25 spp.), *Bazzania* (18 spp.), *Scapania* (18 spp.), *Solenostoma* (17 spp.), *Riccardia* (15 spp.), *Porella* (12 spp.) and *Heteroscyphus* (11 spp.). Twenty-five taxa are hitherto known only from Taiwan. Forty-seven formerly recorded species are dubious and excluded from this checklist. *Chiloscyphus engelii* R.L. Zhu & M.J. Lai *nom. nov.* as well as *Folioceros verruculosus* (J. Haseg.) R.L. Zhu & M.J. Lai *comb. nova*, *Lejeunea barbata* (Herzog) R.L. Zhu & M.J. Lai, *comb. nova* and *Lejeunea luzonensis* (Steph.) R.L. Zhu & M.J. Lai, *comb. nova* are proposed. *Cololejeunea peraffinis* fo. *corticola* (Benedix ex Herzog) S.H. Lin is treated as a new synonym of *Cololejeunea peraffinis* (Schiffn.) Schiffn.

## Introduction

The island of Taiwan is 448 km long, with a maximum width of 144 km, and a total area of 36 000 km<sup>2</sup>. Plains below 100 m a.s.l. occupy 31.3% of the total surface, areas between 100–1000 m occupy 37.2%, and 31.5% of the area is above 1000 m. The highest mountain peak, Mt. Yushan, reaches 3952 m.

Bisected by the Tropic of Cancer and situated between two branches of the warm Kuroshio (or Kuroshio) sea current, Taiwan has a distinct

oceanic subtropical–tropical climate tempered by winds from the sea and by frequent rains and typhoons. Humidity is normally high throughout the year, and the mean annual rainfall is about 2000 mm. With the exception of a few mountain areas, the mean annual temperature is ca. 22 °C, with the lowest temperatures ranging from 12 °C to 17 °C. The abundance of rain, the high temperatures and varied topography make Taiwan rich in vegetation types. The vascular flora consists of more than 4000 species, of which ca. 26% are endemic (Lai 2003). It is estimated that

52% of the land is covered by forests, of which the Tertiary relicts/conifers *Taiwania cryptomerioides* and *Chamaecyparis formosensis* are world-famous.

The studies on liverworts and hornworts of Taiwan can be traced back to the first report on *Marchantia cuneiloba* (= *M. emarginata* subsp. *tosana*) by Stephani (1897) based on A. Henry's collection from the south of Taiwan during the period of 1892–1895 (Kuo & Chiang 1988, Lin 2000). Early collections of Taiwan hepatics were made by R. Oldham, A. Henry, K. Miyake and U. Faurie, and their specimens, studied by Stephani (1900–1924), were described as 24 taxa, including 19 species unknown until then. From then on, the major fieldwork and research on Taiwan liverworts and hornworts were conducted by Japanese bryologists. The main contributions include Hattori (1941, 1942a, 1942b, 1942c, 1943, 1944, 1951b, 1980a, 1980b), Horikawa (1932, 1934a, 1934b, 1935, 1936, 1939, 1951), Inoue (1967, 1978, 1982, 1985), Mizutani (1961, 1964, 1967, 1968), Okamura (1916), Yamada (1979), Yamada and Inoue (1988), and Yamada and Lai (1979). Since the 1950s, many studies on Taiwan liverworts and hornworts have been published by European and Chinese taxonomists and the results were published mainly by Herzog and Noguchi (1955), Huang and Lin (2009), Huang et al. (2009), Inoue and Yang (1966), Kuo and Chiang (1988), Lai (1990, 1995, 2002), Lai and Wang-Yang (1976), Lin (1981, 1990, 2000), Lin and Yang (1992), Schuster (1998), Váña and Inoue (1983), Wu and Yamada (1996, 1997, 2000), Wu et al. (2002), Yang (1960a, 1960b, 1963, 1965, 1966, 2009), Yang and Lin (2008a, 2008b, 2008c, 2009a, 2009b), Zhu and So (1998a, 2000a, 2000b, 2001) and Zuo and Cao (2007).

Four checklists of the liverworts and hornworts of Taiwan have been compiled, by Lai and Wang-Yang (1976), Kuo and Chiang (1988), Piippo (1990) and Lin (2000), respectively. The first of those checklists contained 385 species in 116 genera; Kuo and Chiang (1988) listed 503 species in 124 genera; Piippo (1990) enumerated 489 species in 122 genera; and Lin (2000) presented the most recent checklist with 487 species in 121. Over the past decade, a number of nomenclatural changes and new additions have been reported for the Taiwan liverwort and horn-

wort flora due to taxonomic revisions, floristic investigations, and molecular phylogenetic studies. Therefore publishing an updated checklist is necessary.

## Floristic accounts

The present list contains 512 species of liverworts belonging to 116 genera in 52 families and 19 species of hornworts belonging to six genera in three families. Lejeuneaceae with 121 spp. in 24 genera is the most species-rich family, accounting ca. 23% of the total number of the species. Genera with numerous species are *Frullania* (49 spp.), *Cololejeunea* (41 spp.), *Plagiochila* (36 spp.), *Radula* (26 spp.), *Lejeunea* (25 spp.), *Bazzania* (18 spp.), *Scapania* (18 spp.), *Solenostoma* (17 spp.), *Riccardia* (15 spp.), *Porella* (12 spp.) and *Heteroscyphus* (11 spp.). Forty-seven formerly reported species are considered dubious and excluded from this checklist. Twenty-five taxa, *Anastrophyllum minutum* var. *acuminatum*, *A. striolatum*, *Anthoceros fusiformis* var. *taiwanensis*, *Bazzania serrulatoides*, *B. spinosa*, *Calypogeia formosana*, *Chiloscyphus breviculus*, *Folioceros verruculosus*, *Frullania formosae*, *F. hiroshii*, *F. macrophylla*, *F. sphaerolobulata*, *F. taiheizana*, *F. yuzawana*, *Lejeunea barbata*, *L. chaishanensis*, *L. laii*, *Marchantia formosana*, *Plagiochila multipinnula*, *P. taiwanensis*, *Riccardia formosensis*, *R. submultifida*, *Schistochila minor*, *Targionia formosica* and *Tuyamaella molischii* var. *taiwanensis* are known only from Taiwan.

Wu et al. (2002) analyzed the similarity of the bryophyte floras between Taiwan and the Hengduan Mountains, southwest China, and they considered that almost all species common to those areas are tropical or subtropical. We employed Kroeber's percentage similarity (van Balgooy 1971) to evaluate the floristic affinities of liverworts and hornworts between different regions: Taiwan, Japan (Yamada & Iwatsuki 2006), the Philippines (Tan & Engel 1986), Fujian (Zhang et al. 2011) and Hainan Island (Shi 2008). The Kroeber's percentage of generic similarity of Taiwan and the Philippines is 73.5%, which is lower than the 75.3% with Hainan Island, the 80.0% with Fujian, and the 84.7% with Japan.

This conclusion reveals that the affinity of the liverwort and hornwort flora of Taiwan with Japan is closer than that with the Philippines, as concluded from the moss flora by Higuchi and Lin (2004). The lower Kroeber's percentage of generic similarity of Taiwan and Hainan plus the low number of species and genera of Hainan indicate that more detailed investigations on the liverwort and hornwort flora of Hainan are needed. The relatively high similarity of the liverwort and hornwort flora between Taiwan and Fujian might probably be explained by the fact that Taiwan was connected with Fujian in the mid-Pleistocene (Lai 2003).

## Updated checklist of the liverworts and hornworts of Taiwan

The checklist was compiled according to the records in the published literature. The records given in Piippo (1990) are not repeated here. Species names with references are arranged in alphabetical order as in our earlier paper (Lai *et al.* 2008). The nomenclature mostly follows Crandall-Stotler *et al.* (2009), Frey *et al.* (2009), and recent revisions.

### Liverworts (Marchantiophyta)

- Acrobolbus ciliatus* (Mitt.) Schiffn. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Wu & Wang (2001), Gao & Lai (2003), Gao & Wu (2008).
- Acrolejeunea pusilla* (Steph.) Grolle & Gradst. — Piippo (1990), So & Zhu (1996b), He (1997), Lin (2000), Yang (2009).
- Acromastigum divaricatum* (Nees) A. Evans — Piippo (1990), Lin (2000), Gao (2003), Gao & Lai (2003).
- Anastrepta orcadensis* (Hook.) Schiffn. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008).
- Anastrophyllum bidens* (Reinw. *et al.*) Steph. — Piippo (1990), Lin (2000), Gao & Wu (2008), Hsu & Huang (2010).
- A. michauxii* (F. Weber) H. Buch — Lin (2000), Gao & Wu (2008).
- A. minutum* (Schreb.) R.M. Schust. var. *acuminatum* (Horik.) T. Cao & J. Sun — Piippo (1990 as *A. acuminatum*), Lin (2000 as *A. acuminatum*), Gao & Wu (2008).
- A. minutum* (Schreb.) R.M. Schust. var. *minutum* — Piippo (1990), Lin (2000), Gao & Lai (2003), Gao & Wu (2008).
- A. striolatum* (Horik.) N. Kitag. — Piippo (1990), Lin (2000), Gao & Wu (2008).
- Aneura maxima* (Schiffn.) Steph. — Piippo (1990), Lin (2000).
- A. pinguis* (L.) Dumort. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- Apomarsupella revoluta* (Nees) R.M. Schust. — Piippo (1990 as *Marsupella revoluta*), Bapna & Kachroo (2000b as *M. revoluta*), Gao & Cao (2000 as *M. revoluta*), Lin (2000 as *M. revoluta*), Gao & Wu (2008).
- A. rubida* (Mitt.) R.M. Schust. — Piippo (1990 as *Marsupella rubida*), Bapna & Kachroo (2000b as *M. rubida*), Lin (2000 as *M. rubida*), Gao & Wu (2008).
- Apotreubia nana* (S. Hatt. & Inoue) S. Hatt. & Mizut. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Gao & Lai (2003).
- Archilejeunea planiuscula* (Mitt.) Steph. — Stephani (1912 as *Brachiolejeunea miyakeana*), Horikawa (1934a as *B. miyakeana*).
- Asterella leptophylla* (Mont.) Grolle — Piippo (1990 as *A. yoshinagana*), Lin (2000 as *A. yoshinagana*), Wu & Wang (2001 as *A. yoshinagana*), Gao & Lai (2003 as *A. yoshinagana*), Long (2006).
- A. wallichiana* (Lehm. & Lindenb.) Grolle — Piippo (1990 as *A. monospiris*), Lin (2000 as *A. monospiris*), Long (2006).
- Barbilophozia attenuata* (Mart.) Loeske — Piippo (1990), Lin (2000), Gao & Lai (2003).
- B. barbata* (Schreb.) Loeske — Piippo (1990), Lin (2000), Gao & Lai (2003).
- B. hatcheri* (A. Evans) Loeske — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Wu & Wang (2001).
- Bazzania albifolia* Horik. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003).
- B. angustifolia* Horik. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003), Hsu & Huang (2010).
- B. bidentula* (Steph.) Steph. ex Yasuda — Piippo (1990), Bapna & Kachroo (2000a), Lin (2000).
- B. conophylla* (Sande Lac.) Schiffn. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003).
- B. fauriana* (Steph.) S. Hatt. — Piippo (1990), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- B. imbricata* (Mitt.) S. Hatt. — Long & Grolle (1990), Piippo (1990), Lin (2000), Hsu & Huang (2010).
- B. japonica* (Sande Lac.) Lindb. — Piippo (1990), Lin (2000), Gao & Lai (2003).
- B. magna* Horik. — Piippo (1990), Lin (2000), Gao (2003).
- B. ovistipula* (Steph.) Abeyw. — Long & Grolle (1990), Piippo (1990), Lin (2000), Hsu & Huang (2010).
- B. praerupta* (Reinw. *et al.*) Trevis. — Piippo (1990), Gao & Cao (2000), Lin (2000), Hsu & Huang (2010).
- B. revoluta* (Steph.) N. Kitag. — Piippo (1990), Lin (2000), Gao (2003), Gao & Lai (2003).
- B. serrulatoidea* Horik. — Piippo (1990), Lin (2000), Gao (2003), Gao & Lai (2003).
- B. sikkimensis* (Steph.) Herzog — Piippo (1990), Lai (1995), Lin (2000).
- B. spinosa* S. Okamura — Piippo (1990), Lin (2000).
- B. tricrenata* (Wahlenb.) Lindb. — Piippo (1990), Bapna &

- Kachroo (2000a), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003).
- B. tridens* (Reinw. et al.) Trevis. var. *tridens* — Piippo (1990), Piippo (1990 as *B. oshimensis*), Lai (1995), Bapna & Kachroo (2000a), Lin (2000 as *B. oshimensis*), Gao & Lai (2003), Hsu & Huang (2010).
- B. tridens* (Reinw. et al.) Trevis. var. *assamica* (Steph.) Pócs — Lai (1990), Piippo (1990), Lin (2000).
- B. trilobata* (L.) Gray — Piippo (1990), Lin (2000).
- B. vittata* (Gottsche) Trevis. — Piippo (1990), Lin (2000), Lai (2002), Gao (2003), Gao & Lai (2003), Hsu & Huang (2010).
- Blasia pusilla* L. — Lai (1990), Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Gao & Lai (2003).
- Blepharostoma minus* Horik. — Gao & Cao (2000).
- B. trichophyllum* (L.) Dumort. — Lai (1990), Piippo (1990), Bapna & Kachroo (2000a), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Zhang & Wu (2006), Hsu & Huang (2010).
- Calycularia crispula* Mitt. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- Calypogeia aeruginosa* Mitt. — Yang & Lin (2009a).
- C. arguta* Nees & Mont. — Lai (1990), Piippo (1990), So & Zhu (1996b), Bapna & Kachroo (2000a), Lin (2000), Hsu & Huang (2010).
- C. azurea* Stotler & Crotz — Piippo (1990 as *C. trichomanis*), Lin (2000 as *C. trichomanis*).
- C. fissa* (L.) Raddi — Piippo (1990), Lin (2000), Gao (2003).
- C. formosana* Horik. — Piippo (1990), Lin (2000).
- C. neesiana* (C. Massal. & Carestia) Müll. Frib. — Piippo (1990), Lin (2000).
- C. tosana* (Steph.) Steph. — Lai (1990), Piippo (1990), Lin (2000), Zhu & So (2001), Zhang & Wu (2006).
- Caudalejeunea reniloba* (Gottsche) Steph. — Zhu & Lai (2003 as *C. recurvistipula*).
- Cephalozia bicuspidata* (L.) Dumort. — Piippo (1990).
- C. catenulata* (Huebener) Lindb. subsp. *nipponica* (S. Hatt.) Inoue — Piippo (1990), Lin (2000).
- C. gollanii* Steph. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Long (2005).
- C. hamatiloba* Steph. — Piippo (1990 as *C. otaruensis*), Váña (1988, 1993), Bapna & Kachroo (2000a), Lin (2000 as *C. otaruensis*), Long (2005), Hsu & Huang (2010 as *C. otaruensis*).
- Cephalozia kiaeri* (Austin) Douin — Piippo et al. (2002).
- C. microphylla* (Steph.) Douin — Huang & Lin (2009).
- Ceratolejeunea belangeriana* (Gottsche) Steph. — Piippo (1990), He (1997), Lin (2000), Zhu et al. (2005).
- Cheilolejeunea ceylanica* (Gottsche) R.M. Schust. & Kachroo — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Zhu et al. (2002), Yang (2009).
- C. chenii* R.L. Zhu & M.L. So — Zhu et al. (1999), Lin (2000), Zhu & So (2001), Zhu et al. (2002), Yang (2009).
- C. eximia* (Ast & Tixier) R.L. Zhu & M.L. So — Zhu & So (2001), Zhu et al. (2002).
- C. falsinervis* (Sande Lac.) Kachroo & R.M. Schust. — Yang & Lin (2008a), Yang (2009).
- C. intertexta* (Lindenb.) Steph. — Piippo (1990), Lin (1990), He (1997), Gao & Cao (2000), Lin (2000), Zhu et al. (2002), Gao & Lai (2003), Yang (2009).
- C. obtusilobula* (S. Hatt.) S. Hatt. — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Zhu et al. (2002), Yang (2009).
- C. kitagawae* W. Ye & R.L. Zhu — Piippo (1990 as *Leucolejeunea paroica*), Zhu & So (1999b as *L. paroica*), Lin (2000 as *L. paroica*), Yang (2009 as *L. paroica*).
- C. trapezia* (Nees) Kachroo & R.M. Schust. — Piippo (1990 as *C. imbricata* and *C. longiloba*), Zhu & Hu (1993 as *C. imbricata*), So & Zhu (1996a as *C. imbricata*), He (1997 as *C. longiloba*), Bapna & Kachroo (2000b as *C. imbricata*), Gao & Cao (2000 as *C. longiloba*), Lin (2000 as *Lejeunea imbricata* and *C. imbricata*), Zhu & So (2001 as *C. imbricata* and *C. longiloba*), Gao & Lai (2003 as *C. longiloba*), Yang (2009 as *C. imbricata*), Hsu & Huang (2010 as *C. imbricata*).
- C. trifaria* (Reinw. et al.) Mizut. — Piippo (1990), He (1997), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Zhu et al. (2002), Gao & Lai (2003), Yang (2009).
- C. turgida* (Mitt.) W. Ye & R.L. Zhu — Grolle & Piippo (1990 as *Leucolejeunea turgida*), Wu & Yamada (1997 as *L. turgida*), Zhu & So (1999b as *L. turgida*), Gao & Lai (2003 as *L. turgida*), Zhu & Long (2003 as *L. turgida*), Yang (2009 as *L. turgida*).
- C. xanthocarpa* (Lehm. & Lindenb.) Malombe — Piippo (1990 as *Leucolejeunea xanthocarpa*), He (1997 as *L. xanthocarpa*), Zhu & So (1999b as *L. xanthocarpa*), Bapna & Kachroo (2000b as *L. xanthocarpa*), Lin (2000 as *L. xanthocarpa*), Zhu & So (2001 as *L. xanthocarpa*), Yang (2009 as *L. xanthocarpa*).
- Chiloscyphus breviculus* B.Y. Yang & W.C. Lee — Piippo (1990), Lin (2000).
- C. ciliolatus* (Nees) J.J. Engel & R.M. Schust. — Piippo (1990), Lin (2000), Gao & Wu (2008).
- C. costatus* (Nees) J.J. Engel & R.M. Schust. — Piippo (1990), Piippo (1996), Lin (2000), Gao & Wu (2008).
- C. engelii* R.L. Zhu & M.J. Lai, *nom. nov.* Basionym: *Lophocolea javanica* Schiffn., Denkschr. Kais. Akad. Wiss. Wien Math.-Nat. Clas. 70: 195. 1900. (non *Chiloscyphus javanicus* Steph., Sp. Hepat. 6: 308. 1922). *Chiloscyphus schiffneri* J.J. Engel & R.M. Schust., Nova Hedwigia 39: 422. 1985[1984], *nom. illeg.* (non *Chiloscyphus schiffneri* S.W. Arnell, Österr. Akad. Wiss., Math.-Naturwiss. Kl. 111: 71. 1964). — Gao & Wu (2008 as *Chiloscyphus schiffneri*).
- C. polyanthos* (L.) Corda var. *polyanthos* — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- C. polyanthos* (L.) Corda var. *fragilis* (Roth) K. Müll. — Piippo (1990), Lin (2000).
- C. sikkimensis* (Steph.) J.J. Engel & R.M. Schust. — Kitagawa (1979 as *Lophocolea sikkimensis*), Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Gao et al. (2004 as *Heteroscyphus acutangulus*), Gao & Wu (2008 as *H. acutangulus*), Wu & Gao (2008).
- Cololejeunea aequabilis* (Sande Lac.) Schiffn. — Zhu et al. (1999 as *C. yulensis*), Gao & Cao (2000 as *C. yulensis*), Zhu & So (2001 as *C. yulensis*).
- C. angustiflora* (Steph.) Mizut. — Zhu & So (2000b as *C. mackeeana*), Zhu & So (2001 as *C. mackeeana*), Zhu &



- So (2002), Yang (2009 as *C. mackeeana*).
- C. appressa* (A. Evans) Benedix — Piippo (1990), Zhu (1995), He (1997), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Zhu & Lai (2003), Yang (2009).
- C. ceratilibula* (P.C. Chen) R.M. Schust. — Piippo (1990 as *C. reinecheana*), He (1997 as *C. reinecheana*), Zhu & So (1998b), Bapna & Kachroo (2000b as *C. formosana*), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Yang (2009).
- C. denticulata* (Horik.) S. Hatt. — Piippo (1990), Zhu & Hu (1993), He (1997), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009), Hsu & Huang (2010).
- C. diaphana* A. Evans — Piippo (1990 as *Aphanolejeunea truncatifolia*), He (1997 as *A. truncatifolia*), Bapna & Kachroo (2000b as *C. truncatifolia*), Gao & Cao (2000 as *A. truncatifolia*), Lin (2000 as *A. truncatifolia*), Zhu & So (2001 as *A. truncatifolia*), Zhu & Long (2003 as *A. truncatifolia*), Yang (2009 as *A. truncatifolia*).
- C. dozyana* (Sandé Lac.) Schiffn. — Zhu & So (1998a), Zhu & So (1999a), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Zhu & Long (2003), Yang (2009).
- C. falcata* (Horik.) Benedix — Piippo (1990), Lin (2000), Zhu & So (2001), Yang (2009).
- C. floccosa* (Lehm. & Lindenb.) Schiffn. — Piippo (1990), He (1997), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. gottschei* (Steph.) Pandé *et al.* — Zhu & Lai (2003).
- C. grossepapillosa* (Horik.) N. Kitag. — Piippo (1990 as *Aphanolejeunea grossepapillosa*), He (1997 as *A. grossepapillosa*), Gao & Cao (2000 as *A. grossepapillosa*), Lin (2000 as *A. grossepapillosa*), Zhu & So (2001 as *A. grossepapillosa*), Yang (2009 as *A. grossepapillosa*).
- C. hainanensis* R.L. Zhu — Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. haskariana* (Lehm.) Schiffn. — Piippo (1990), Zhu & Hu (1993), He (1997), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. inflata* Steph. — Piippo (1990 as *C. oshimensis*), Zhu & Hu (1993 as *C. oshimensis*), He (1997 as *C. oshimensis*), Gao & Cao (2000), Lin (2000 as *C. oshimensis*), Zhu & So (2001), Gao & Lai (2003), Zhu & Lai (2003), Yang (2009).
- C. lanciloba* Steph. — Piippo (1990), Zhu & Hu (1993), He (1997), Lin (2000), Gao & Lai (2003), Zhu & So (2001), Zhu & Lai (2003), Yang (2009).
- C. latilobula* (Herzog) Tixier — Piippo (1990), Lin & Yang (1992), Zhu & Hu (1993), He (1997), Lin (2000), Yang (2009).
- C. longifolia* (Mitt.) Benedix *ex* Mizut. — Piippo (1990), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Zhu & Long (2003), Yang (2009).
- C. macounii* (Spruce *ex* Underw.) A. Evans — Piippo (1990), Zhu (1995), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- C. magnilobula* (Horik.) S. Hatt. — Piippo (1990), Zhu (1995), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- C. magnistyla* (Horik.) Mizut. — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- C. minutissima* (Sm.) Schiffn. — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- C. obliqua* (Nees & Mont.) Schiffn. — Piippo (1990 as *C. scabriflora*), He (1997 as *C. scabriflora*), Lin (2000 as *C. scabriflora*), Zhu & So (2001), Yang (2009).
- C. ocellata* (Horik.) Benedix — Piippo (1990), Zhu & Hu (1993), Zhu (1995), He (1997), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. ocelloides* (Horik.) Mizut. — Piippo (1990), Zhu & Hu (1993), Zhu (1995), He (1997), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. peraffinis* (Schiffn.) Schiffn. — Piippo (1990), Piippo (1990 as *Taeniolejeunea peraffinis* fo. *corticola*, 1990 as *C. magnipapillosa*), Zhu & Hu (1993 as *C. magnipapillosa*), He (1997 as *C. magnipapillosa*), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000 as *C. magnipapillosa*), Lin (2000 as *C. peraffinis* fo. *corticola*), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. planissima* (Mitt.) Abeyw. — Piippo (1990 as *C. latilobula*), Zhu & Hu (1993), He (1997), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. pseudofloccosa* (Horik.) Benedix — Piippo (1990), Zhu & Hu (1993), Zhu (1995), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Yang (2009).
- C. raduliloba* Steph. — Piippo (1990), Zhu (1995), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- C. rotundilobula* (P.C. Wu & P.J. Lin) Piippo — Zhu & So (2000b), Zhu & So (2001), Yang (2009).
- C. schmidtii* Steph. — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- C. schwabei* Herzog — Piippo (1990), Lin & Yang (1992), Zhu (1995), He (1997), Zhu & So (2001), Yang (2009).
- C. shikokiana* (Horik.) S. Hatt. — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- C. sigmoidea* Ast & Tixier — Zhu & So (1998a), Lin (2000), Zhu & So (2001), Yang (2009).
- C. sintenisii* (Steph.) Pócs — Zhu & So (2001 as *Aphanolejeunea angustiloba*), Yang (2009 as *A. angustiloba*).
- C. spinosa* (Horik.) Pandé & Misra — Piippo (1990), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Zhu & Lai (2003), Yang (2009).
- C. stylosa* (Steph.) A. Evans — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- C. subocelloides* Mizut. — Zhu & So (2001).
- C. tenella* Benedix — Zhu & So (1999a), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- C. trichomanis* (Gottsche) Steph. — Piippo (1990 as *C. goebelii*), He (1997 as *C. goebelii*), Bapna & Kachroo (2000b as *C. goebelii*), Gao & Cao (2000 as *C. goebelii*), Lin (2000 as *C. goebelii*), Zhu & So (2000c as *C. goebelii*), Zhu & So (2001 as *C. goebelii*), Gao & Lai (2003 as *C. goebelii*), Zhu & Lai (2003 as *C. goebelii*), Zhang & Wu (2006 as *C. goebelii*), Yang (2009 as *C. goebelii*), Hsu & Huang (2010 as *C. goebelii*).
- C. verrucosa* Steph. — Piippo (1990), He (1997), Lin (2000),

- Zhu & So (2001), Yang (2009).
- C. yakusimensis* (S. Hatt.) Mizut. — Zhu & Lai (2003).
- Colura acroloba* (Mont. ex Steph.) Ast — Zhu & So (2001), Yang (2009).
- C. inuii* Horik. — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- C. tenuicornis* (A. Evans) Steph. — Piippo (1990), Zhu & Hu (1993), He (1997), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009), Hsu & Huang (2010).
- Conocephalum conicum* (L.) Dumort. — Li (1990), Piippo (1990), Lin (2000), Hsu & Huang (2010).
- C. japonicum* (Thunb.) Grolle — Lai (1990), Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- Cryptomitrium tenerum* (Hook. ex Kunth) Austin ex Underw. — Piippo (1990 as *Mannia subpilosa*), Wei & Wu (1991 as *M. subpilosa*), Lin (2000 as *M. subpilosa*), Katagiri & Deguchi (2011).
- Cyathodium smaragdinum* Schiffn. ex Keissl. — Piippo (1990), Lin (2000), Gao (2003).
- Cylindrocolea recurvifolia* (Steph.) Inoue — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Huang & Lin (2009).
- Denotarisia linguifolia* (De Not.) Grolle — Piippo (1990), Lin (2000), Lin (2000 as *Jamesoniella pulchra*), Gao & Wu (2008).
- Diplasiolejeunea cavifolia* Steph. — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- Diplophyllum albicans* (L.) Dumort. — Piippo (1990), Lin (2000), Hsu & Huang (2010).
- D. andrewsii* A. Evans — Piippo (1990), Lin (2000).
- D. obtusifolium* (Hook.) Dumort. — Piippo (1990), Wu & Wang (2001).
- D. serrulatum* (Müll. Frib.) Steph. — Piippo (1990), Lin (2000), Gao & Wu (2008).
- D. taxifolium* (Wahlenb.) Dumort. — Piippo (1990), Lin (2000), Gao & Lai (2003).
- Drepanolejeunea angustifolia* (Mitt.) Grolle — Piippo (1990), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Zhu & Long (2003), Yang (2009).
- D. commutata* Grolle & R.L. Zhu — Zhu & So (2001), Yang (2009).
- D. dactylophora* (Nees et al.) Schiffn. — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- D. erecta* (Steph.) Mizut. — Piippo (1990), Zhu & Hu (1993), He (1997), Lin (2000), Zhu & So (2001), Zhu & Long (2003), Yang (2009).
- D. foliicola* Horik. — Piippo (1990 as *Leptolejeunea yangii*), Zhu & Hu (1993 as *L. yangii*), He (1997 as *L. yangii*), Bapna & Kachroo (2000b as *Rhaphidolejeunea foliicola*), Gao & Cao (2000 as *R. foliicola*), Grolle & Zhu (2000), Lin (2000 as *R. foliicola*), Zhu & So (2001), Zhu & Lai (2003), Yang (2009), Hsu & Huang (2010 as *L. yangii*).
- D. pentadactyla* (Mont.) Steph. — Piippo (1990), He (1997), Gao & Cao (2000), Lin (2000), Wu & Wang (2001), Zhu & So (2001), Yang (2009).
- D. ternatensis* (Gottsche) Schiffn. — Piippo (1990), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Yang (2009).
- D. vesiculosa* (Mitt.) Steph. — Piippo (1990), Lin et al. (1992), He (1997), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- D. yunnanensis* (P.C. Chen) Grolle & R.L. Zhu — Piippo (1990 as *Rhaphidolejeunea yunnanensis*), Zhu & So (2001), Gao & Lai (2003 as *R. yunnanensis*), Yang (2009).
- Dumortiera hirsuta* (Sw.) Nees — Lai (1990), Piippo (1990), Lin (2000), Hsu & Huang (2010).
- Eocalypogeia quelpaertensis* (S. Hatt. & Inoue) R.M. Schust. — Lin (2000 as *Metacalypogeia quelpaertensis*).
- Fossombronia caledonica* Steph. — Huang et al. (2009).
- F. japonica* Schiffn. — Piippo (1990 as *F. australi-nipponica*), Lin 2000 as *F. australi-nipponica*), Kraysky et al. (2005), Huang et al. (2009).
- F. pusilla* (L.) Nees — Piippo (1990), Lin (2000), Gao & Lai (2003), Kraysky et al. (2005), Huang et al. (2009).
- Frullania acutiloba* Mitt. — Lin & Chen (1997), Lin (2000).
- F. amamiensis* Kamim. — Piippo (1990 as *F. caduca*), Lin (2000 as *F. caduca*).
- F. amplificaria* Steph. — Piippo (1990), Lin (2000), Hsu & Huang (2010).
- F. aoshimensis* Horik. — Piippo (1990), Lin (2000), Zhu & So (2001).
- F. arecae* (Spreng.) Gottsche — Piippo (1990), Yuzawa (1991), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- F. berthoumieuui* Steph. — Piippo (1990).
- F. bonincola* S. Hatt. — Lin & Chen (1997), Lin (2000).
- F. davurica* Hampe — Piippo (1990), Gao & Cao (2000), Lin & Chen (1997), Lin (2000), Gao & Lai (2003), Ahonen (2004), Hsu & Huang (2010 as *F. jackii* subsp. *japonica*).
- F. davurica* Hampe var. *chichibuensis* (Kamim.) S. Hatt. — Piippo (1990), Lin (2000), Hsu & Huang (2010 as *F. jackii* subsp. *japonica* var. *chichibuensis*).
- F. densiloba* Steph. ex A. Evans — Piippo (1990), Lin & Chen (1997), Lin (2000).
- F. dilatata* (L.) Dumort. — Piippo (1990).
- F. diversitexta* Steph. — Piippo (1990), Lin (2000).
- F. ericoides* (Nees) Mont. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin & Chen (1997), Lin (2000), Gao & Lai (2003).
- F. formosae* Steph. — Piippo (1990), Lin (2000).
- F. gaudichaudii* (Nees & Mont.) Nees & Mont. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000).
- F. giraldiana* C. Massal. — Lin & Chen (1997), Lin (2000).
- F. gracilis* (Reinw. et al.) Dumort. — Lin (2000).
- F. hamatiloba* Steph. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- F. hiroshii* S. Hatt. — Piippo (1990), Lin & Chen (1997), Lin (2000).
- F. hypoleuca* Nees — Piippo (1990), Lin (2000), Gao & Cao (2000), Wu & Wang (2001).
- F. inflata* Gottsche — Gao & Cao (2000).
- F. inflexa* Mitt. — Gao & Cao (2000), Lin (2000), Gao & Lai (2003).

- F. inouei* S. Hatt. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- F. jackii* Gottsche — Piippo (1990 as *F. davurica* subsp. *jackii*), Gao & Cao (2000 as *F. davurica* subsp. *jackii*).
- F. kagoshimensis* Steph. — Piippo (1990), Lin & Chen (1997), Lin (2000).
- F. macrophylla* S. Hatt. — Piippo (1990), Lin (2000).
- F. meyeniana* Lindenb. — Piippo (1990), Lin & Chen (1997), Lin (2000).
- F. moniliata* (Reinw. *et al.*) Mont. — Li (1990 as *F. tamarisci* subsp. *moniliata*), Piippo (1990), Bapna & Kachroo (2000b as *F. tamarisci* subsp. *obscura* and *F. tamarisci* subsp. *obscura* var. *balansae*), Gao & Cao (2000), Lin & Chen (1997), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Hsu & Huang (2010).
- F. monocera* (Hook. f. & Taylor) Taylor var. *depauperata* S. Hatt. — Piippo (1990), Lin (2000).
- F. monocera* (Hook. f. & Taylor) Taylor var. *monocera* — Piippo (1990), Bapna & Kachroo (2000b), Lin & Chen (1997), Lin (2000), Ahonen (2004).
- F. monocera* (Hook. f. & Taylor) Taylor var. *schiffneri* (Verd.) S. Hatt. — Piippo (1990 as *F. acutiloba* var. *schiffneri*).
- F. motoyana* Steph. — Piippo (1990), Gao & Cao (2000), Lin & Chen (1997), Lin (2000).
- F. muscicola* Steph. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin & Chen (1997), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Hsu & Huang (2010 as *F. pedicellata*).
- F. nepalensis* (Spreng.) Lehm. & Lindenb. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin & Chen (1997), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- F. orientalis* Sande Lac. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- F. osumiensis* (S. Hatt.) S. Hatt. — Piippo (1990), Piippo (1990 as *F. kagoshimensis* var. *minor*), Lin & Chen (1997), Lin (2000), Lin (2000 as *F. kagoshimensis* var. *minor*).
- F. pariharui* S. Hatt. & Thaitong — Lin & Chen (1997), Lin (2000).
- F. parvistipula* Steph. — Lin & Chen (1997), Lin (2000).
- F. ramuligera* (Nees) Mont. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- F. riojaneirensis* (Raddi) Ångstr. — Piippo (1990), Piippo (1990 as *F. galeata*), Yuzawa (1991), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- F. sackawana* Steph. — Lin & Chen (1997), Lin (2000).
- F. schensiana* C. Massal. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- F. serrata* Gottsche — Piippo (1990), Gao & Cao (2000), Lin & Chen (1997), Lin (2000), Gao & Lai (2003).
- F. sphaerolobulata* S.H. Lin — Lin & Chen (1997), Lin (2000).
- F. taiheizana* Horik. — Piippo (1990), Lin & Chen (1997), Lin (2000).
- F. tamarisci* (L.) Dumort. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Ahonen (2004), Hsu & Huang (2010).
- F. tamsuina* Steph. — Piippo (1990), Lin (2000).
- F. ternatensis* Gottsche var. *non-appendiculata* S. Hatt. — Piippo (1990), Lin (2000).
- F. ternatensis* Gottsche var. *ternatensis* — Piippo (1990), Lin & Chen (1997), Lin (2000), Hsu & Huang (2010).
- F. trichodes* Mitt. — Piippo (1990), Lin & Chen (1997), Lin (2000).
- F. valida* Steph. — Lin & Chen (1997), Lin (2000).
- F. yuennanensis* Steph. — Piippo (1990), Gao & Cao (2000), Gao & Lai (2003), Lin & Chen (1997), Lin (2000).
- F. yuzawana* S. Hatt. — Piippo (1990), Lin & Chen (1997), Lin (2000).
- Gottschelia schizopleura* (Spruce) Grolle — Piippo (1990), Lin (2000), Gao & Wu (2008).
- Gymnomitrium concinatum* (Lightf.) Corda — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008).
- Haplomitrium blumii* (Nees) R.M. Schust. — Piippo (1990), Lin (2000), Gao (2003), Lai (2007).
- H. mnioides* (Lindb.) R.M. Schust. — Lai (1990), Piippo (1990), Lai (1995), Lin (2000), Zhang & Wu (2006), Lai (2007), Hsu & Huang (2010).
- Harpalejeunea filicuspis* (Steph.) Mizut. — Piippo (1990), He (1997), Lin (2000).
- Herbertus aduncus* (Dicks.) Gray — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003), Juslén (2004), Juslén (2006), Hsu & Huang (2010).
- H. dicranus* (Taylor *ex* Gottsche *et al.*) Trevis. — Piippo (1990), Piippo (1990 as *H. girdaldianus* and *H. longifissus*), Piippo (1990 as *H. sakuraii*), Hong *et al.* (1993 as *H. sakuraii* subsp. *sakuraii*), Bapna & Kachroo (2000a as *H. girdaldianus* and *H. longifolius*), Lin (2000 as *H. girdaldianus*, *H. longifissus* and *H. sakuraii*), Gao (2003), Juslén (2006), Hsu & Huang (2010), Hsu & Huang (2010 as *H. sakuraii* and *H. pinnatus*).
- H. kurzii* (Steph.) R.S. Chopra — Piippo (1990), Long & Grolle (1990), Bapna & Kachroo (2000a), Lin (2000).
- H. longispinus* J.B. Jack & Steph. — Juslén (2006).
- Heteroscyphus argutus* (Reinw. *et al.*) Schiffn. var. *argutus* — Lai (1990), Piippo (1990), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Zhang & Wu (2006), Gao & Wu (2008), Hsu & Huang (2010).
- H. argutus* (Reinw. *et al.*) Schiffn. var. *brevidens* (Schiffn.) Herzog & Nog. — Piippo (1990), Lin (2000).
- H. aselliformis* (Reinw. *et al.*) Schiffn. — Piippo (1990), Lin (2000), Gao & Wu (2008).
- H. coalitus* (Hook.) Schiffn. — Lai (1990 as *H. bescherellei*), Piippo (1990), Lai (1995 as *H. bescherellei*), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Gao & Wu (2008), Hsu & Huang (2010 as *H. bescherelli*).
- H. lophocoleoides* S. Hatt. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Wu (2008).
- H. planus* (Mitt.) Schiffn. — Lai (1990), Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008), Hsu & Huang (2010).
- H. saccogynoides* Herzog — Piippo (1990), Lin (2000), Gao & Wu (2008).
- H. splendens* (Lehm. & Lindenb.) Grolle — Piippo (1990).
- H. tener* (Steph.) Schiffn. — Piippo (1990), Bapna &

- Kachroo (2000b), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- H. tridentatus* (Sande Lac.) Grolle — Gao et al. (2004), Gao & Wu (2008).
- H. turgidus* (Schiffn.) Schiffn. — Gao et al. (2004).
- H. zollingeri* (Gottsche) Schiffn. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008).
- Isotachis japonica* Steph. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao et al. (2003 “2002”), Gao & Wu (2008), Hsu & Huang (2010).
- Jackiella javanica* Schiffn. var. *cavifolia* Schiffn. — Piippo (1990), Bapna & Kachroo (2000a), Lin (2000).
- J. javanica* Schiffn. var. *javanica* — Piippo (1990), So & Zhu (1996b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Zhang & Wu (2006).
- Jubula hutchinsiae* (Hook.) Dumort. — Ahonen (2004), Wilson et al. (2004), Katagiri et al. (2010).
- J. javanica* Steph. — Piippo (1990 as *J. hutchinsiae* subsp. *javanica*), Bapna & Kachroo (2000b), Gao & Cao (2000 as *J. hutchinsiae* subsp. *javanica*), Lin (2000 as *J. hutchinsiae* subsp. *javanica*), Gao & Lai (2003 as *J. hutchinsiae* subsp. *javanica*), Katagiri et al. (2010), Hsu & Huang (2010).
- J. japonica* Steph. — Piippo (1990), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Katagiri et al. (2010).
- Jungermannia atrovirens* Dumort. — Piippo (1990), Lin (2000).
- J. exsertifolia* Steph. — Piippo (1990 as *J. exsertifolia* subsp. *cordifolia*), Bapna & Kachroo (2000a), Lin (2000 as *J. exsertifolia* subsp. *cordifolia*).
- Kurzia abietinella* (Herzog) Grolle — Wu & Yamada (2000).
- K. borneensis* Mizut. — Wu & Yamada (2000).
- K. gonyotricha* (Sande Lac.) Grolle — Piippo (1990), Lin (2000).
- K. makinoana* (Steph.) Grolle — Piippo (1990), Bapna & Kachroo (2000a), Lin (2000), Hsu & Huang (2010).
- K. pauciflora* (Dicks.) Grolle — Piippo (1990), Bapna & Kachroo (2000a), Gao & Lai (2003), Hsu & Huang (2010 as *Lepidozia setacea*).
- Leiocolea morrisoncola* (Horik.) Furuki & Mizut. — Gao & Cao (2000 as *Lophozia morrisoncola*), Lin (2000 as *L. morrisoncola*), Bakalin (2003 as *Hattoriella morrisoncola*), Gao & Wu (2008 as *L. morrisoncola*).
- Lejeunea anisophylla* Mont. — Herzog & Noguchi (1955 as *L. subigiensis*), Piippo (1990 as *L. catanduana*), He (1997 as *L. catanduana*), Lin (2000 as *L. catanduana*), Zhu & So (2001), Yang (2009), Hsu & Huang (2010 as *L. bornensis*).
- Lejeunea apiculata* Sande Lac. — Tixier (1971 as *Stenolejeunea apiculata*), Piippo (1990 as *S. apiculata*, He (1997 as *S. apiculata*), Lin (2000 as *S. apiculata*), Zhu & So (2001 as *S. apiculata*), Yang (2009 as *P. ungulata*).
- L. aquatica* Horik. — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- Lejeunea barbata*** (Herzog) R.L. Zhu & M.J. Lai, *comb. nova*. Basionym: *Rectolejeunea barbata* Herzog in Herzog & Noguchi, J. Hattori Bot. Lab. 14: 49. 1955. The type specimen of *R. barbata* belongs to *Lejeunea* rather than *Rectolejeunea*. — Piippo (1990 as *Rectolejeunea barbata*), He (1997 as *R. barbata*), Lin (2000 as *R. barbata*).
- L. bidentula* Herzog — Yang & Lin (2008b), Yang (2009).
- L. chaishanensis* S.H. Lin — Lin & Yang (1992), Yang (2009).
- L. cocoes* Mitt. — Piippo (1990 as *L. proliferans*), He (1997 as *L. proliferans*), So & Zhu (1998), Zhu & So (1999a), Gao & Cao (2000), Lin (2000 as *L. proliferans*), Zhu & So (2001), Gao & Lai (2003), Yang (2009).
- L. compacta* (Steph.) Steph. — Piippo (1990), He (1997), Zhu & So (1999a), Lin (2000), Gao & Lai (2003), Yang (2009).
- L. curviloba* Steph. — Zhu & So (2001), Zhu & Long (2003).
- L. discreta* Lindenb. — Piippo (1990), He (1997), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Yang (2009), Hsu & Huang (2010 as *L. vaginata*).
- L. eifrigii* Mizut. — Lin (2000), Yang (2009).
- L. exilis* (Reinw. et al.) Grolle — Piippo (1990 as *Drepanolejeunea subacuta*), He (1997 as *D. subacuta*), Lin (2000 as *D. subacuta*), Zhu & Grolle (2003), Yang (2009).
- L. flava* (Sw.) Nees — Lai (1990), Piippo (1990), Zhu & Hu (1993), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2000b), Zhu & So (2001), Zhu & Lai (2003), Yang (2009), Hsu & Huang (2010).
- L. japonica* Mitt. — Piippo (1990), He (1997), Lin (2000), So (2001), Yang (2009), Hsu & Huang (2010).
- L. laii* R.L. Zhu — Piippo (1990 as *L. ramulosa*), He (1997 as *Microlejeunea ramulosa*), Lin (2000 as *L. ramulosa*), Wang & Zhu (2008), Yang (2009).
- L. leratii* (Steph.) Mizut. — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- Lejeunea luzonensis*** (Steph.) R.L. Zhu & M.J. Lai, *comb. nova*. Basionym: *Taxilejeunea luzonensis* Steph., Hedwigia 35: 134. 1896. The examination of the type material of *T. luzonensis* revealed the taxon is better placed in *Lejeunea*. — Piippo (1990 as *Taxilejeunea luzonensis*), He (1997 as *T. luzonensis*), Lin (2000 as *T. luzonensis*).
- L. obscura* Mitt. — Herzog & Noguchi (1955 as *T. subcompressiuscula*), Piippo (1990 as *T. subcompressiuscula*), He (1997 *T. subcompressiuscula*), Lin (2000 as *T. subcompressiuscula*).
- L. parva* (S. Hatt.) Mizut. — Piippo (1990), Zhu & Hu (1993), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- L. punctiformis* Taylor — Lai & Wang-Yang (1976 as *Microlejeunea punctiformis*), Kuo & Chiang (1988 as *M. punctiformis*), Bapna & Kachroo (2000b), Gao & Cao (2000), Zhu & So (2001), Gao & Lai (2003), Zhu & Lai (2003).
- L. riparia* Mitt. — Lin (2000).
- L. sordida* (Nees) Nees — Yang (2009), Yang & Lin (2009b).
- L. subacuta* Mitt. — He (1997).
- L. ulicina* (Taylor) Gottsche et al. — Lai (1990), Piippo (1990), Piippo (1990 as *L. lunulatiloba*), Zhu & Hu (1993), He (1997 as *Microlejeunea lunulatiloba* and *M. ulicina*), Bapna & Kachroo (2000b), Lin (2000), Lin (2000 as *L. lunulatiloba*), Yang (2009), Yang (2009 as *L. lunulatiloba*).
- L. wightii* Lindenb. — Piippo (1990), He (1997), Zhu & So (1999a), Bapna & Kachroo (2000b), Gao & Cao (2000),



- Lin (2000), Yang (2009).
- Lepicolea yakusimensis* (S. Hatt.) S. Hatt. — Horikawa (1934a as *L. scolopendra*), Piippo (1990), Lin (2000), Gao (2003 as *L. scolopendra*), Gao & Lai (2003), Lai (2007), Hsu & Huang (2010).
- Lepidolejeunea bidentula* (Steph.) R.M. Schust. — Piippo (1990), He (1997), Lin (2000), Zhu & So (2001), Yang (2009).
- Lepidozia fauriana* Steph. — Piippo (1990), Lai (1995), Lin (2000), Paul *et al.* (2001), Gao & Lai (2003), Hsu & Huang (2010).
- L. filamentosa* (Lehm. & Lindenb.) Gottsche *et al.* — Piippo (1990), Lin (2000), Hsu & Huang (2010).
- L. pearsonii* Spruce — Hsu & Huang (2010).
- L. reptans* (L.) Dumort. — Piippo (1990), Bapna & Kachroo (2000a), Lin (2000), Gao & Bai (2002), Gao & Lai (2003).
- L. subtransversa* Steph. — Piippo (1990), Lin (2000).
- L. trichodes* (Reinw. *et al.*) Nees — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Bai (2002), Gao (2003), Gao & Lai (2003).
- L. vitrea* Steph. — Lai (1990), Piippo (1990), Gao & Cao (2000), Lin (2000), Paul *et al.* (2001), Gao & Bai (2002), Gao (2003), Gao & Lai (2003), Hsu & Huang (2010).
- Leptolejeunea amphiophthalmia* Zwichel — Yang & Lin (2008c as *L. picta*), Yang (2009 as *L. picta*).
- L. apiculata* (Horik.) S. Hatt. — Piippo (1990 as *L. foliicola*), Gao & Cao (2000), Lin (2000 as *L. foliicola*), Zhu & So (2001), Gao & Lai (2003), Yang (2009), Hsu & Huang (2010).
- L. elliptica* (Lehm. & Lindenb.) Schiffn. — Piippo (1990), Zhu & Hu (1993), Lai (1995), He (1997), Bapna & Kachroo (2000b), Bapna & Kachroo (2000b as *L. subacuta*), Lin (2000), Zhu & So (2001), Zhu & Lai (2003), Yang (2009), Hsu & Huang (2010).
- L. epiphylla* (Mitt.) Steph. — Piippo (1990), He (1997), Zhu & So (1999a), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Gao & Cao (2000), Yang (2009).
- L. maculata* (Mitt.) Schiffn. — Piippo (1990 as *L. radiata*), He (1997 as *L. radiata*), Bapna & Kachroo (2000b as *L. schiffneri*), Lin (2000 as *L. radiata*), Zhu & So (2001), Yang (2009).
- L. truncatifolia* Steph. — Piippo (1990 as *L. aberrantia*), He (1997 as *L. aberrantia*), Lin (2000 as *L. aberrantia*), Zhu & So (2001), Yang (2009).
- Lioclaena subulata* (A. Evans) Schljakov — Piippo (1990 as *Jungermannia subulata*), Bapna & Kachroo (2000a as *J. subulata*), Lin (2000 as *J. subulata*), Váňa *et al.* (2005 as *J. subulata*).
- Lophocolea bidentata* (L.) Dumort. — Piippo (1990 as *Chiloscyphus latifolius* and *C. cuspidatus*), Gao & Cao (2000 as *C. latifolius* and *C. cuspidatus*), Lin (2000 as *C. latifolius* and *C. cuspidatus*), Gao & Lai (2003 as *C. latifolius* and *C. cuspidatus*), Gao & Wu (2008 as *C. latifolius* and *C. cuspidatus*).
- L. fragrans* (Moris. & De Not.) Gottsche *et al.* — Gao & Wu (2008 as *Chiloscyphus fragrans*).
- L. heterophylla* (Schrad.) Dumort. — Lai (1990), Piippo (1990 as *Chiloscyphus profundus*), Lin (2000 as *C. profundus*).
- L. horikawana* S. Hatt. — Piippo (1990 as *Chiloscyphus horikawanus*), Lin (2000).
- L. minor* Nees — Piippo (1990 as *Chiloscyphus minor*), Gao & Cao (2000 as *C. minor*), Lin (2000 as *C. minor*).
- L. muricata* (Lehm.) Nees — Piippo (1990 as *Chiloscyphus muricatus*), Lin (2000 as *C. muricatus*), Wu & Wang (2001), Gao & Wu (2008 as *C. muricatus*).
- Lopholejeunea eulopha* (Taylor) Schiffn. — Piippo (1990 as *L. nicobarica*), He (1997 as *L. nicobarica*), Bapna & Kachroo (2000b as *L. nicobarica*), Lin (2000 as *L. nicobarica*), Zhu & So (2001), Zhu & Gradstein (2005), Yang (2009).
- L. nigricans* (Lindenb.) Schiffn. — Piippo (1990 as *L. brunnea*, *L. javanica* and *L. muensis*), He (1997 as *L. brunnea*, *L. javanica* and *L. muensis*), Lin (2000 as *L. javanica* and *L. muensis*), Zhu & So (2000b as *L. javanica*), Zhu & So (2001), Zhu & Gradstein (2005), Yang (2009).
- L. subfusca* (Nees) Schiffn. — Piippo (1990), Lin & Yang (1992), Lai (1995), He (1997), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Zhu & Gradstein (2005), Yang (2009).
- L. zollingeri* (Steph.) Schiffn. — Piippo (1990 as *L. nipponica*), He (1997), Zhu & Gradstein (2005), Yang (2009).
- Lophozia excisa* (Dicks.) Dumort. — Piippo (1990), Lin (2000).
- L. lacerata* N. Kitag. — Piippo (1990), Lin (2000).
- L. wenzelii* (Nees) Steph. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Gao & Lai (2003), Gao & Wu (2008).
- Lunularia cruciata* (L.) Dumort. *ex* Lindb. — Piippo (1990), Lin (2000).
- Makinoa crispata* (Steph.) Miyake — Piippo (1990), Lin (2000), Hsu & Huang (2010).
- Marchantia emarginata* Reinw. *et al.* subsp. *tosana* (Steph.) Bischl. — Iwatsuki & Mizutani (1972 as *M. cuneiloba*), Lai (1990), Piippo (1990), Lai (1995 as *M. tosana*), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Zhang & Wu (2006 as *M. tosana*), Hsu & Huang (2010 as *M. cuneiloba* and *M. tosana*).
- M. formsana* Horik. — Piippo (1990), Lin (2000), Hsu & Huang (2010).
- M. paleacea* Bertol. subsp. *diptera* (Nees & Mont.) Inoue — Piippo (1990 as *M. paleacea* var. *diptera*), Gao & Cao (2000), Lin (2000 as *M. paleacea* var. *diptera*), Gao & Lai (2003 as *M. paleacea* var. *diptera*).
- M. paleacea* Bertol. subsp. *paleacea* — Piippo (1990), Lai (1995), Lin (2000).
- M. pinnata* Steph. — Piippo (1990), Lin (2000).
- M. polymorpha* L. — Li (1990), Piippo (1990), Bapna & Kachroo (2000b), Lin (2000).
- Marsupella emarginata* (Ehrh.) Dumort. — Lai (1990), Piippo (1990 as *M. emarginata* subsp. *tubulosa*), Bapna & Kachroo (2000b), Lin (2000 as *M. emarginata* subsp. *tubulosa*), Váňa *et al.* (2005).
- M. pseudofunckii* S. Hatt. — Piippo (1990), Lin (2000), Gao & Wu (2008).
- Marsupidium knightii* Mitt. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu

- (2008).
- Mastigolejeunea auriculata* (Wilson) Schiffn. — Piippo (1990), He (1997), Bapna & Kachroo (2000b as *M. humilis*), Lin (2000), Yang (2009).
- M. repleta* (Taylor) A. Evans — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- Mastigophora diclados* (Brid. ex F. Weber) Nees — Piippo (1990), Lin (2000), Gao (2003), Gao & Lai (2003), Lai (2007), Hsu & Huang (2010).
- M. woodsii* (Hook.) Nees — Piippo (1990), Lin (2000), Gao & Lai (2003), Lai (2007), Hsu & Huang (2010).
- Metacalypogeia alternifolia* (Gottsche et al.) Grolle — Piippo (1990), Bapna & Kachroo (2000a), Gao & Cao (2000), Lin (2000).
- M. cordifolia* (Steph.) Inoue — Piippo (1990), Lin (2000).
- Metalejeunea cucullata* (Reinw. et al.) Grolle — Piippo (1990 as *Lejeunea cucullata*), He (1997), Lin (2000 as *L. cucullata*), Zhu & So (2001), Yang (2009).
- Metzgeria consanguinea* Schiffn. — Lin (2000), So (2003b).
- M. foliicola* Schiffn. — So (2003b).
- M. furcata* (L.) Corda — Lai (1990), Piippo (1990 as *M. fauriana*), Bapna & Kachroo (2000b), Lin (2000 as *M. decipiens* and *M. fauriana*), Zhu & So (2001), Gao & Lai (2003), Zhang & Wu (2006), Hsu & Huang (2010).
- M. leptoneura* Spruce — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Hsu & Huang (2010 as *M. hamata*).
- M. lindbergii* Schiffn. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001).
- M. pubescens* (Schrank) Raddi — Piippo (1990 as *Apometzgeria pubescens*), Lin (2000 as *A. pubescens*), Hsu & Huang (2010 as *A. pubescens*).
- M. temperata* Kuwah. — Piippo (1990 as *M. fruticulosa*), Lin (2000 as *M. fruticulosa*).
- Mnioloma fuscum* (Lehm.) R.M. Schust. — Gao et al. (2002).
- Monosolenium tenerum* Griff. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Zhang & Wu (2006).
- Mylia taylorii* (Hook.) Gray — Piippo (1990), Lin (2000).
- M. verrucosa* Lindb. subsp. *nuda* (Inoue & B.Y. Yang) Potemkin & Kazanovsky — Piippo (1990 as *M. nuda*), Lin (2000 as *M. nuda*), Gao & Bai (2001 as *M. nuda*), Gao (2003 as *M. nuda*), Gao & Lai (2003 as *M. nuda*), Engel & Braggins (2005 as *M. nuda*), Hsu & Huang (2010 as *M. nuda*).
- M. verrucosa* Lindb. subsp. *verrucosa* — Lai (1990), Piippo (1990), Lin (2000), Gao & Bai (2001), Gao (2003), Gao & Lai (2003), Hsu & Huang (2010).
- Nardia assamica* (Mitt.) Amakawa — Lai (1990), Piippo (1990), Bapna & Kachroo (2000a), Lin (2000), Váňa et al. (2005).
- N. scalaris* Gray — Lai (1990), Piippo (1990), Lin (2000).
- Nipponolejeunea pilifera* (Steph.) S. Hatt. — Piippo (1990), He (1997), Lin (2000), Hsu & Huang (2010).
- N. subalpina* (Horik.) S. Hatt. — Piippo (1990), He (1997), Lin (2000).
- Notoscyphus lutescens* (Lehm. & Lindenb.) Mitt. — Piippo (1990), So & Zhu (1996b), Gao et al. (1999), Bapna & Kachroo (2000a as *N. parvicus*), Lin (2000), Váňa et al. (2005), Zhang & Wu (2006).
- Nowellia curvifolia* (Dicks.) Mitt. — Piippo (1990), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- Odontoschisma denudatum* (Mart.) Dumort. — Lai (1990), Piippo (1990), Lin (2000), Zhu & So (2001).
- O. grosseverrucosum* Steph. — Piippo (1990), Gao & Cao (2000), Gao & Lai (2003), Lin (2000), Yi & Gao (2001), So (2004), Hsu & Huang (2010).
- Pallavicinia ambigua* (Mitt.) Steph. — Piippo (1990), Lin (2000), Gao & Lai (2003).
- P. levieri* Schiffn. — Piippo (1990), Lin (2000).
- P. lyellii* (Hook.) Carruth. — Piippo (1990), Lai (1995), Lin (2000), Zhang & Wu (2006).
- P. subciliata* (Austin) Steph. — Lai (1990), Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010 as *P. longispina*).
- Pellia endiviifolia* (Dicks.) Dumort. — Piippo (1990), Lin (2000).
- P. epiphylla* (L.) Corda — Piippo (1990), Lin (2000).
- P. neesiana* (Gottsche) Limpr. — Piippo (1990), Lin (2000).
- Phaeolejeunea latistipula* (Schiffn.) Mizut. — Piippo (1990), He (1997), Lin (2000).
- Plagiochasma appendiculatum* Lehm. & Lindenb. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000).
- P. cordatum* Lehm. & Lindenb. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000).
- P. intermedium* Lindenb. & Gottsche — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- P. japonicum* (Steph.) C. Massal. — Piippo (1990), Lin (2000).
- P. pterospermum* C. Massal. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000).
- P. rupestre* (J.R. Forst. & G. Forst.) Steph. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000).
- Plagiochila arbuscula* (Brid. ex Lehm. & Lindenb.) Lindenb. — Piippo (1990), Lin (2000), So (2001), Gao & Lai (2003), Gao & Wu (2008).
- P. bantamensis* (Reinw. et al.) Mont. — Lin (2000).
- P. chinensis* Steph. — Piippo (1990), Lin (2000), So (2001), Gao & Wu (2008).
- P. delavayi* Steph. — Piippo (1990 as *P. sikutzuisana*), Lin (2000), So (2001).
- P. dendroides* (Nees) Lindenb. — Piippo (1990), Lin (2000), So (2001), Gao & Wu (2008).
- P. durelii* Schiffn. — Piippo (1990 as *P. vietnamica* and *P. sawadae*), Grolle & So (1999a), Lin (2000), So (2001), Gao & Wu (2008).
- P. duthiana* Steph. — Gao & Lai (2003).
- P. elegans* Mitt. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), So (2001), Gao & Wu (2008).
- P. flexuosa* Mitt. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), So (2001), Gao & Wu (2008).
- P. fordiana* Steph. — Piippo (1990), Grolle & So (1999b), Lin (2000), So (2001), Zhu & So (2001), Gao & Lai (2003).
- P. fruticosa* Mitt. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), So (2001), Gao & Wu (2008), Hsu & Huang (2010).
- P. gracilis* Lindenb. & Gottsche — Piippo (1990), Bapna

- & Kachroo (2000b), Bapna & Kachroo (2000b as *P. firma* var. *acicularis*), Lin (2000), So (2001), Gao & Wu (2008).
- P. gymnoclada* Sande Lac. — Piippo (1990), Lin (2000), So (2001).
- P. javanica* (Sw.) Nees & Mont. — Lin (2000).
- P. junghuhniana* Sande Lac. — Piippo (1990 as *P. massalongoana*), So & Grolle (1999), Lin (2000), So (2001), Gao & Wu (2008).
- P. khasiana* Mitt. — Lin (2000), So (2001), Gao & Wu (2008).
- P. kurzii* Steph. — Piippo (1990 as *P. tobagensis*), Grolle & So (1999c), So (2001).
- P. magna* Inoue — Piippo (1990), Lin (2000), So (2001), Gao & Wu (2008).
- P. multipinnula* Herzog & S. Hatt. — Piippo (1990), Grolle (1999d), Lin (2000), So (2001), Gao & Wu (2008).
- P. nepalensis* Lindenb. — Bapna & Kachroo (2000b as *P. gollanii*), Lin (2000), So (2001), Zhu & So (2001), Gao & Wu (2008), Hsu & Huang (2010 as *P. gollanii*).
- P. obtusa* Lindenb. — Piippo (1990 as *P. eberhardtii*), Grolle & So (1999c), Lin (2000), So (2001).
- P. ovalifolia* Mitt. — Piippo (1990), Lai (1995), Lin (2000), So (2001), Gao & Wu (2008), Hsu & Huang (2010).
- P. parvifolia* Lindenb. — Piippo (1990 as *P. yokogurensis*), Grolle & So (1999b), Lin (2000), So (2001), Gao & Wu (2008).
- P. peculiaris* Schiffn. — Piippo (1990), Piippo (1990 as *P. griffithiana*), Grolle & So (1999c), Lin (2000), So (2001), Gao & Wu (2008), Hsu & Huang (2010).
- P. perserrata* Herzog — Piippo (1990), Grolle & So (1999c), Bapna & Kachroo (2000b), Lin (2000), So (2001).
- P. porelloides* (Torr. ex Nees) Lindenb. — Lin (2000).
- P. pulcherrima* Horik. — Piippo (1990), Lin (2000), So (2001), Gao & Wu (2008), Hsu & Huang (2010).
- P. sciophila* Nees ex Lindenb. — Piippo (1990), Lai (1995), So & Grolle (1999), Lin (2000), So (2001), Gao & Wu (2008), Hsu & Huang (2010 as *P. acanthophylla* subsp. *japonica*).
- P. secretifolia* Mitt. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), So (2001), Gao & Wu (2008).
- P. semidecurrans* (Lehm. & Lindenb.) Lindenb. — Piippo (1990), So & Grolle (1999), Lin (2000), So (2001), Zhang & Wu (2006), Gao & Wu (2008).
- P. singularis* Schiffn. — Piippo (1990), Lin (2000), So (2001).
- P. taiwanensis* Inoue — Piippo (1990), Lin (2000), So (2001), Gao & Wu (2008).
- P. trabeculata* Steph. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Hsu & Huang (2010).
- P. vexans* Schiffn. ex Steph. — So (2001), Gao & Wu (2008).
- P. wangii* Inoue — Gao & Cao (2000), Lin (2000), So (2001), Gao & Lai (2003), Gao & Wu (2008).
- P. zonata* Steph. — Piippo (1990), Lin (2000), So (2001).
- Plagiochilium braunianum* (Nees) S. Hatt. — Piippo (1990), Gao & Cao (2000), Gao & Lai (2003).
- P. mayebarae* S. Hatt. — Grolle (1966 as *P. mayebarae* fo. *integerrimus*), Piippo (1990), Bapna & Kachroo (2000b), Bapna & Kachroo (2000b as *P. mayebarae* fo. *integerrimus*), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Zheng *et al.* (2005).
- P. oppositum* (Reinw. *et al.*) S. Hatt. — Piippo (1990), Lin (2000).
- Pleurozia acinosa* (Mitt.) Trevis. — Piippo (1990), Thiers (1993), Bai & Li (1998), Lin (2000), Gao & Cao (2000), Gao & Lai (2003), Lai (2007), Gao & Wu (2008), Hsu & Huang (2010).
- P. gigantea* (F. Weber) Lindb. — Bai & Li (1998), Hsu & Huang (2010).
- P. purpurea* Lindb. — Piippo (1990), Thiers (1993), Bai & Li (1998), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Lai (2007), Hsu & Huang (2010).
- P. subinflata* (Austin) Austin — Piippo (1990 as *Eopleurozia giganteoides*), Thiers (1993), Bai & Li (1998), Gao & Cao (2000 as *E. giganteoides*), Lin (2000 as *E. giganteoides*), Gao & Lai (2003 as *E. giganteoides*), Lai (2007).
- Plicanthus birmensis* (Steph.) R.M. Schust. — Piippo (1990 as *Chandonanthus birmensis*), Bapna & Kachroo (2000b as *C. birmensis*), Lin (2000 as *C. birmensis*), Gao & Lai (2003 as *C. birmensis*), Gao & Wu (2008 as *C. birmensis*).
- P. hirtellus* (F. Weber) R.M. Schust. — Piippo (1990 as *Chandonanthus hirtellus*), Bapna & Kachroo (2000b as *C. hirtellus*), Gao & Cao (2000 as *C. hirtellus*), Lin (2000 as *C. hirtellus*), Wu (2000 as *C. hirtellus*), Gao & Lai (2003 as *C. hirtellus*), Gao & Wu (2008 as *C. hirtellus*), Hsu & Huang (2010 as *C. hirtellus*).
- Porella acutifolia* (Lehm. & Lindenb.) Trevis. subsp. *acutifolia* — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- P. acutifolia* (Lehm. & Lindb.) Trevis. subsp. *tosana* (Steph.) S. Hatt. — Hattori (1978), Piippo (1990), Gao & Cao (2000), Lin (2000), So (2002), Gao & Lai (2003).
- P. arboris-vitae* (With.) Grolle — Piippo (1990).
- P. caespitans* (Steph.) S. Hatt. var. *caespitans* — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- P. caespitans* (Steph.) S. Hatt. var. *cordifolia* (Steph.) S. Hatt. — Hattori (1970 as *P. caespitans* var. *setigera*), Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010 as *P. caespitans* var. *setigera*).
- P. caespitans* (Steph.) S. Hatt. var. *nipponica* S. Hatt. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- P. densifolia* (Steph.) S. Hatt. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- P. gracillima* Mitt. subsp. *gracillima* — Piippo (1990), Gao & Cao (2000), Gao & Lai (2003).
- P. gracillima* Mitt. subsp. *urogea* (C. Massal.) S. Hatt. & M.X. Zhang — Hattori (1971 as *P. gracillima* var. *urogea*), Hattori (1976b as *P. gracillima* var. *urogea*), Piippo (1990), Bapna & Kachroo (2000b), Lin (2000).
- P. grandiloba* Lindb. — Piippo (1990), Lin (2000).
- P. japonica* (Sande Lac.) Mitt. — Hattori (1967), Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Hsu &

- Huang (2010).
- P. obtusata* (Taylor) Trevis. var. *macroloba* (Steph.) S. Hatt. & M.X. Zhang — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- P. perrottetiana* (Mont.) Trevis. — Piippo (1990), Lai (1995), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- P. plumosa* (Mitt.) Parihar — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- P. ulophylla* (Steph.) S. Hatt. — Piippo (1990 as *Macvicaria ulophylla*), Gao & Cao (2000 as *M. ulophylla*), Lin (2000 as *M. ulophylla*), Gao & Lai (2003 as *M. ulophylla*).
- P. vernicosa* Lindb. — Piippo (1990), Lin (2000).
- Preiszia quadrata* (Scop.) Nees — Piippo (1990), Lin (2000).
- Pseudolepicolea andoi* (R.M. Schust.) Inoue — Piippo (1990), Bapna & Kachroo (2000a as *P. trollii* subsp. *andoi*), Lin (2000).
- Ptychanthus striatus* (Lehm. & Lindenb.) Nees — Piippo (1990), Lin (1990), Lai (1995), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Yang (2009), Hsu & Huang (2010).
- Radula acuminata* Steph. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Zhu & Lai (2003), Hsu & Huang (2010).
- R. acuta* Mitt. — Piippo (1990 as *R. apiculata*), Lin (2000 as *R. apiculata*), Gao & Wu (2008 as *R. apiculata*).
- R. anceps* Sande Lac. — Piippo (1990), Lin (2000).
- R. auriculata* Steph. — Piippo (1990), Gao & Wu (2008).
- R. campanigera* Mont. — Piippo (1990), Lin (2000), Hsu & Huang (2010).
- R. cavifolia* Hampe ex Gottsche et al. — Piippo (1990), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Gao & Wu (2008).
- R. chinensis* Steph. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- R. constricta* Steph. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- R. formosa* (C.F.W. Meissn. ex Spreng.) Nees — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008).
- R. fulvifolia* (Hook. f. & Taylor) Gottsche et al. — Gao & Cao (2000 as *R. meyeri*), Gao & Lai (2003 as *R. meyeri*).
- R. inouei* K. Yamada — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008).
- R. japonica* Gottsche ex Steph. — Piippo (1990), Lin (2000).
- R. javanica* Gottsche — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Hsu & Huang (2010), Hsu & Huang (2010 as *R. variabilis*).
- R. kojana* Steph. — Lai (1990), Piippo (1990), Lin (2000), Zhu & So (2001), Gao & Wu (2008).
- R. nymanii* Steph. — Piippo (1990), Lin (2000), Gao & Wu (2008).
- R. obscura* Mitt. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Hsu & Huang (2010).
- R. okamurana* Steph. — Piippo (1990), Lin (2000).
- R. onraedtii* K. Yamada — Lin (2000), Gao & Wu (2008).
- R. oyamensis* Steph. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008), Hsu & Huang (2010).
- R. perrottetii* Gottsche ex Steph. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Gao & Lai (2003), Hsu & Huang (2010), Hsu & Huang (2010 as *R. valida*).
- R. philippinensis* K. Yamada — Piippo (1990), Lin (2000), Gao & Wu (2008).
- R. reflexa* Nees & Mont. — Hsu & Huang (2010).
- R. retroflexa* Taylor — Piippo (1990), Piippo (1990 as *R. retroflexa* var. *fauciloba*), Lai (1995), Bapna & Kachroo (2000b), Lin (2000), Lin (2000 as *R. retroflexa* var. *fauciloba*), Gao & Wu (2008), Gao & Wu (2008 as *R. retroflexa* var. *fauciloba*), Hsu & Huang (2010).
- R. tjobodensis* K.I. Goebel — Zhu & So (2001).
- R. tokiensis* Steph. — Piippo (1990), Piippo (1990 as *R. kanemarui*), Bapna & Kachroo (2000b), Lin (2000).
- R. yangii* K. Yamada — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Gao & Wu (2008).
- Reboulia hemisphaerica* (L.) Raddi — Lai (1990), Piippo (1990), Lai (1995), Lin (2000), Hsu & Huang (2010).
- Riccardia angustata* Horik. — Piippo (1990), Lin (2000).
- R. chamedryfolia* (With.) Grolle — Piippo (1990), Lin (2000).
- R. crassa* (Schwägr.) Carrington & Pearson — Yao & Lin (2010).
- R. crenulata* Schiffn. — Piippo (1990), Lin (2000).
- R. flavovirens* Furuki — Yao & Lin (2010).
- R. formosensis* (Steph.) Horik. — Piippo (1990), Lin (2000).
- R. graeffei* (Steph.) Hewson — Piippo (1990 as *R. platyclada*), Lin (2000 as *R. platyclada*).
- R. kodamae* Mizut. & S. Hatt. — Yao & Lin (2010).
- R. latifrons* (Lindb.) Lindb. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- R. marginata* (Colenso) Pearson var. *pacifica* Furuki — Yao & Lin (2010).
- R. multifida* (L.) Gray — Lai (1990), Lai (1995), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- R. nagasakiensis* (Steph.) S. Hatt. — Piippo (1990 as *R. jackii*).
- R. palmata* (Hedw.) Carruth. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- R. pumila* Furuki — Yao & Lin (2010).
- R. submultifida* Horik. — Piippo (1990), Lin (2000).
- Riccia fluitans* L. — Lai (1990), Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Zhang & Wu (2006).
- R. glauca* L. — Lai (1990), Piippo (1990), Lin & Yang (1992), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- Ricciocarpos natans* (L.) Corda — Lai (1990), Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- Saccogynidium irregularospinum* C. Gao, T. Cao & M.J. Lai — Gao et al. (2001), Gao & Wu (2008).
- S. muricellum* (De Not.) Grolle — Piippo (1990), Lai (1995), Lin (2000), Gao et al. (2001), Gao & Wu (2008).
- S. rigidulum* (Nees) Grolle — Piippo (1990), Lin (2000), Gao



- et al.* (2001), Gao & Lai (2003), Gao & Wu (2008).
- Scapania ampliata* Steph. — Zuo & Cao (2007), Zuo *et al.* (2008).
- S. bolanderi* Austin — Piippo (1990), Lin (2000), Gao & Wu (2008), Zuo *et al.* (2008).
- S. ciliata* Sande Lac. — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Wu (2008), Zuo *et al.* (2008), Hsu & Huang (2010).
- S. ciliatospinosa* Horik. — Piippo (1990), Potemkin (1999), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Gao & Wu (2008), Zuo *et al.* (2008).
- S. curta* (Mart.) Dumort. — Gao & Wu (2008), Zuo *et al.* (2008).
- S. ferruginea* (Lehm. & Lindenb.) Gottsche *et al.* — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Wu (2008), Zuo *et al.* (2008).
- S. gracilis* Lindb. — Piippo (1990).
- S. griffithii* Schiffn. — Gao & Wu (2008), Zuo *et al.* (2008).
- S. javanica* Gottsche — Piippo (1990), Lin (2000).
- S. ligulata* Steph. subsp. *ligulata* — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000), Zuo *et al.* (2008).
- S. ligulata* Steph. subsp. *stephanii* (Müll. Frib.) Potemkin *et al.* — Piippo (1990 as *S. stephanii*), Lai (1995 as *S. stephanii*), Lin (2000 as *S. stephanii*), Potemkin *et al.* (2004), Gao & Wu (2008 as *S. stephanii*), Zuo *et al.* (2008 as *S. stephanii*), Hsu & Huang (2010), Hsu & Huang (2010 as *S. stephanii*).
- S. maxima* Horik. — Bapna & Kachroo (2000b), Potemkin (2002), Hsu & Huang (2010).
- S. nemorea* (L.) Grolle — Piippo (1990), Lin (2000), Zuo *et al.* (2008).
- S. nimbosea* Taylor *ex* Lehm. — Zuo *et al.* (2008).
- S. ornithopodioides* (With.) Waddell — Piippo (1990), Lin (2000), Gao & Wu (2008), Zuo *et al.* (2008), Hsu & Huang (2010).
- S. parvixta* Steph. — Piippo (1990), Piippo (1990 as *S. glaucoviridis*), Bapna & Kachroo (2000b as *S. glaucoviridis*), Lin (2000 as *S. glaucoviridis*), Gao & Wu (2008 as *S. parvidens* and *S. glaucoviridis*), Gao & Wu (2008), Zuo *et al.* (2008), Zuo *et al.* (2008 as *S. glaucoviridis* and *S. parvidens*), Hsu & Huang (2010 as *S. glaucoviridis*).
- S. subnimbosea* Steph. — Piippo (1990), Gao & Cao (2000), Lin (2000 as *S. robusta*), Lin (2000), Gao & Lai (2003), Gao & Wu (2008), Zuo *et al.* (2008), Hsu & Huang (2010 as *S. robusta*).
- S. umbrosa* (Schrad.) Dumort. — Zuo *et al.* (2008).
- S. undulata* (L.) Dumort. — Piippo (1990), Lin (2000), Gao & Wu (2008).
- Schiffneria hyalina* Steph. — Lai (1990), Piippo (1990), Bapna & Kachroo (2000a), Lin (2000), Wu & Wang (2001), Gao & Lai (2003), Long (2005), Hsu & Huang (2010).
- Schiffneriolejeunea tumida* (Nees) Gradst. var. *haskarliana* (Gottsche) Gradst. & Terken — Piippo (1990 as *S. tumida*), He (1997 as *S. tumida*), Lin (2000 as *S. tumida*), Yang (2009).
- Schistochila acuminata* Steph. — Piippo (1990), Lin (2000), Wu & Jong (2001), Gao & Lai (2003), So (2003a), Gao & Wu (2004), Lai (2007), Gao & Wu (2008).
- S. aligera* (Nees & Blume) J.B. Jack & Steph. — Piippo (1990 as *Gottschea aligera*), Bapna & Kachroo (2000b), Gao & Cao (2000 as *G. aligera*), Lin (2000 as *G. aligera*), Gao & Lai (2003 as *G. aligera*), So (2003a), Gao & Wu (2004 as *G. aligera*), Lai (2007 as *G. aligera* and *G. philippinensis*), Gao & Wu (2008 as *G. aligera* and *G. philippinensis*), Hsu & Huang (2010 as *Paraschistochila aligera* and *P. philippinensis*).
- S. blumei* (Nees) Trevis. — Piippo (1990), Lin (2000), Gao & Lai (2003), So (2003a), Lai (2007), Gao & Wu (2008).
- S. minor* C. Gao & Y.H. Wu — Gao & Wu (2004), Gao & Wu (2008).
- S. nuda* Horik. — Piippo (1990 as *Gottschea nuda*), Gao & Cao (2000 as *G. nuda*), Lin (2000 as *G. nuda*), Gao & Lai (2003 as *G. nuda*), So (2003a), Lai (2007 as *G. nuda*), Gao & Wu (2008 as *G. nuda*).
- Schistochilopsis cornuta* (Steph.) Konstant. — Piippo (1990 as *Lophozia cornuta*), Gao & Cao (2000 as *L. cornuta*), Lin (2000 as *L. cornuta*).
- S. incisa* (Schrad.) Konstant. — Piippo (1990 as *Lophozia incisa*), Bapna & Kachroo (2000b as *L. incisa*), Gao & Cao (2000 as *L. incisa*), Lin (2000 as *L. incisa*), Gao & Wu (2008 as *L. incisa*).
- S. nakanishii* (Inoue) Konstant. — Piippo (1990 as *Lophozia nakanishii*), Lin (2000 as *L. nakanishii*).
- Solenostoma appressifolium* (Mitt.) Váňa & D.G. Long — Piippo (1990 as *Jungermannia appressifolia*), Bapna & Kachroo (2000a as *J. appressifolia*), Gao & Cao (2000 as *J. appressifolia*), Lin (2000 as *J. appressifolia*), Gao & Bai (2001 as *Haplozia rotundifolia*), Gao (2003 as *J. appressifolia*), Gao & Lai (2003 as *J. appressifolia*).
- S. caespiticium* (Lindenb.) Steph. — Hsu & Huang (2010 as *Aplozia caespiticia*).
- S. comatum* (Nees) C. Gao — Piippo (1990 as *Jungermannia comata*), Bapna & Kachroo (2000a as *J. comata*), Lin (2000 as *J. comata*), Váňa *et al.* (2005 as *J. comata*).
- S. gracillimum* (Sm.) R.M. Schust. — Piippo (1990 as *Jungermannia gracillima*), Hsu & Huang (2010 as *Plectocolea crenulata*).
- S. hasskarlianum* (Nees) R.M. Schust. *ex* Váňa & D.G. Long — Piippo (1990 as *Jungermannia hasskarliana*), Bapna & Kachroo (2000a as *J. hasskarliana*), Lin (2000 as *J. hasskarliana*).
- S. hyalinum* (Lyell) Mitt. — Piippo (1990 as *Jungermannia hyalina*), Lin (2000 as *J. hyalina*).
- S. infusum* (Mitt.) Hentschel — Lai (1990 as *Jungermannia infusca*), Piippo (1990 as *J. infusca*), Bapna & Kachroo (2000a as *J. infusca*), Lin (2000 as *J. infusca*), Váňa *et al.* (2005 as *J. infusca*).
- S. pseudocyclops* (Inoue) Váňa & D.G. Long — Piippo (1990 as *Jungermannia pseudocyclops*), Bapna & Kachroo (2000a as *J. pseudocyclops*), Gao & Cao (2000 as *J. pseudocyclops*), Lin (2000 as *J. pseudocyclops*), Gao & Bai (2001 as *J. pseudocyclops*), Gao & Lai (2003 as *J. pseudocyclops*).
- S. pyriformum* Steph. — Lai (1990 as *Jungermannia pyriforma*), Piippo (1990 as *J. pyriforma*), Piippo *et al.* (1997 as *J. pyriforma*), Bapna & Kachroo (2000a as *J. pyriforma*), Lin (2000 as *J. pyriforma*).

- S. radiculosum* Mitt. — Gao (2003 as *Jungermannia radiculosa*).
- S. schaulianum* (Steph.) Váňa & D.G. Long — Piippo (1990 as *Jungermannia schauliana*), Bapna & Kachroo (2000a as *J. schauliana*), Gao & Cao (2000 as *J. schauliana*), Lin (2000 as *J. schauliana*), Gao & Bai (2001 as *J. schauliana*), Gao (2003 as *J. schauliana*), Gao & Lai (2003 as *J. schauliana*).
- S. speciosum* (Horik.) Hentschel et al. — Piippo (1990 as *Scaphophyllum speciosum*), Schuster (1998 as *S. speciosum* subsp. *speciosum*), Lai (1999 as *S. speciosum*), Bapna & Kachroo (2000a as *S. speciosum*), Gao & Cao (2000 as *S. speciosum*), Lin (2000 as *S. speciosum*), Gao & Bai (2001 as *S. speciosum*), Gao (2003 as *S. speciosum*), Gao & Lai (2003 as *S. speciosum*), Lai (2007 as *S. speciosum*).
- S. subrubrum* (Schiffn. ex Steph.) Váňa & D.G. Long — Piippo (1990 as *Jungermannia subrubra*), Bapna & Kachroo (2000a as *J. subrubra*), Lin (2000 as *J. subrubra*).
- S. tetragonum* (Lindenb.) R.M. Schust. ex Váňa & D.G. Long — Piippo (1990 as *Jungermannia tetragona*), Gao & Cao (2000 as *J. tetragona*), Lin (2000 as *J. tetragona*), Gao & Bai (2001 as *J. tetragona*), Gao & Lai (2003 as *J. tetragona*), Váňa et al. (2005 as *J. tetragona*).
- S. truncatum* (Nees) R.M. Schust. ex Váňa & D.G. Long — Lai (1990 as *Jungermannia truncata*), Piippo (1990 as *J. truncata*), Lai (1995 as *J. truncata*), Bapna & Kachroo (2000a as *J. truncata*), Lin (2000 as *J. truncata*), Gao & Lai (2003 as *J. truncata*), Váňa et al. (2005 as *J. truncata*).
- S. virgatum* (Mitt.) Váňa & D.G. Long — Piippo (1990 as *Jungermannia virgata*), Lin (2000 as *J. virgata*), Váňa et al. (2005 as *J. virgata*).
- S. vulcanicola* (Schiffn.) Nyushko — Lai (1990 as *Jungermannia vulcanicola*).
- Sphenolobopsis pearsonii* (Spruce) R.M. Schust. — Piippo (1990), Lin (2000), Gao & Wu (2008), Huang & Lin (2009 as *Cephalozicopsis pearsonii*).
- Spruceanthus polymorphus* (Sande Lac.) Verd. — Piippo (1990), He (1997 as *Archilejeunea polymorpha*), Bapna & Kachroo (2000b as *A. polymorpha*), Gao & Cao (2000 as *A. polymorpha*), Lin (2000), Gao & Lai (2003 as *A. polymorpha*), Yang (2009 as *A. polymorpha*), Hsu & Huang (2010).
- S. semirepandus* (Nees) Verd. — Piippo (1990), He (1997), Bapna & Kachroo (2000b), Lin (2000), Zhu & So (2001), Yang (2009), Hsu & Huang (2010).
- Syzygiella autumnalis* (DC.) Feldberg et al. — Piippo (1990 as *Jamesoniella autumnalis*), Bapna & Kachroo (2000b as *J. autumnalis*), Lin (2000 as *J. autumnalis*), Zhu & So (2001 as *J. autumnalis*), Hsu & Huang (2010 as *J. autumnalis*).
- S. nipponica* (S. Hatt.) Feldberg et al. — Piippo (1990 as *J. nipponica*), Bapna & Kachroo (2000b as *J. nipponica*), Gao & Cao (2000 as *J. nipponica*), Lin (2000 as *J. nipponica*).
- S. securifolia* (Nees ex Lindenb.) Inoue — Piippo (1990), Lin (2000), Gao & Wu (2008).
- Targionia formosica* Horik. — Piippo (1990), Lin (2000).
- T. hypophylla* L. — Piippo (1990), Bapna & Kachroo (2000b), Lin (2000).
- Telaranea neesii* (Lindenb.) Fulford — Wu & Yamada (1996).
- T. planifolia* (Steph.) J.J. Engel & G.L. Merr. — Engel & Merrill (2004).
- T. wallichiana* (Gottsche) R.M. Schust. — Piippo (1990 as *Lepidozia wallichiana*), Lai (1995 as *L. wallichiana*), Bapna & Kachroo (2000a as *L. wallichiana*), Lin (2000 as *L. wallichiana*), Gao & Lai (2003 as *L. wallichiana*), Engel & Merrill (2004), Hsu & Huang (2010 as *L. wallichiana*).
- Temnoma setigerum* (Lindenb.) R.M. Schust. — Long & Grolle (1990), Piippo (1990), Bapna & Kachroo (2000a), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003).
- Tetralophozia filiformis* (Steph.) Urmi — Piippo (1990), Wang (1993 as *Chandonanthus pusillus*), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Wu (2008), Hsu & Huang (2010 as *C. pusillus*).
- T. setiformis* (Ehrh.) Schljakov — Piippo (1990).
- Thysananthus aculeatus* Herzog — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- T. flavescens* (S. Hatt.) Gradst. — Lin (2000), Yang (2009).
- T. spathulistipus* (Reinw. et al.) Lindenb. — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- Trichocolea merrilliana* Steph. — Piippo (1990), Gao & Cao (2000), Lin (2000), Wu & Wang (2001 as *T. lumbrioides*), Gao (2003), Gao & Lai (2003), Lai (2007), Hsu & Huang (2010).
- T. tomentella* (Ehrh.) Dumort. — Piippo (1990), Lai (1995), Bapna & Kachroo (2000a), Lin (2000), Gao & Lai (2003), Lai (2007), Lai (2007 as *T. pluma*), Hsu & Huang (2010), Hsu & Huang (2010 as *T. pluma*).
- Trichocoleopsis sacculata* (Mitt.) S. Okamura — Piippo (1990), Lin (2000), Gao & Lai (2003), Lai (2007), Hsu & Huang (2010).
- Tritomaria exsecta* (Schmidel ex Schrad.) Schiffn. ex Loeske — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Wu (2008).
- T. quinquentata* (Huds.) H. Buch — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).
- Trocholejeunea infuscata* (Mitt.) Verd. — Piippo (1990), Bapna & Kachroo (2000b).
- T. sandvicensis* (Gottsche) Mizut. — Lai (1990), Piippo (1990), Lai (1995), He (1997), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Zhu & So (2001), Zhang & Wu (2006), Yang (2009), Hsu & Huang (2010).
- Tuyamaella molischii* (Schiffn.) S. Hatt. var. *molischii* — Piippo (1990), He (1997), Lin (2000), Yang (2009).
- T. molischii* (Schiffn.) S. Hatt. var. *taiwanensis* R.L. Zhu & M.L. So — Zhu & So (2000a), Zhu & So (2001).
- Wetsteinia inversa* (Sande Lac.) Schiffn. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003).
- W. rotundifolia* (Horik.) Grolle — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao (2003), Gao & Lai (2003).
- Wiesnerella denudata* (Mitt.) Steph. — Piippo (1990), Bapna

& Kachroo (2000b), Lin (2000), Gao & Lai (2003).  
*Xenochila integrifolia* (Mitt.) Inoue — Piippo (1990), Bapna & Kachroo (2000b), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).  
*Zoopsis liukuensis* Horik. — Piippo (1990), Gao & Cao (2000), Lin (2000), Gao & Lai (2003).

## Hornworts (Anthocerotophyta)

*Anthoceros agrestis* Paton — Lai & Wang-Yang (1976 as *A. nagasakiensis*).  
*A. angustus* Steph. — Long & Grolle (1990 as *A. formosae*), Piippo (1990 as *A. formosae*), Hasegawa (1992), Hasegawa (1993), Bapna & Kachroo (2000a as *A. formosae*), Gao & Cao (2000 as *A. formosae*), Lin (2000), Zhang & Wu (2006 as *A. formosae*).  
*A. fusiformis* Austin var. *fusiformis* — Piippo (1990).  
*A. fusiformis* Austin var. *taiwanensis* J. Haseg. — Hasegawa (1993).  
*A. punctatus* L. — Piippo (1990), Hasegawa (1993), So (1995), Lin (2000).  
*Dendroceros japonicus* Steph. — Piippo (1990), Hasegawa (1993), Gao & Cao (2000), Lin (2000), Wu & Wang (2001), Gao & Lai (2003).  
*D. javanicus* (Nees) Nees — Piippo (1990), Hasegawa (1993), Lin (2000).  
*D. tubercularis* S. Hatt. — Piippo (1990), Hasegawa (1993), Lin (2000).  
*Folioceros amboinensis* (Schiffn.) Piippo — Hasegawa (1993 as *Anthoceros amboinensis*).  
*F. appendiculatus* (Steph.) J. Haseg. — Hasegawa (1993 as *Anthoceros* aff. *appendiculatus*).  
*F. fuciiformis* (Mont.) D.C. Bhardwaj — Piippo (1990 as *F. vesiculosus*), Hasegawa (1993 as *Anthoceros fuciiformis*), Lin (2000 as *F. vesiculosus*).  
*F. glandulosus* (Lehm. & Lindenb.) D.C. Bharadwaj — Piippo (1990), Lin (2000).  
***Folioceros verruculosus*** (J. Haseg.) R.L. Zhu & M.J. Lai, *comb. nova*. Basionym: *Anthoceros verruculosus* J. Haseg., Acta Phytotax. Geobot. 44(2): 103. 1993. According to the description and illustration by Hasegawa (1993), *A. verruculosus* has ecostate, cavernous thali, and strongly thick-walled elaters. Hence it is better to place it in the genus *Folioceros*. — Hasegawa (1993 as *Anthoceros verruculosus*).  
*Megaceros flagellaris* (Mitt.) Steph. — Piippo (1990), Hasegawa (1993), Gao & Cao (2000), Lin (2000), Gao & Lai (2003), Zhang & Wu (2006).  
*Notothylas javanica* (Sande Lac.) Gottsche — Piippo (1990), Hasegawa (1993), Lin (2000).  
*N. orbicularis* (Schwein.) Sull. ex A. Gray — Piippo (1990), Lin (2000), Zhang & Wu (2006 as *N. japonica*).  
*Phaeoceros carolinianus* (Michx.) Prosk. — Lai (1990), Piippo (1990 as *P. laevis* subsp. *carolinianus*), Hasegawa (1993), Lin (2000 as *P. laevis* subsp. *carolinianus*), Guo & Zhang (2001 “1999” as *P. laevis* subsp. *carolinianus*).  
*P. exiguus* (Steph.) J. Haseg. — Hasegawa (1993).

*P. laevis* (L.) Prosk. — Piippo (1990), Piippo (1990 as *P. miyakeanus*), Hasegawa (1991), So (1995), Bapna & Kachroo (2000a), Lin (2000 as *P. miyakeanus*), Zhang & Wu (2006).  
*P. pearsonii* (M. Howe) Prosk. — Piippo (1990 as *Anthoceros pearsonii*).

## Doubtful and excluded records

*Asterella tenella* (L.) Beauv. — All records of *A. tenella* from outside of North America are wrong (Grolle 1989).  
*Bazzania pearsonii* Steph. — The record of this species in Taiwan by Lai and Wang-Yang (1976) and Lai (1977) is erroneous (Yamada & Lai 1979).  
*B. mayebarae* S. Hatt. — The record of this species in Taiwan by Lai and Wang-Yang (1976) and Lai (1977) is erroneous (Yamada & Lai 1979).  
*Cheilolejeunea mariana* (Gottsche) B.M. Thiers & Gradst. — This species has been excluded from China and the records from China were erroneous identifications, and the Taiwan specimen actually is assignable to *Lopholejeunea javanica* (Zhu & So 2000b), which is a synonym of *L. nigricans* (Zhu & Gradstein 2005).  
*C. minutissima* Nakanishi — *C. minutissima* was included in Kuo and Chiang (1988), Piippo (1990) and He (1997). The name “*Cheilolejeunea minutissima*” (Nakanishi 1964: 72) was a misprinting of *Cololejeunea minutissima* (Sm.) Schiffn. Hence, the species *Cheilolejeunea minutissima* Nakanishi does not exist (Zhu *et al.* 2002).  
*Cololejeunea filicis* (Herzog) Piippo — The record in Taiwan (He 1997) was an erroneous citation.  
*C. reinecheana* Steph. — All records of *C. reinecheana* from China are actually *C. formosana*, a synonym of *C. ceratitilobula* (Zhu & So 1998b).  
*Conocephalum riccardia* B.Y. Yang, *nom. nud.*  
*Diplastolejeunea pellucida* (Meissn.) Schiffn. — *D. pellucida* has to be excluded from the bryoflora of Taiwan, because the vouchers from Taiwan are assignable to *D. cavifolia* (Zhu & So 2001). The reliable records of the true *D. pellucida* are restricted to the Neotropics (Grolle 1995).  
*Fossombronia cristula* Austin — This species is endemic to the eastern North America. All records from Taiwan by Piippo (1990), Piippo (1990 as *F. australi-nipponica*) and Lin (2000 as *F. australi-nipponica*) are erroneous determinations (Krayesky *et al.* 2005).  
*Frullania fauriana* Steph. — The records of *F. fauriana* from Taiwan are doubtful (Kamimura 1961).  
*Herbertus longifissus* Steph. — *H. longifissus* is a species occurring only in Melanesia (So 2003c, Juslén 2006). Asian specimens named as *H. longifissus* revealed to be *H. sakuraii* (So 2003c), which is a synonym of *H. dicranus* (Hodgetts 2003).  
*H. maximus* B.Y. Yang, *nom. nud.*  
*Heteroscyphus acutangulus* (Schiffn.) Schiffn. — The record of this species in Taiwan is actually *Chiloscyphus sikimensis* (Wu & Gao 2008).  
*H. bescherellei* (Steph.) S. Hatt. fo. *tenella* Herzog, *nom. nud.*

- H. zollingeri* (Gottsche) Schiffn. fo. *pluridentatus* Herzog — *H. zollingeri* fo. *pluridentatus* was synonymized with *H. zollingeri* by Gao & Wu (2008).
- Jungermannia ariadne* Taylor ex Lehm. — The records of this species by Piippo (1990) and Bapna and Kachroo (2000a) in Taiwan belong to *J. hasskartiiana* (Váňa & Inoue 1983).
- J. kyushuensis* Amakawa — The specimen mentioned as *J. kyushuensis* by Lai and Wang-Yang (1976) and Hsu & Huang (2010) belongs to *J. infusca* (Piippo 1990), which is a synonym of *Solenostoma infusum* (Hentschel et al. 2007a).
- Kurzia sylvatica* (A. Evans) Grolle — The record from Taiwan is questionable (Piippo 1990).
- Lejeunea planiloba* A. Evans — Horikawa (1934a) reported an epiphyllous occurrence of *L. planiloba* in Taiwan, but the voucher does not contain that species. Hence, the occurrence of this species in Taiwan remains doubtful (Zhu & So 2001).
- L. subigiensis* (Steph.) Steph. — The record of this species in Taiwan by Herzog and Noguchi (1955) is assignable to *L. anisophylla* Mont. (present study).
- Lepicolea scolopendra* (Hook.) Dumort. — The only record in China was based on Horikawa's collection from Taiwan, which actually was assignable to *L. yakusimensis* by Inoue (1978).
- Leptolejeunea emarginata* (Horik.) S. Hatt. — The type specimen of this species might be missing. Based on the protologue, *L. emarginata* appears to be closely related to *L. apiculata* (Zhu & So 2001).
- L. foliicola* Steph. — This name may be synonym of *L. elliptica* (Zhu & So 2001).
- Lophocolea semiteres* (Lehm.) Mitt. — *L. semiteres* (Piippo 1990, Lin 2000 as *Chiloscyphus semiteres*) does not occur in China (Piippo 1996).
- Lopholejeunea aberrantia* Horik. — The material in HIRO (including the type) may have been destroyed in 1945 when Hiroshima was bombed (H. Deguchi and T. Yamaguch pers. comm.). Judging from the original description and illustrations, this species may be a member of *Archilejeunea* because of its 5–6-keeled perianths without laciniae (Zhu & Gradstein 2005).
- L. applanata* (Reinw. et al.) Schiffn. — The records in Taiwan are *L. nigricans* (Zhu & Gradstein 2005).
- Marsupella commutata* (Limpr.) Bernet — The record in Taiwan by Bapna and Kachroo (2000b) is uncertain.
- Metzgeria albinea* Spruce — The record in China and Asia was excluded by So (2003b), the voucher specimens in China are referable to *M. foliicola*. *Metzgeria albinea* is known only from New Zealand (So 2003b).
- M. conjugata* Lindb. — All records of this species from Taiwan and mainland China are treated here as *M. lindbergii* Schiffn.
- Plagiochila blepharophora* (Nees) Lindenb. — Excluded from China by So (2001).
- P. debilis* Mitt. — According to So (2001), the record of *P. debilis* in Taiwan by Bapna and Kachroo (2000b) is uncertain.
- P. grandis* B.Y. Yang, *nom. nud.*
- P. hattoriana* Inoue, *nom. inval.* (So 2001).
- P. integrilobula* Schiffn. — The record in Taiwan is based on *P. tobagensis* (Herzog & Noguchi 1955), which is a synonym of *P. kurzii* (So 2001).
- P. microdonta* Mitt. — The record in China is based on erroneous identification (So 2001).
- P. oblonga* Inoue — The record in China (Taiwan) is based on erroneous identification (So 2001).
- P. philippinensis* Steph. — The record in China (Taiwan) is based on erroneous identification (So 2001).
- P. semidecurrens* var. *longifolia* Inoue — According to So (2001), the record of this species in Taiwan by Bapna and Kachroo (2000b) is uncertain.
- Porella revoluta* (Lehm.) Grolle — The record in Taiwan by Bapna and Kachroo (2000b) is uncertain.
- Pseudolepicolea trollii* (Herzog) Grolle & Ando — The records of this species by Piippo (1990) and Bapna and Kachroo (2000a) in Taiwan belong to *P. andoi* (Inoue 1978).
- Radula borneensis* Steph. — The record of *R. borneensis* from Taiwan is uncertain (Yamada & Inoue 1988).
- R. complanata* (L.) Dumort. — The record in Taiwan is uncertain (Yamada & Inoue 1988).
- R. lindenberghiana* Gottsche ex C. Hartm. — The records of *R. lindenberghiana* from Asia are based on erroneous determinations, this species is rejected from Taiwan (Yamada 1979).
- R. portoricensis* Steph. — According to Yamada & Inoue (1988), the occurrence of Latin American *R. guatemalensis* in Taiwan is improbable.
- R. protensa* Lindenb. — The specimens of *R. protensa* from Taiwan mentioned by Horikawa (1934a) belong to *R. acuminata* (Yamada & Inoue 1988).
- Treubia insignis* Goebel emend. R.M. Schust. & Scott — Inoue (1967) supposed that the specimens reported as *T. insignis* from Taiwan belong to *Apotreubia nana*.

## Synonyms

- Anastrophyllum acuminatum* (Horik.) N. Kitag. Accepted name: *A. minutum* (Schreb.) R.M. Schust. var. *acuminatum* (Horik.) T. Cao & J. Sun (Gao & Wu 2008).
- Anthoceros amboinensis* Schiffn. Accepted name: *Folioceros amboinensis* (Schiffn.) Piippo (Piippo 1993).
- Anthoceros appendiculatus* Steph. Accepted name: *Folioceros appendiculatus* (Steph.) J. Haseg. (Hasegawa 1986).
- Anthoceros formosae* Steph. Accepted name: *A. angustus* Steph. (Hasegawa 1992).
- Anthoceros fuciformis* Mont. Accepted name: *Folioceros fuciformis* (Mont.) D.C. Bhardwaj (Bhardwaj 1975).
- Anthoceros nagasakiensis* Steph. Accepted name: *A. agrestis* Paton (Stotler & Crandall-Stotler 2003).
- Anthoceros pearsonii* M. Howe. Accepted name: *Phaeoceros pearsonii* (M. Howe) Prosk. (Stotler & Crandall-Stotler 2005).
- Anthoceros verruculosus* J. Haseg. Accepted name: *Folioceros verruculosus* (J. Haseg.) R.L. Zhu & M.J. Lai (present paper).
- Aphanolejeunea angustiloba* Horik. Accepted name: *Colole-*



- jeunea sintenisii* (Steph.) Pócs (Pócs & Bernecker 2009).  
*Aphanolejeunea grossepapillosa* Horik. Accepted name: *Cololejeunea grossepapillosa* (Horik.) N. Kitag. (Pócs & Bernecker 2009).  
*Aphanolejeunea truncatifolia* Horik. Accepted name: *Cololejeunea diaphana* A. Evans (Pócs & Bernecker 2009).  
*Aplozia caespiticia* (Lindenb.) Dumort. — *Jungermannia caespiticia* Lindenb. (Váňa 1974). Accepted name: *Solenostoma caespiticium* (Lindenb.) Steph. (Váňa & Long 2009).  
*Apometzgeria pubescens* (Schrank) Kuwah. Accepted name: *Metzgeria pubescens* (Schrank) Raddi (So 2003b).  
*Archilejeunea polymorpha* (Sande Lac.) B.M. Thiers & Gradst. Accepted name: *Spruceanthus polymorphus* (Sande Lac.) Verd. (Gradstein *et al.* 2002).  
*Asterella monospiris* (Horik.) Horik. Accepted name: *A. wallichiana* (Lehm. & Lindenb.) Grolle (Long 2006).  
*Asterella yoshinagana* (Horik.) S. Hatt. Accepted name: *A. leptophylla* (Mont.) Grolle (Long 2006).  
*Bazzania oshimensis* (Steph.) Horik. Accepted name: *B. tridens* (Reinw. *et al.*) Trevis. (Inoue 1989).  
*Brachiolejeunea miyakeana* Steph. — *Archilejeunea mariana* auct. non (Gottsche) Steph. (Verdoorn 1934). Accepted name: *A. planiuscula* (Mitt.) Steph. (Thiers & Gradstein 1989).  
*Calypogeia trichomanis* (L.) Corda. Accepted name: *C. azurea* Stotler & Crotz (Stotler & Crotz 1983).  
*Caudalejeunea recurvistipula* (Gottsche) Schiffn. Accepted name: *Caudalejeunea reniloba* (Gottsche) Steph. (Zhu *et al.* 2011).  
*Cephalozia otaruensis* Steph. Accepted name: *C. hamatiloba* Steph. (Váňa 1988, 1993).  
*Chandonanthus birmensis* Steph. — *Temnoma birmense* (Steph.) Horik. (Horikawa 1951). Accepted name: *Plicanthus birmensis* (Steph.) R.M. Schust. (Schuster 2002).  
*Chandonanthus hirtellus* (F. Weber) Mitt. Accepted name: *Plicanthus hirtellus* (F. Weber) R.M. Schust. (Schuster 2002).  
*Chandonanthus pusillus* Steph. — *Temnoma setiforme* (Ehrh.) Howe (Hattori 1948). Accepted name: *Tetralophozia filiformis* (Steph.) Urmí (Urmí 1983).  
*Cheilolejeunea longiloba* (Hoffm.) J.J. Engel & B.C. Tan. — *C. imbricata* (Nees) S. Hatt. (Zhu *et al.* 2002). Accepted name: *C. trapezia* (Nees) Kachroo & R.M. Schust. (Zhu & Grolle 2004).  
*Chiloscyphus cuspidatus* (Nees) J.J. Engel & R.M. Schust. — *Lophocolea cuspidata* (Nees) Limpr. Accepted name: *L. bidentata* (L.) Dumort. (Furuki & Mizutani 2004).  
*Chiloscyphus fragrans* (Moris & De Not.) J.J. Engel & R.M. Schust. Accepted name: *Lophocolea fragrans* (Moris & De Not.) Gottsche *et al.* (Grolle & Long 2000).  
*Chiloscyphus horikawanus* (S. Hatt.) J.J. Engel & R.M. Schust. Accepted name: *Lophocolea horikawana* S. Hatt. (Inoue 1959).  
*Chiloscyphus latifolius* (Nees) J.J. Engel & R.M. Schust. Accepted name: *Lophocolea bidentata* (L.) Dumort. (Grolle & Long 2000).  
*Chiloscyphus minor* (Nees) J.J. Engel & R.M. Schust. Accepted name: *Lophocolea minor* Nees (Inoue 1959).  
*Chiloscyphus muricatus* (Lehm.) J.J. Engel & R.M. Schust. Accepted name: *Lophocolea muricata* (Lehm.) Nees (Grolle 1995).  
*Chiloscyphus profundus* (Nees) J.J. Engel & R.M. Schust. Accepted name: *Lophocolea heterophylla* (Schrad.) Dumort. (Grolle & Long 2000).  
*Chiloscyphus schiffneri* J.J. Engel & R.M. Schust., *nom. illeg.* Accepted name: *Chiloscyphus engelii* R.L. Zhu & M.J. Lai (present paper).  
*Chiloscyphus semiteres* (Lehm.) Lindenb. Accepted name: *Lophocolea semiteres* (Lehm.) Mitt. (Grolle & Long 2000).  
*Cololejeunea formosana* Mizut. Accepted name: *C. ceratitlobula* (P.C. Chen) R.M. Schust. (Zhu & So 1998b).  
*Cololejeunea goebelii* (Gottsche ex Schiffn.) Schiffn. Accepted name: *C. trichomanis* (Gottsche) Steph. (Grolle 2001).  
*Cololejeunea mackeeana* Tixier. Accepted name: *C. angustiflora* (Steph.) Mizut. (Zhu & So 2002).  
*Cololejeunea magnipapillosa* (Kamim.) P.C. Chen & P.C. Wu. Accepted name: *C. peraffinis* (Schiffn.) Schiffn. (Zhu & So 2001).  
*Cololejeunea oshimensis* (Horik.) Benedix. Accepted name: *C. inflata* Steph. (Zhu & So 2001).  
*Cololejeunea peraffinis* (Schiffn.) Schiffn., Consp. Hepat. Arch. Ind.: 245. 1898. — *C. peraffinis* fo. *corticola* (Benedix ex Herzog) S.H. Lin, Liverw. Fl. Taiwan: 401. 2000, *syn. nov.* *Taeniolejeunea peraffinis* (Schiffn.) Zwick. fo. *corticola* Benedix ex Herzog, J. Hattori Bot. Lab. 14: 55. 1955. TYPE: China. Taiwan, Botel Tobago, Am Kamm, ca. 460 m, auf Borke, G.H. Schwabe 123 p.p. (holotype JE!).  
*Cololejeunea scabriflora* Gottsche ex Steph. Accepted name: *C. obliqua* (Nees & Mont.) Schiffn. (Long & Grolle 1990).  
*Cololejeunea yulensis* (Steph.) Benedix. Accepted name: *C. aequabilis* (Sande Lac.) Schiffn. (Zhu 2005).  
*Drepanolejeunea formosana* Horik. — *Stenolejeunea apiculata* (Sande Lac.) R.M. Schust. (Zhu & So 2001). Accepted name: *Lejeunea apiculata* Sande Lac. (Söderström *et al.* 2010).  
*Drepanolejeunea subacuta* (Horik.) H.A. Miller *et al.* Accepted name: *Lejeunea exilis* (Reinw. *et al.*) Grolle (Zhu & Grolle 2003).  
*Eopleurozia giganteoides* (Horik.) Inoue. Accepted name: *Pleurozia subinflata* (Austin) Austin (Thiers 1993).  
*Folioceros vesiculosus* (Austin) D.C. Bharadwaj. Accepted name: *F. fuciformis* (Mont.) D.C. Bharadwaj (Hasegawa 1993).  
*Fossombronina australi-nipponica* Horik. Accepted name: *F. japonica* Schiffn. (Krayesky *et al.* 2005).  
*Frullania acutiloba* Mitt. var. *schiffneri* Verd. Accepted name: *F. monocera* (Hook.f. & Taylor) Taylor var. *schiffneri* (Verd.) S. Hatt. (Hattori 1979a).  
*Frullania caduca* S. Hatt. Accepted name: *F. amamiensis* Kamim. (Yuzawa 2001).  
*Frullania davurica* Hampe subsp. *jackii* (Gottsche) S. Hatt. Accepted name: *F. jackii* Gottsche (Grolle & Long 2000).  
*Frullania galeata* (Reinw. *et al.*) Dumort. Accepted name:

- F. riojaneirensis* (Raddi) Ångstr. (Svihla 1956, Hattori & Lin 1985).
- Frullania jackii* Gottsche subsp. *japonica* (Sande Lac.) S. Hatt. Accepted name: *F. davurica* Hampe (Hattori 1976a).
- Frullania jackii* Gottsche subsp. *japonica* (Sande Lac.) S. Hatt. var. *chichibuensis* Kamim. Accepted name: *F. davurica* Hampe var. *chichibuensis* (Kamim.) S. Hatt. (Hattori 1976a).
- Frullania kagoshimensis* Steph. var. *minor* Kamim. Accepted name: *F. osumiensis* (S. Hatt.) S. Hatt. (Yuzawa 2001).
- Frullania pedicellata* auct. non Steph. Accepted name: *F. muscicola* Steph. (Hattori & Lin 1985).
- Frullania tamarisci* subsp. *moniliata* (Reinw. et al.) Kamim. Accepted name: *F. moniliata* (Reinw. et al.) Mont. (Crandall-Stotler et al. 1987).
- Frullania tamarisci* ssp. *obscura* (Verd.) S. Hatt. Accepted name: *F. moniliata* (Reinw. et al.) Mont. (Crandall-Stotler et al. 1987).
- Frullania tamarisci* ssp. *obscura* var. *balansae* (Steph.) S. Hatt. Accepted name: *F. moniliata* (Reinw. et al.) Mont. (Crandall-Stotler et al. 1987).
- Gottschea aligera* (Nees & Blume) Nees. Accepted name: *Schistochila aligera* (Nees & Blume) J.B. Jack & Steph. (So 2003a).
- Gottschea nuda* (Horik.) Grolle & Zijlstra. Accepted name: *Schistochila nuda* Horik. (So 2003a).
- Gottschea philippinensis* Mont. Accepted name: *Schistochila aligera* (Nees & Blume) J.B. Jack & Steph. (So 2003a).
- Herbertus girdaldianus* (Steph.) Nichols. Accepted name: *H. dicranus* (Taylor ex Gottsche et al.) Trevis. (Feldberg & Heinrichs 2005).
- Herbertus longifolius* Horik. Accepted name: *H. dicranus* (Taylor ex Gottsche et al.) Trevis. (Inoue 1977).
- Herbertus pinnatus* (Steph.) H.A. Mill. Accepted name: *H. dicranus* (Taylor ex Gottsche et al.) Trevis. (Juslén 2006).
- Herbertus sakuraii* (Warnst.) S. Hatt. Accepted name: *H. dicranus* (Taylor ex Gottsche et al.) Trevis. (Hodgetts 2003).
- Heteroscyphus bescherellei* (Steph.) S. Hatt. Accepted name: *H. coalitus* (Hook.) Schiffn. (Inoue 1988).
- Jamesoniella autumnalis* (DC.) Steph. Accepted name: *Syzygiella autumnalis* (DC.) Feldberg et al. (Feldberg et al. 2010).
- Jamesoniella nipponica* S. Hatt. Accepted name: *Syzygiella nipponica* (S. Hatt.) Feldberg et al. (Feldberg et al. 2010).
- Jamesoniella pulchra* N. Kitag. Accepted name: *Denotarisia linguifolia* (De Not.) Grolle (Váňa 1991).
- Jubula hutchinsiae* (Hook.) Dumort. subsp. *javanica* (Steph.) Verd. Accepted name: *J. javanica* Steph. (Katagiri et al. 2010).
- Jungermannia appressifolia* Mitt. Accepted name: *Solenostoma appressifolium* (Mitt.) Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia comata* Nees. Accepted name: *Solenostoma comatum* (Nees) C. Gao (Váňa & Long 2009).
- Jungermannia exsertifolia* Steph. subsp. *cordifolia* (Dumort.) Váňa. Accepted name: *J. exsertifolia* Steph. (Schuster 1988).
- Jungermannia gracillima* Sm. Accepted name: *Solenostoma gracillimum* (Sm.) R.M. Schust. (Váňa & Long 2009).
- Jungermannia hasskarliana* (Nees) Steph. Accepted name: *Solenostoma hasskarlianum* (Nees) R.M. Schust. ex Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia hyalina* Lyell. Accepted name: *Solenostoma hyalinum* (Lyell) Mitt. (Váňa & Long 2009).
- Jungermannia infusca* (Mitt.) Steph. Accepted name: *Solenostoma infusum* (Mitt.) Hentschel (Hentschel et al. 2007a).
- Jungermannia pseudocyclops* Inoue. Accepted name: *Solenostoma pseudocyclops* (Inoue) Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia pyriflora* Steph. Accepted name: *Solenostoma pyriflorum* Steph. (Váňa & Long 2009).
- Jungermannia radiculosa* (Mitt.) Steph. Accepted name: *Solenostoma radiculosum* Mitt. (Váňa & Long 2009).
- Jungermannia schauliana* Steph. Accepted name: *Solenostoma schaulianum* (Steph.) Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia subrubra* Steph. Accepted name: *Solenostoma subrubrum* (Schiffn. ex Steph.) Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia subulata* A. Evans. Accepted name: *Liochlarena subulata* (A. Evans) Schljakov (Váňa & Long 2009).
- Jungermannia tetragona* Lindenb. Accepted name: *Solenostoma tetragonum* (Lindenb.) R.M. Schust. ex Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia truncata* Nees. Accepted name: *Solenostoma truncatum* (Nees) R.M. Schust. ex Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia virgata* (Mitt.) Steph. Accepted name: *Solenostoma virgatum* (Mitt.) Váňa & D.G. Long (Váňa & Long 2009).
- Jungermannia vulcanicola* (Schiffn.) Steph. Accepted name: *Solenostoma vulcanicola* (Schiffn.) Nyushko (Potemkin & Sofronova 2009).
- Lejeunea borneensis* Steph. — *Lejeunea catanduana* (Steph.) H.A. Mill. et al. (Kitagawa 1981). Accepted name: *L. anisophylla* Mont. (Hürlimann 1993).
- Lejeunea catanduana* (Steph.) H.A. Mill. et al. Accepted name: *L. anisophylla* Mont. (Hürlimann 1993).
- Lejeunea cucullata* (Reinw. et al.) Nees. Accepted name: *Metalejeunea cucullata* (Reinw. et al.) Grolle (Grolle 1995).
- Lejeunea obscura* Mitt. — *Taxilejeunea subcompressiuscula* Herzog, Journal of the Hattori Botanical Laboratory 14: 47. 1955, *nom. inval.*, *syn. nov.* Based on study of the original material, we conclude that this name is a synonym of *Lejeunea obscura*.
- Lejeunea proliferans* Herzog. Accepted name: *L. cocoes* Mitt. (So & Zhu 1998).
- Lejeunea vaginata* Steph. Accepted name: *L. discreta* Lindenb. (Mizutani 1970).
- Lepidozia setacea* auct. Accepted name: *Kurzia pauciflora* (Dicks.) Grolle (Grolle 1963).
- Lepidozia wallichiana* Gottsche. Accepted name: *Telaranea wallichiana* (Gottsche) R.M. Schust. (Schuster 1980).
- Leptolejeunea aberrantia* Horik. Accepted name: *L. truncati-*

- folia* Steph. (Zhu & So 2001).
- Leptolejeunea picta* Herzog. Accepted name: *Leptolejeunea amphiophthalmia* Zwichel (Schuster 1984).
- Leptolejeunea radiata* (Mitt.) Steph. Accepted name: *L. maculata* (Mitt.) Schiffn. (Grolle 1976).
- Leptolejeunea schiffneri* (Steph.) Steph. Accepted name: *L. maculata* (Mitt.) Schiffn. (Grolle 1976).
- Leptolejeunea subacuta* A. Evans. Accepted name: *L. elliptica* (Lehm. & Lindenb.) Schiffn. (Mizutani & Hattori 1969).
- Leptolejeunea yangii* M.J. Lai. Accepted name: *Drepanolejeunea foliicola* Horik. (Grolle & Zhu 2000).
- Leucolejeunea paroica* N. Kitag. Accepted name: *Cheilolejeunea kitagawae* W. Ye & R.L. Zhu (Ye & Zhu 2010).
- Leucolejeunea turgida* (Mitt.) Verd. Accepted name: *Cheilolejeunea turgida* (Mitt.) W. Ye & R.L. Zhu (Ye & Zhu 2010).
- Leucolejeunea xanthocarpa* (Lehm. & Lindenb.) A. Evans. Accepted name: *Cheilolejeunea xanthocarpa* (Lehm. & Lindenb.) Malombe (Malombe 2009).
- Lopholejeunea brunnea* Horik. Accepted name: *L. nigricans* (Lindenb.) Schiffn. (Amakawa 1970).
- Lopholejeunea javanica* (Nees) Schiffn. Accepted name: *L. nigricans* (Lindenb.) Schiffn. (Zhu & Gradstein 2005).
- Lopholejeunea muenensis* Steph. Accepted name: *L. nigricans* (Lindenb.) Schiffn. (Zhu & Gradstein 2005).
- Lopholejeunea nicobarica* Steph. Accepted name: *L. eulopha* (Taylor) Schiffn. (Zhu & Gradstein 2005).
- Lopholejeunea nipponica* Horik. Accepted name: *L. zollingeri* (Steph.) Schiffn. (Zhu & Gradstein 2005).
- Lophozia cornuta* (Steph.) S. Hatt. Accepted name: *Schistochilopsis cornuta* (Steph.) Konstant. (Konstantinova & Vasiljev 1994).
- Lophozia incisa* (Schrad.) Dumort. Accepted name: *Schistochilopsis incisa* (Schrad.) Konstant. (Konstantinova & Vasiljev 1994).
- Lophozia morrisoncola* Horik. Accepted name: *Leiocolea morrisoncola* (Horik.) Furuki & Mizut. (Furuki & Mizutani 2004).
- Lophozia nakanishii* Inoue. Accepted name: *Schistochilopsis nakanishii* (Inoue) Konstant. (Konstantinova & Vasiljev 1994).
- Macvicaria ulophylla* (Steph.) S. Hatt. Accepted name: *Porella ulophylla* (Steph.) S. Hatt. (Hentschel *et al.* 2007b).
- Mannia subpilosa* (Horik.) Horik. Accepted name: *Cryptomitrium tenerum* (Hook. ex Kunth) Austin ex Underw (Katagiri & Deguchi 2011).
- Marchantia cuneiloba* Steph. Accepted name: *M. emarginata* Reinw. *et al.* subsp. *tosana* (Steph.) Bischl. (Bischler 1989).
- Marchantia paleacea* Bertol. var. *diptera* (Nees & Mont.) S. Hatt. Accepted name: *M. paleacea* Bertol. subsp. *diptera* (Nees & Mont.) S. Hatt. ex Inoue (Inoue 1989).
- Marchantia tosana* Steph. Accepted name: *M. emarginata* Reinw. *et al.* subsp. *tosana* (Steph.) Bischl. (Bischler 1989).
- Marsupella emarginata* (Ehrh.) Dumort. subsp. *tubulosa* (Steph.) N. Kitag. Accepted name: *M. emarginata* (Ehrh.) Dumort. (Váňa *et al.* 2005).
- Marsupella revoluta* (Nees) Dumort. Accepted name: *Apomarsupella revoluta* (Nees) Dumort. (Schuster 1996).
- Marsupella rubida* (Mitt.) Grolle Accepted name: *Apomarsupella rubida* (Mitt.) R.M. Schust. (Schuster 1996).
- Mastigolejeunea humilis* (Gottsche) Schiffn. Accepted name: *M. auriculata* (Wilson) Schiffn. (Thiers & Gradstein 1989).
- Metacalypogeia quelpaertensis* S. Hatt. & Inoue. Accepted name: *Eocalypogeia quelpaertensis* (S. Hatt. & Inoue) R.M. Schust. (Schuster & Konstantinova 1996).
- Metzgeria decipiens* (C. Massal.) Schiffn. Accepted name: *M. furcata* (L.) Dumort. (Grolle 2002).
- Metzgeria fauriana* Steph. Accepted name: *M. furcata* (L.) Dumort. (So 2003b).
- Metzgeria fruticulosa* auct. non (Disks.) A. Evans. Accepted name: *M. temperata* Kuwah. (Kuwahara 1976).
- Metzgeria hamata* auct. Accepted name: *M. leptoneura* Spruce (Engel 1976).
- Microlejeunea lunulatiloba* Horik. — *Lejeunea lunulatiloba* (Horik.) Mizut. (Mizutani 1961). Accepted name: *L. ulicina* (Taylor) Gottsche *et al.* (Schuster 1962).
- Microlejeunea ramulosa* Herzog. — *Lejeunea ramulosa* (Herzog) R.M. Schust. (Schuster 1963). Accepted name: *L. laii* R.L. Zhu (Wang & Zhu 2008).
- Microlejeunea punctiformis* (Taylor) Spruce. Accepted name: *Lejeunea punctiformis* Taylor (Schuster 1962).
- Microlejeunea ulicina* (Taylor) A. Evans. Accepted name: *Lejeunea ulicina* (Taylor) Gottsche *et al.* (Inoue 1974).
- Mylia nuda* Inoue & B.Y. Yang. Accepted name: *M. verrucosa* Lindb. subsp. *nuda* (Inoue & B.Y. Yang) Potemkin & Kazanovsky (Potemkin & Kazanovsky 1993).
- Notoscyphus paroicus* Schiffn. Accepted name: *N. lutescens* (Lehm. & Lindenb.) Mitt. (Váňa & Inoue 1983).
- Notothylas japonica* Horik. Accepted name: *N. orbicularis* (Schwein.) Sull. ex A. Gray (Inoue 1976).
- Pallavicinia longispina* Steph. Accepted name: *P. subciliata* (Austin) Steph. (Inoue 1981).
- Phaeoceros laevis* (L.) Prosk. subsp. *carolinianus* (Michx.) Prosk. Accepted name: *P. carolinianus* (Michx.) Prosk. (Hasegawa 1991).
- Paraschistochila aligera* (Nees & Blume) R.M. Schust. — *Gottschea aligera* (Nees & Blume) Nees (Grolle & Zijlstra 1984). Accepted name: *Schistochila aligera* (Nees & Blume) J.B. Jack & Steph. (So 2003a).
- Paraschistochila philippinensis* (Mont.) R.M. Schust. — *Gottschea philippinensis* Mont. (Grolle & Zijlstra 1984). Accepted name: *Schistochila aligera* (Nees & Blume) J.B. Jack & Steph. (So 2003a).
- Phaeoceros miyakeanus* (Schiffn.) S. Hatt. Accepted name: *P. laevis* (L.) Prosk. (Hasegawa 1991).
- Plagiochila acanthophylla* Gottsche. Accepted name: *P. sciophila* Nees ex Lindenb. (Inoue 1982).
- Plagiochila eberhardtii* Steph. Accepted name: *P. obtusa* Lindenb. (Grolle & So 1999c).
- Plagiochila firma* var. *acicularis* (Herzog) Inoue. Accepted name: *P. gracilis* Lindenb. & Gottsche (Inoue 1984).
- Plagiochila gollanii* Steph. Accepted name: *P. nepalensis* Lindenb. (Grolle & So 1999d).
- Plagiochila griffithiana* Steph. Accepted name: *P. peculiaris*

Schiffn. (Grolle & So 1999c).  
*Plagiochila massalongoana* Schiffn. Accepted name: *P. junghuhniana* Sande Lac. (So & Grolle 1999).  
*Plagiochila sawadae* Inoue. Accepted name: *P. durelii* Schiffn. (Grolle & So 1999a).  
*Plagiochila sikutzuisana* C. Massal. Accepted name: *P. delavayi* Steph. (So 2001).  
*Plagiochila tobagensis* Herzog & S. Hatt. Accepted name: *P. kurzii* Steph. (Grolle & So 1999c).  
*Plagiochila vietnamica* Inoue. Accepted name: *P. durelii* Schiffn. (Grolle & So 1999a).  
*Plagiochila yokogurensis* Steph. Accepted name: *P. parvifolia* Lindenb. (Grolle & So 1999b).  
*Plectocolea crenulata* (Mitt.) A. Evans. — *Jungermannia gracillima* Sm. (Váňa 1974) Accepted name: *Solenostoma gracillimum* (Sm.) R.M. Schust. (Váňa & Long 2009).  
*Porella caespitans* var. *setigera* (Steph.) S. Hatt. Accepted name: *P. caespitans* var. *cordifolia* (Steph.) S. Hatt. (Hattori 1979b).  
*Porella gracillima* Mitt. var. *urogea* (C. Massal.) S. Hatt. Accepted name: *P. gracillima* Mitt. subsp. *urogea* (C. Massal.) S. Hatt. & M.X. Zhang (Hattori 1971).  
*Prionolejeunea ungulata* Herzog. — *Stenolejeunea apiculata* (Sande Lac.) R.M. Schust. (Zhu & So 2001). Accepted name: *Lejeunea apiculata* Sande Lac. (Söderström et al. 2010).  
*Pseudolepicolea trollii* subsp. *andoi* (R.M. Schust.) S. Hatt. & Mizut. Accepted name: *P. andoi* (R.M. Schust.) Inoue (Inoue 1978).  
*Radula apiculata* Sande Lac. ex Steph. Accepted name: *R. acuta* Mitt. (So 2006).  
*Radula kanemarui* S. Hatt. Accepted name: *R. tokiensis* Steph. (Hattori 1958).  
*Radula meyeri* Steph. Accepted name: *R. fulvifolia* (Hook.f. & Taylor) Gottsche et al. (Grolle 2001).  
*Radula retroflexa* Taylor var. *fauciloba* (Steph.) K. Yamada Accepted name: *R. retroflexa* Taylor (Yamada & Inoue 1988).  
*Radula valida* Steph. Accepted name: *R. perrottetii* Gottsche ex Steph. (Yamada 1975).  
*Radula variabilis* S. Hatt. Accepted name: *R. javanica* Gottsche (Yamada 1979).  
*Rectolejeunea barbata* Herzog. Accepted name: *Lejeunea barbata* (Herzog) R.L. Zhu & M.J. Lai (present paper).  
*Rhaphidolejeunea foliicola* (Horik.) P.C. Chen. Accepted name: *Drepanolejeunea foliicola* Horik. (Grolle & Zhu 2000).  
*Rhaphidolejeunea yunnanensis* P.C. Chen. Accepted name: *Drepanolejeunea yunnanensis* (P.C. Chen) Grolle & R.L. Zhu (Grolle & Zhu 2000).  
*Riccardia jackii* auct. non. Schiffn. Accepted name: *R. nagsakiensis* (Steph.) S. Hatt. (Furuki 1991).  
*Riccardia platyclada* Schiffn. Accepted name: *R. graeffei* (Steph.) Hewson (Furuki 1991).  
*Scapania glaucoviridis* Horik. Accepted name: *S. parvixtexta* Steph. (Potemkin 2002).  
*Scapania parvidens* Steph. Accepted name: *S. parvixtexta* Steph. (Potemkin 2002).  
*Scapania robusta* Horik. Accepted name: *S. subnimbosa*

Steph. (Potemkin 2002).  
*Scapania stephanii* Müll. Frib. Accepted name: *S. ligulata* subsp. *stephanii* (Müll. Frib.) Potemkin et al. (Potemkin et al. 2004).  
*Scaphophyllum speciosum* (Horik.) Inoue. Accepted name: *Solenostoma speciosum* (Horik.) Hentschel et al. (Feldberg et al. 2009).  
*Stenolejeunea apiculata* (Sande Lac.) R.M. Schust. Accepted name: *Lejeunea apiculata* Sande Lac. (Söderström et al. 2010).  
*Taeniolejeunea peraffinis* fo. *corticola* Herzog. Accepted name: *C. peraffinis* (Schiffn.) Schiffn. (present paper).  
*Taxilejeunea luzonensis* Steph. Accepted name: *Lejeunea luzonensis* (Steph.) R.L. Zhu & M.J. Lai (present paper).  
*Trichocolea lumbricoides* Horik. Accepted name: *T. merrillana* Steph. (Kitagawa 1973).

## Acknowledgements

Thanks go to the Early Land Plants Today project (M. von Konrat, L. Söderström, & A. Hagborg), Y.-C. Hsu, D.G. Long, J. Váňa and T. Yamaguchi for providing literature. This research was sponsored by the National Natural Science Foundation of China (nos. 30825004, 30670155), the Program of Shanghai Subject Chief Scientist (no. 08XD14016), 211 Project for the East China Normal University and PhD Program Scholarship Fund of ECNU 2009. The special thanks are due to J. Zündorf (JE) and M. Price (G) for making specimens including types available for this study, and to W. Frey for reading and correcting the earlier manuscript.

## References

- Ahonen, I. 2004: Molecular phylogeny of the liverwort order Porellales (Marchantiophyta, Jungermanniopsida). — *Monographs in Systematic Botany from the Missouri Botanical Garden* 98: 168–188.
- Amakawa, T. 1970: [On *Lopholejeunea brunnea*]. — *Miscellanea Bryologica et Lichenologica* 5: 103–104. [In Japanese].
- Bai, Z.-K. & Li, D.-K. 1998: A study on Pleuroziaceae (Hepaticae) of China. — *Acta Botanica Yunnanica* 20: 174–178.
- Bakalin, V. A. 2003: The status and treatment of the genus *Hattoriella* (H. Inoue) H. Inoue. — *Arctoa* 12: 91–96.
- Bapna, K. R. & Kachroo, P. 2000a: *Hepaticology of India I*. — Himanshu Publications, Udaipur.
- Bapna, K. R. & Kachroo, P. 2000b: *Hepaticology of India II*. — Himanshu Publications, Udaipur.
- Bharadwaj, D. C. 1975: Validation of some new combinations under *Folioceros* Bharadwaj. — *Geophytology* 5: 227–228.
- Bischler, H. 1989: *Marchantia* L. subg. *Chiamidium* (Nees) Bischl., sect. *Papiliatae* Bischl. sect. nov. en Asie et en Océanie. — *Cryptogamie: Bryologie, Lichénologie* 10: 61–79.



- Crandall-Stotler, B., Stotler, R. E. & Geissler, P. 1987: A biosystematic study of the subspecies of *Frullania tamarisci* (L.). — *The Bryologist* 90: 287–308.
- Crandall-Stotler, B., Stotler, R. E. & Long, D. G. 2009: Phylogeny and classification of the Marchantiophyta. — *Edinburgh Journal of Botany* 66: 155–198.
- Engel, J. J. 1976: *Metzgeria hamata* Lindb., an illegitimate name of Hepaticae. — *Lindbergia* 3: 219–220.
- Engel, J. J. & Braggins, J. E. 2005: Are *Mylia* and *Trabacellula* (Hepaticae) related? Unsuspected links revealed by cell wall morphology, with the transfer of *Mylia anomala* to a new genus (*Leiomylia* J.J. Engel & Braggins) of Jungermanniaceae. — *Taxon* 54: 665–680.
- Engel, J. J. & Merrill, G. L. S. 2004: Austral Hepaticae. 35. A taxonomic and phylogenetic study of *Telaranea* (Lepidoziaceae), with a monograph of the genus in temperate Australasia and commentary on extra-Australasian taxa. — *Fieldiana: Botany, New Series* 44: 1–265.
- Feldberg, K. & Heinrichs, J. 2005: On the identity of *Herbertus borealis* (Jungermanniopsida: Herbertaceae), with notes on the possible origin of *H. sendtneri*. — *Journal of Bryology* 27: 343–350.
- Feldberg, K., Vána, J., Hentschel, J. & Heinrichs, J. 2010: Currently accepted species and new combinations in Jamesonielloideae (Adelanthaceae, Jungermanniales). — *Cryptogamie Bryologie* 31: 141–146.
- Feldberg, K., Hentschel, J., Bombosch, A., Long, D. G., Vána, J. & Heinrichs, J. 2009: Transfer of *Gottschelia grollei*, *G. patoniae* and *Scaphophyllum speciosum* to *Solenostoma* based on chloroplast DNA *rbcL* sequences. — *Plant Systematics and Evolution* 280: 243–250.
- Frey, W., Stech, M. & Fischer, E. (eds.) 2009: *Syllabus of plant families — A. Engler's Syllabus der Pflanzenfamilien 3: Bryophytes and seedless vascular plants*. — Gebr. Bornträger Verlagsbuchhandlung, Berlin.
- Furuki, T. 1991: A taxonomical revision of the Aneuraceae (Hepaticae) of Japan. — *Journal of the Hattori Botanical Laboratory* 70: 293–397.
- Furuki, T. & Mizutani, M. 2004: [Checklist of Japanese Hepaticae and Anthocerotae, 2004]. — *Bryological Research* 8: 296–316. [In Japanese].
- Gao, C. (ed.) 2003: *Flora Bryophytorum Sinicorum* 9. — Science Press, Beijing. [In Chinese].
- Gao, C. & Bai, X.-L. 2001: A synoptic revision of family Jungermanniaceae (Hepaticae) in China including some taxa nova. — *The Philippine Scientist* 38: 111–170.
- Gao, C. & Cao, T. (eds.) 2000: *Flora Yunnanica* 17. — Science Press, Beijing. [In Chinese].
- Gao, C. & Lai, M.-J. 2003: [*Illustrations of Bryophytes of China*]. — SMC Publishing Inc., Taipei. [In Chinese].
- Gao, C. & Wu, Y.-H. 2004: On seven species of the Schistochilaceae (Hepaticae) in China, including one new species and one new combination. — *Journal of the Hattori Botanical Laboratory* 95: 263–270.
- Gao, C. & Wu, Y.-H. (eds.) 2008: *Flora Bryophytorum Sinicorum* 10. — Science Press, Beijing. [In Chinese].
- Gao, C., Cao, T. & Lai, M.-J. 2001: The genus *Saccogynidium* (Geocalyceae, Hepaticae) in China. — *The Bryologist* 104: 126–129.
- Gao, C., Cao, T. & Sun, J. 2002: The genus *Mnioloma* (Hepaticae, Calypogeiaceae) new to China discovered from Taiwan. — *Arctoa* 11: 23–26.
- Gao, C., Cao, T. & Sun, J. 2003 “2002”: The genus *Isotachis* (Hepaticae, Balantiopsaceae) in China. — *The Bryologist* 105: 693–698.
- Gao, C., Jia, X.-Y. & Cao, T. 1999: A taxonomic study of *Notoscyphus* Mitt. (Hepaticae) in China. — *Bulletin of Botanical Research* 19: 361–367.
- Gao, C., Cao, T., Wu, Y.-H., Yu, J. & Chen, Y. 2004: A new species and three new records of *Heteroscyphus* (Jungermanniopsida: Geocalyceae) to China. — *Journal of Bryology* 26: 97–102.
- Gradstein, S. R., He, X.-L., Piippo, S. & Mizutani, M. 2002: Bryophyte flora of the Huon Peninsula, Papua New Guinea. LXVIII. Lejeuneaceae subfamily Ptychanthoideae (Hepaticae). — *Acta Botanica Fennica* 174: 1–88.
- Grolle, R. 1963: Über *Kurzia* v. Martens. — *Revue Bryologique et Lichénologique* 32: 166–180.
- Grolle, R. 1966: Die Lebermoose Nepals. — *Khumbu Himal, Ergebnisse des Forschungsunternehmens Nepal Himalaya* 1: 262–298.
- Grolle, R. 1976: *Drepanolejeunea* subgen. *Kolpolejeunea* Eine neue Untergattung aus der Paläotropis. — *Journal of the Hattori Botanical Laboratory* 40: 191–216.
- Grolle, R. 1989: Miscellanea hepaticologica. 271–280. — *Journal of the Hattori Botanical Laboratory* 67: 249–254.
- Grolle, R. 1995: The Hepaticae and Anthocerotae of the East African Islands: An annotated catalogue. — *Bryophytorum Bibliotheca* 48: 1–178.
- Grolle, R. 2001: Miscellanea hepaticologica 291–300. — *Haussknechtia* 8: 59–69.
- Grolle, R. 2002: The Hepaticae and Anthocerotae of the subantarctic and temperate islands in the eastern southern hemisphere (90°E to 0°): an annotated catalogue. — *Journal of Bryology* 24: 57–80.
- Grolle, R. & Long, D. G. 2000: An annotated checklist of the Hepaticae and Anthocerotae of Europe and Macronesia. — *Journal of Bryology* 22: 103–140.
- Grolle, R. & Piippo, S. 1990: Bryophyte flora of the Huon Peninsula, Papua New Guinea. XXXVII. *Leucolejeunea* (Lejeuneaceae, Hepaticae). — *Annales Botanici Fennici* 27: 119–129.
- Grolle, R. & So, M. L. 1999a: On the *Plagiochila* species of sect. *Zonatae* with paraphyllia or mammillose stems (Hepaticae). — *Systematic Botany* 24: 297–310.
- Grolle, R. & So, M. L. 1999b: Studies of *Plagiochila* sect. *Subtropicae* in Asia. — *The Bryologist* 102: 67–75.
- Grolle, R. & So, M. L. 1999c: Studies on *Plagiochila* in China III. — *The Bryologist* 102: 200–207.
- Grolle, R. & So, M. L. 1999d: Studies on *Plagiochila* sect. *Contiguae* in East and South Asia. — *The Bryologist* 102: 294–303.
- Grolle, R. & Zhu, R.-L. 2000: A study of *Drepanolejeunea* subgen. *Rhaphidolejeunea* (Herzog) Grolle & R.L. Zhu, *stat. nov.* (Hepaticae, Lejeuneaceae) in China with notes on its species elsewhere. — *Nova Hedwigia* 70: 373–396.
- Grolle, R. & Zijlstra, G. 1984: On the nomenclature of *Gottschea* Nees *ex* Mont. 1843 (Schistochilaceae,

- Jungermanniales). — *Taxon* 33: 87–93.
- Hasegawa, J. 1986: The Anthocerotae collected by Drs. Z. Iwatsuki and N. Kitagawa in New Caledonia and Fiji. — *Journal of the Hattori Botanical Laboratory* 60: 379–390.
- Hasegawa, J. 1991: Taxonomy of *Phaeoceros laevis* subsp. *carolinianus* and its allied taxa in Japan and its adjacent region. — *Journal of the Hattori Botanical Laboratory* 69: 101–106.
- Hasegawa, J. 1992: [Distribution and taxonomic status of *Anthoceros formosae*]. — *Acta Phytotaxonomica et Geobotanica* 43: 44. [In Japanese].
- Hasegawa, J. 1993: Taxonomical studies on Asian Anthocerotae V. A short revision of Taiwanese Anthocerotae. — *Acta Phytotaxonomica et Geobotanica* 44: 97–112.
- Hattori, S. 1941: Notulae de Hepaticis Japonicis. I. — *Journal of Japanese Botany* 17: 457–466.
- Hattori, S. 1942a: Notulae de Hepaticis Japonicis. II. — *Journal of Japanese Botany* 18: 66–77.
- Hattori, S. 1942b: Notulae de Hepaticis Japonicis. III. — *Journal of Japanese Botany* 18: 469–477.
- Hattori, S. 1942c: Notulae de Hepaticis Japonicis. IV. — *Journal of Japanese Botany* 18: 653–660.
- Hattori, S. 1943: Hepaticarum species novae et minus cognitae nipponenses I. — *Botanical Magazine* 57: 357–363.
- Hattori, S. 1944: Contributio ad Floram Hepaticarum Austro-Kiushiensem. — *Bulletin of the Tokyo Science Museum* 11: 1–203.
- Hattori, S. 1948: Notulae de Hepaticis Japonicis. X. — *Journal of Japanese Botany* 22: 107–111.
- Hattori, S. 1951a: Notulae de Hepaticis Japonicis. 13. — *Journal of Japanese Botany* 26: 179–183.
- Hattori, S. 1951b: Oil bodies of Japanese Hepaticae (1). — *Journal of the Hattori Botanical Laboratory* 5: 69–97.
- Hattori, S. 1958: The Hepaticae of Ontake Mountain, middle Japan. — *Journal of the Hattori Botanical Laboratory* 28: 33–53.
- Hattori, S. 1967: Studies of the Asiatic species of the genus *Porella* (Hepaticae). I. Some new or little known Asiatic species of *Porella*. — *Journal of the Hattori Botanical Laboratory* 30: 129–151.
- Hattori, S. 1970: Studies of the Asiatic species of the genus *Porella* (Hepaticae). III. — *Journal of the Hattori Botanical Laboratory* 33: 41–87.
- Hattori, S. 1971: Studies of the Asiatic species of the genus *Porella* (Hepaticae). IV. — *Journal of the Hattori Botanical Laboratory* 34: 411–428.
- Hattori, S. 1976a: Notes on the Asiatic species of the genus *Frullania*, Hepaticae. IX. — *Bulletin of the National Science Museum B* 2: 7–22.
- Hattori, S. 1976b: Studies on the Asiatic species of the genus *Porella* (Hepaticae). VI. — *Journal of the Hattori Botanical Laboratory* 40: 121–138.
- Hattori, S. 1978: Studies on the Asiatic species of the genus *Porella* (Hepaticae). VII. A synopsis of Asiatic Porellaceae. — *Journal of the Hattori Botanical Laboratory* 44: 91–120.
- Hattori, S. 1979a: A revision of the Australasian species of the genus *Frullania*, Hepaticae. II. — *Journal of the Hattori Botanical Laboratory* 46: 119–153.
- Hattori, S. 1979b: [Corrections of names of *Porella* taxa]. — *Miscellanea Bryologica et Lichenologica* 8: 79. [In Japanese].
- Hattori, S. 1980a: A revision of the subgenus *Homotropantha* of the genus *Frullania*, Hepaticae. — *Journal of the Hattori Botanical Laboratory* 47: 165–236.
- Hattori, S. 1980b: Dr. H. Inoue's *Frullania* collections made in Formosa. — *Bulletin of the National Science Museum B* 6: 33–40.
- Hattori, S. & Lin, P.-J. 1985: A preliminary study of Chinese *Frullania* flora. — *Journal of the Hattori Botanical Laboratory* 59: 123–169.
- He, X.-L. 1997: A review and checklist of the Lejeuneaceae in China. — *Abstracta Botanica* 21: 69–77.
- Hentschel, J., Paton, J. A., Schneider, H. & Heinrichs, H. 2007a: Acceptance of *Liochlaena* Nees and *Solenostoma* Mitt., the systematic position of *Eremonotus* Pearson and notes on *Jungermannia* L. s.l. (Jungermanniidae) based on chloroplast DNA sequence data. — *Plant Systematics and Evolution* 268: 147–157.
- Hentschel, J., Zhu, R.-L., Long, D. G., Davison, P. G., Schneider, H., Gradstein, S. R. & Heinrichs, J. 2007b: A phylogeny of *Porella* (Porellaceae, Jungermanniopsida) based on nuclear and chloroplast DNA sequences. — *Molecular Phylogenetics and Evolution* 45: 693–705.
- Herzog, T. & Noguchi, A. 1955: Beitrag zur Kenntnis der Bryophytenflora von Formosa und den benachbarten Inseln Botel Tabago und Kwashyoto. — *Journal of the Hattori Botanical Laboratory* 14: 29–70.
- Higuchi, M. & Lin, S.-H. 2004: Correlations between the Moss Floras of Japan and Taiwan. — *National Science Museum Monographs* 24: 227–240.
- Hodgetts, N. G. 2003: Some synonyms of *Herbertus dicranus* (Taylor ex Gottsche et al.) Trevis. — *Journal of Bryology* 25: 138–140.
- Hong, W. S., Deffinbaugh, B. & Sparrow, B. 1993: The genus *Herbertus* in western North America. — *Lindbergia* 18: 41–45.
- Horikawa, Y. 1932: Studies on the Hepaticae of Japan. VI. — *Journal of Science of Hiroshima University B, Division 2 (Botany)* 1: 77–94.
- Horikawa, Y. 1934a: Monographia Hepaticarum Australi-Japonicarum. — *Journal of Science of Hiroshima University B, Division 2 (Botany)* 2: 101–325.
- Horikawa, Y. 1934b: Symbolae Florae Bryophytae Orientali-Asiae et Micronesiae. II. — *Hikobia* 48: 599–609.
- Horikawa, Y. 1935: Symbolae Florae Bryophytae Orientali-Asiae et Micronesiae. VI. — *Hikobia* 48: 588–595.
- Horikawa, Y. 1936: Symbolae Florae Bryophytae Orientali-Asiae et Micronesiae. X. — *Hikobia* 50: 556–561.
- Horikawa, Y. 1939: Contributions to the bryological flora of Eastern Asia VI. — *Journal of Japanese Botany* 15: 359–368.
- Horikawa, Y. 1951: Symbolae Florae Bryophytae Orientali-Asiae et Micronesiae. XII. — *Hikobia* 1: 78–100.
- Hsu, Y.-C. & Huang, Y.-C. 2010: [A preliminary examination of the specimens of lichen and bryophyte collected by professor Ming-Jou Lai]. — *Journal of the National Taiwan Museum* 63: 47–82. [In Chinese].
- Huang, S.-F. & Lin, S.-H. 2009: Materials to the bryophyte

- flora of Taiwan (I) — *Cephaloziella microphylla* (Steph.) Douin (Cephaloziellaceae). — *Taiwania* 54: 97–99.
- Huang, S.-F., Wu, Y.-C. & Lin, S.-H. 2009: Materials to the bryophyte flora of Taiwan (II) — *Fossombronina calcedonica* Steph. (Fossombroniaceae). — *Taiwania* 54: 187–189.
- Hürlimann, H. 1993: Hepaticae aus dem Gebiete des Südlichen Pazifik XII. — *Bauhinia* 11: 3–17.
- Inoue, H. 1959: A review of Japanese species of *Lophocolea* Dumort. — *Journal of the Hattori Botanical Laboratory* 21: 214–230.
- Inoue, H. 1967: *Apotreubia nana* from Taiwan with reference to the chromosome number. — *Bulletin of the National Science Museum* 10: 365–367.
- Inoue, H. 1974: [Illustrations of Japanese Hepaticae 1]. — Tsukiji Shokan, Tokyo. [In Japanese].
- Inoue, H. 1976: [Illustrations of Japanese Hepaticae 2]. — Tsukiji Shokan, Tokyo. [In Japanese].
- Inoue, H. 1977: Studies on Taiwan Hepaticae. II. Herbertaceae. — *Bulletin of the National Science Museum B* 3: 1–11.
- Inoue, H. 1978: Studies on Taiwan Hepaticae. III. Subord. Herbertinae and subord. Ptilidiinae. — *Bulletin of the National Science Museum B* 4: 93–103.
- Inoue, H. 1981: Taxonomic miscellany on hepatics (1). — *Journal of Japanese Botany* 56: 105–110.
- Inoue, H. 1982: Studies on Taiwan Hepaticae. IV. Plagiochilaceae. — *Bulletin of the National Science Museum B* 8: 125–144.
- Inoue, H. 1984: *The genus Plagiochila (Dum.) Dum. in Southeast Asia*. — Academic Scientific Book Inc., Tokyo.
- Inoue, H. 1985: Studies on Taiwan Hepaticae. VI. Schistochilaceae. — *Bulletin of the National Science Museum B* 11: 109–118.
- Inoue, H. 1988: Taxonomic miscellany on hepatics. 5. — *Journal of Japanese Botany* 63: 393–398.
- Inoue, H. 1989: Taxonomic miscellany on hepatics. 6. — *Journal of Japanese Botany* 64: 193–198.
- Inoue, H. & Yang, B.-Y. 1966: The genus *Mylia* Gray in Taiwan. — *Taiwania* 12: 35–37.
- Iwatsuki, Z. & Mizutani, M. 1972: *Coloured illustrations of bryophytes of Japan*. — Hoikusha Publishing Co., Ltd., Osaka.
- Juslén, A. 2004: Bryophyte flora of Hunan Province, China. 7. *Herbertus* (Herbertaceae, Hepaticae). — *Annales Botanici Fennici* 41: 393–404.
- Juslén, A. 2006: Revision of Asian *Herbertus* (Herbertaceae, Marchantiophyta). — *Annales Botanici Fennici* 43: 409–436.
- Kamimura, M. 1961: A monograph of Japanese Frullaniaceae. — *Journal of the Hattori Botanical Laboratory* 24: 1–108.
- Katagiri, T., Yamaguchi, T. & Deguchi, H. 2010: Taxonomic studies on the liverwort *Jubula* Dumort. (Marchantiophyta: Jubulaceae) in Japan and Taiwan. — *Hikobia* 15(4): 463–472.
- Katagiri, T. & H. Deguchi. 2011: *Mannia subpilosa* (Horik.) Horik. a new synonym of *Cryptomitrium tenerum* (Hook.) Austin. — *Tropical Bryology* 33: 1–5.
- Kitagawa, N. 1973: Miscellaneous notes on little-known species of Hepaticae, 26–50. — *Journal of the Hattori Botanical Laboratory* 37: 263–273.
- Kitagawa, N. 1981: Miscellaneous notes on little-known species of Hepaticae, 51–70. — *Hikobia* suppl. 1: 67–72.
- Konstantinova, N. A. & Vasiljev, A. N. 1994: On the hepatic flora of Sayan Mountains (South Siberia). — *Arctoa* 3: 123–132.
- Krayesky, D. M., Crandal-Stotler, B. & Stotler, R. E. 2005: A revision of the genus *Fossombronina* Raddi in East Asia and Oceania. — *Journal of the Hattori Botanical Laboratory* 98: 1–45.
- Kuwahara, Y. 1976: *Metzgeria temperata*, a new holarctic species of Hepaticae. — *Journal of the Hattori Botanical Laboratory* 40: 217–220.
- Kuo, C.-M. & Chiang, T.-Y. 1988: Index of Taiwan Hepaticae. — *Taiwania* 33: 1–46.
- Lai, M.-J. 1977: Bryoflora of Yuanyang Lake Natural Reserve, Taiwan. — *The Bryologist* 80: 153–155.
- Lai, M.-J. 1990: [Investigation of the bryophyte and lichen plant resource of Yangmingshan National Park]. — Yangmingshan National Park, Taipei. [In Chinese].
- Lai, M.-J. 1995: [Handbook of Bryophytic study]. — Administration of the Experimental Forest of National Taiwan University, Nantou. [In Chinese].
- Lai, M.-J. 2002: *Bazzania vittata* found in Taiwan. — *Taiwania* 23: 75–76.
- Lai, M.-J. 2003: [The flora and vegetation of Taiwan]. — Morning Star Publishing Inc., Taizhong. [In Chinese].
- Lai, M.-J. 2007: Study of bryophytes in Taiwan (I). — *National Taiwan Museum Special Publication Series* 12: 1–201.
- Lai, M.-J. & Wang-Yang, J.-R. 1976: Index bryoflorae formosensis. — *Taiwania* 21: 159–203.
- Lai, M.-J., Zhu, R.-L. & Chantanaorrapint, S. 2008: Liverworts and hornworts of Thailand: an updated checklist and bryofloristic accounts. — *Annales Botanici Fennici* 45: 321–341.
- Li, D.-K. 1990: [Preliminary report on the epiphyllous liverworts in Hunan Province, China]. — *Investigatio et Studium Naturae* 10: 137–138. [In Chinese].
- Lin, S.-H. 1981: Exsiccatae of the bryophytes of Taiwan. — *The Bryologist* 84: 359–362.
- Lin, S.-H. 1990: [Investigation of the bryophyte of Yushan National Park I]. — Yushan National Park Administration, Department of construction, Ministry of Interior, Nantou. [In Chinese].
- Lin, S.-H. 2000: [The liverwort flora of Taiwan]. — The Council of Agriculture, The Executive Yuan, Taipei. [In Chinese].
- Lin, S.-H. & Chen, Y.-Y. 1997: A taxonomic study of Frullaniaceae from Taiwan. — *Tunghai Journal* 38: 79–125.
- Lin, S.-H. & Yang, C.-S. 1992: Bryophytes of Chaishan, Taiwan. — *Yushania* 9: 1–12.
- Lin, P.-J., Piippo, S., Koponen, T. & Wu, P.-C. 1992: Bryophyte flora of Jianfengling Mts., Hainan Island, China. — *Bryobrothera* 1: 195–214.
- Long, D. G. 2005: Notes on Himalayan Hepaticae 2: New records and extensions of range for some Himalayan leafy liverworts. — *Cryptogamie Bryologie* 26: 97–107.

- Long, D. G. 2006: Revision of the genus *Asterella* P. Beauv. in Eurasia. — *Bryophytorum Bibliotheca* 63: 1–299.
- Long, D.G. & Grolle, R. 1990: Hepaticae of Bhutan II. — *Journal of the Hattori Botanical Laboratory* 68: 381–440.
- Malombe, I. 2009: Studies on African *Cheilolejeunea* (Lejeuneaceae) I: New species and new combinations. — *Acta Botanica Hungarica* 51: 315–328.
- Mizutani, M. 1961: A revision of Japanese Lejeuneaceae. — *Journal of the Hattori Botanical Laboratory* 24: 115–302.
- Mizutani, M. 1964: Studies of little known Asiatic species of Hepaticae in the Stephani herbarium. 1. On some little known Southeast Asiatic species of the family Lejeuneaceae. — *Journal of the Hattori Botanical Laboratory* 27: 139–148.
- Mizutani, M. 1967: Studies of little known Asiatic species of Hepaticae in the Stephani herbarium. 3. On some little known species of *Cheilolejeunea*, *Euosmolejeunea* and *Pycnolejeunea*. — *Journal of the Hattori Botanical Laboratory* 30: 171–180.
- Mizutani, M. 1968: Studies of little known Asiatic species of Hepaticae in the Stephani herbarium. 4. *Phaeolejeunea*, a new genus of Lejeuneaceae. — *Journal of the Hattori Botanical Laboratory* 31: 152–158.
- Mizutani, M. 1970: Lejeuneaceae, subfamilies Lejeuneoideae and Cololejeuneoideae from Sabah (North Borneo). — *Journal of the Hattori Botanical Laboratory* 33: 225–265.
- Mizutani, M. & Hattori, S. 1969: [Checklist of Japanese Hepaticae and Anthocerotae]. — *Miscellanea Bryologica et Lichenologica* 5: 33–43. [In Japanese].
- Nakanishi, S. 1964: Epiphytic communities in the alpine zone of Mt. Yushan (Mt. Niitaka), Taiwan. — *Miscellanea Bryologica et Lichenologica* 3: 71–73.
- Okamura, S. 1916: Contributiones novae ad Floram Bryophyton Japonicam. II. — *Journal of the College of Science, Imperial University of Tokyo* 38: 1–100.
- Paul, C., König, W. A. & Wu, C.-L. 2001: Sesquiterpenoid constituents of the liverworts *Lepidozia fauriana* and *Lepidozia vitrea*. — *Phytochemistry* 58: 789–798.
- Piippo, S. 1990: Annotated catalogue of Chinese Hepaticae and Anthocerotae. — *Journal of the Hattori Botanical Laboratory* 68: 1–192.
- Piippo, S. 1993: Bryophyte flora of the Huon Peninsula, Papua New Guinea. LIV. Anthocerotophyta. — *Acta Botanica Fennica* 148: 27–51.
- Piippo, S. 1996: Notes on Chinese Geocalycaceae (Hepaticae). 1. — *Annales Botanici Fennici* 33: 45–49.
- Piippo, S., He, X.-L. & Koponen, T. 1997: Hepatics from northwestern Sichuan, China, with a checklist of Sichuan hepatics. — *Annales Botanici Fennici* 34: 51–63.
- Piippo, S., He, X.-L., Juslén, A., Tan, B. C., Murphy, D. H. & Pócs, T. 2002: Hepatic and hornwort flora of Singapore. — *Annales Botanici Fennici* 39: 101–127.
- Pócs, T. & Bernecker, A. 2009: Overview of *Aphanolejeunea* (Jungermanniopsida) after 25 years. — *Polish Botanical Journal* 54: 1–11.
- Potemkin, A. D. 1999: On the evolution and classification of *Scapania* (Hepaticae) — subgenus *Plicatocalyx*. — *Annales Botanici Fennici* 36: 127–136.
- Potemkin, A. D. 2002: Phylogenetic system and classification of the family Scapaniaceae Mig. emend. Potemkin (Hepaticae). — *Annales Botanici Fennici* 39: 309–334.
- Potemkin, A. D. & Kazanovsky, S. G. 1993: On the genus *Mylia* S. Gray (Hepaticae, Jungermanniaceae, Mylioidae). — *Arctoa* 2: 1–11.
- Potemkin, A. D. & Sofronova, E. V. 2009: *Liverworts and Hornworts of Russia, vol. 1.* — Boston-Spectr, St. Petersburg. [In Russian with English summary].
- Potemkin, A. D., Piippo, S. & Koponen, T. 2004: Bryophyte flora of Hunan Province, China. 4. Diplophyllaceae and Scapaniaceae (Hepaticae). — *Annales Botanici Fennici* 41: 415–427.
- Schuster, R. M. 1962: North American Lejeuneaceae VIII. *Lejeunea*, subgenera *Microlejeunea* and *Chaetolejeunea*. — *Journal of the Hattori Botanical Laboratory* 25: 1–80.
- Schuster, R. M. 1963: An annotated synopsis of the genera and subgenera of Lejeuneaceae. I. Introduction; annotated keys to subfamilies and genera. — *Beihefte zur Nova Hedwigia* 9: 1–203.
- Schuster, R. M. 1980: New combinations and taxa of Hepaticae. I. — *Phytologia* 45: 415–437.
- Schuster, R. M. 1984: Diagnoses of some new taxa of Hepaticae. — *Phytologia* 56: 65–74.
- Schuster, R. M. 1988: The Hepaticae of South Greenland. — *Beihefte zur Nova Hedwigia* 92: 1–255.
- Schuster, R. M. 1996: Studies on Antipodal Hepaticae. XII. Gymnomitriaceae. — *Journal of the Hattori Botanical Laboratory* 80: 1–147.
- Schuster, R. M. 1998: On the genus *Scaphophyllum* (Jungermanniaceae). — *The Bryologist* 101: 428–434.
- Schuster, R. M. 2002: Revisionary studies of the Chandonanthoideae (Jungermanniales, Jungermanniaceae). — *Nova Hedwigia* 74: 465–496.
- Schuster, R. M. & Konstantinova, N. A. 1996: Studies on the distribution of critical arctic/subarctic Hepaticae with special reference to taxa found in Russia. — *Lindbergia* 21: 26–48.
- Shi, R.-P. 2008: [Species diversity and distribution of liverworts and hornworts in Hainan Island, China]. — M.Sc. thesis, East China Normal University, Shanghai. [In Chinese].
- So, M. L. 1995: Hornworts of Hong Kong. — *Hikobia* 12: 19–22.
- So, M. L. 2001: *Plagiochila* (Hepaticae, Plagiochilaceae) in China. — *Systematic Botany Monographs* 60: 1–214.
- So, M. L. 2002: The genus *Porella* (Porellaceae, Hepaticae) in Australasia and the South Pacific. — *Systematic Botany* 27: 4–13.
- So, M. L. 2003a: The genus *Schistochila* in Asia. — *Journal of the Hattori Botanical Laboratory* 93: 79–100.
- So, M. L. 2003b: The genus *Metzgeria* (Hepaticae) in Asia. — *Journal of the Hattori Botanical Laboratory* 94: 159–177.
- So, M. L. 2003c: The genus *Herbertus* (Hepaticae) in Australasia and the South Pacific. — *Systematic Botany* 28: 12–23.
- So, M. L. 2004: Studies on the type specimens of *Odonto-*



- schisma* (Hepaticae) in Asia and the Pacific. — *Journal of the Hattori Botanical Laboratory* 95: 249–261.
- So, M. L. 2006: *Radula* (Radulaceae, Marchantiophyta) in the South Pacific. — *Journal of the Hattori Botanical Laboratory* 99: 207–232.
- So, M. L. & Grolle, R. 1999: Studies on *Plagiochila* in Asia: Supplements to sections *Abietinae*, *Annotinae*, *Ciliatae*, *Contiguae*, *Cucullatae*, *Poeltiae*, *Subtropicae* and *Zonatae*. — *Cryptogamie Bryologie* 20: 167–179.
- So, M. L. & Zhu, R.-L. 1996a: Studies on Hong Kong *Cheilolejeunea* with two species new to China. — *Tropical Bryology* 12: 5–10.
- So, M. L. & Zhu, R.-L. 1996b: Studies on Hong Kong hepatics II. Notes on some newly recorded liverworts from Hong Kong. — *Tropical Bryology* 12: 11–19.
- So, M. L. & Zhu, R.-L. 1998: On six species of the genus *Lejeunea* in China, including one new species. — *The Bryologist* 101: 137–143.
- Söderström, L., Gradstein, S. R. & Hagborg, A. 2010: Checklist of the hornworts and liverworts of Java. — *Phytotaxa* 9: 53–149.
- Stephani, F. 1897: Hepaticae Japonicae. — *Bulletin de l'Herbier Boissier* 5: 76–108.
- Stephani, F. 1900–1924: *Species Hepaticarum*, 1–6. — Georg et Cie, même Maison, Genève et Bale, Lyon.
- Stotler, R. E. & Crandall-Stotler, B. J. 2003: (1596–1597) Proposals to conserve the name *Anthoceros* with a conserved type and the name *Anthoceros agrestis* against *A. nagsakiensis* (Anthocerotophyta). — *Taxon* 52: 628–629.
- Stotler, R. E. & Crandall-Stotler, B. 2005: A revised classification of the Anthocerotophyta and a checklist of the hornworts of North America, North of Mexico. — *The Bryologist* 108: 16–26.
- Stotler, R. E. & Croz, D. K. 1983: On *Mnium trichomanis* Linnaeus (Hepatophyta). — *Taxon* 32: 74–75.
- Svihla, D. E. 1956: The Frullaniaceae of Burma. I. — *The Bryologist* 59: 277–280.
- Tan, B. C. & Engel, J. J. 1986: An annotated checklist of Philippine Hepaticae. — *Journal of the Hattori Botanical Laboratory* 60: 283–355.
- Thiers, B. M. 1993: A monograph of *Pleurozia* (Hepaticae; Pleuroziaceae). — *The Bryologist* 96: 517–554.
- Thiers, B. M. & Gradstein, S. R. 1989: Lejeuneaceae (Hepaticae) of Australia. I. Subfamily Ptychanthoideae. — *Memoirs of the New York Botanical Garden* 52: 1–82.
- Tixier, P. 1971: A contribution to bryological knowledge of Fraser's Hill Station (Malaysia). — *The Gardens' Bulletin Singapore* 25: 335–353.
- Urmi, E. 1983: *Tetralophozia filiformis* (Steph.) *comb. nov.* in Europe. — *Journal of Bryology* 12: 393–401.
- van Balgooy, M. M. J. 1971: Plant geography of the Pacific as based on the distribution of phanerogam genera. — *Blumea Supplement* 6: 1–222.
- Váňa, J. 1974: Studien über die Jungermannioideae (Hepaticae). 6. *Jungermannia* subg. *Solenostoma*: Europäische und nordamerikanische Arten. — *Folia Geobotanica et Phytotaxonomica* 9: 369–423.
- Váňa, J. 1988: *Cephalozia* (Dum.) Dum. in Africa, with notes on the genus (Notes on some African hepatic genera 10). — *Beihfte zur Nova Hedwigia* 90: 179–198.
- Váňa, J. 1991: The bryophytes of Sabah (North Borneo) with special reference to the BRYOTROP transect of Mount Kinabalu. XV. Lophoziaaceae (Hepaticae). — *Willdenowia* 20: 199–219.
- Váňa, J. 1993: The bryophytes of Sabah (North Borneo) with special reference to the BRYOTROP transect of Mount Kinabalu. XVIII. Cephaloziaaceae (Hepaticopsida, Jungermanniales). — *Willdenowia* 23: 245–255.
- Váňa, J. & Inoue, H. 1983: Studies in Taiwan Hepaticae. V. Jungermanniaceae. — *Bulletin of the National Science Museum B* 9: 125–142.
- Váňa, J. & Long, D. G. 2009: Jungermanniaceae of the Sino-Himalayan region. — *Nova Hedwigia* 89: 485–517.
- Váňa, J., Piippo, S. & Koponen, T. 2005: Bryophyte flora of Hunan Province, China. 6. Jungermanniaceae and Gymnomitriaceae (Hepaticae). — *Acta Botanica Fennica* 178: 57–78.
- Verdoorn, F. 1934: Studien über Asiatische Jubuleae (De Frullaniaceis XV–XVII) mit einer Einleitung Bryologie und Hepaticologie Ihre Methodik und Zukunft. — *Annales Bryologici, Supplement* 4: 1–231.
- Wang, J. & Zhu, R.-L. 2008: *Lejeunea laii* *nom. nov.* for *Lejeunea ramulosa* (Herzog) R.M. Schust. (Jungermanniopsida: Lejeuneaceae) from Taiwan. — *Journal of Bryology* 30: 173–174.
- Wang, M.-Z. 1993: A preliminary study of Hepaticae from Mt. Jiufu of Guangxi, south China. — *Chenia* 1: 125–131.
- Wei, H.-C. & Wu, C.-L. 1991: A new macrocyclic bis-benzyl diether, marchantiquinone, from the liverwort *Mannia subpilosa*. — *Journal of Chemical Research, Synopses* 130–231.
- Wilson, R., Gradstein, S. R., Heinrichs, J., Groth, H., Ilkiu-Borges, A. L. & Hartmann, F. A. 2004: Phylogeny of Lejeuneaceae: A cladistic analysis of chloroplast gene *rbcl* sequences and morphology with preliminary comments on the mitochondrial NAD4-2 spacer region. — *Monographs in Systematic Botany from the Missouri Botanical Garden* 98: 189–202.
- Wu, C.-L. & Jong, J. R. 2001: A Cyclic Peroxide of Clerodenoic Acid from the Taiwanese liverwort *Schistochila acuminata*. — *Journal of Asian Natural Products Research* 3: 241–246.
- Wu, C.-L. & Yamada, K. 1996: *Telaranea neesii* (liverwort) new to Taiwan. — *Proceedings of the Bryological Society of Japan* 6: 246.
- Wu, C.-L. & Yamada, K. 1997: *Leucolejeunea turgida* (Mitt.) Verd., new to the hepatic flora of Taiwan. — *Bryological Research* 7: 26.
- Wu, C.-L. & Yamada, K. 2000: Two species of the genus *Kurzia* new to Taiwan. — *Bryological Research* 7: 349.
- Wu, P.-C. (ed.) 2000: [*Bryoflora of Hengduan Mts. (Southwest China)*]. — Science Press, Beijing. [In Chinese].
- Wu, P.-C. & Wang, M.-Z. 2001: Relationship of the tropical elements of the bryophytes between Hengduan Mts., southwest China and Taiwan province, southeast China. — *Guizhou Science* 19: 6–9. [In Chinese with English abstract].
- Wu, P.-C., Jia, Y. & Wang, M.-Z. 2002: The bryological

- relationship between the tropical bryophytes of Taiwan, southeast China and Hengduan Mts., southwest China. — *Chenia* 7: 187–191.
- Wu, Y.-H. & Gao, C. 2008: *Chiloscyphus sikkimensis* (Geocalycaceae, Hepaticae), new to Mainland of China. — *Bulletin of Botanical Research* 28: 520–522. [In Chinese].
- Yamada, K. 1975: Memoranda on the type specimens of *Radula* taxa from the southeast Asia (2). — *Journal of the Hattori Botanical Laboratory* 39: 71–84.
- Yamada, K. 1979: A revision of Asian taxa of *Radula*, Hepaticae. — *Journal of the Hattori Botanical Laboratory* 45: 201–322.
- Yamada, K. 1981: Notes on the type specimens of *Radula* taxa from Latin America. 2. — *Journal of the Hattori Botanical Laboratory* 49: 385–398.
- Yamada, K. & Inoue, H. 1988: Studies on Taiwan Hepaticae. VII. Radulaceae. — *Bulletin of the National Science Museum B* 14: 41–51.
- Yamada, K. & Iwatsuki, Z. 2006: Catalog of the Hepatics of Japan. — *Journal of the Hattori Botanical Laboratory* 99: 1–106.
- Yamada, K. & Lai, M.-J. 1979: Five *Bazzania* species new to Taiwan. — *Miscellanea Bryologica et Lichenologica* 8: 87.
- Yang, B.-Y. 1960a: Studies on Taiwan Hepaticae. I. A preliminary list of the Hepaticae of Taiwan. — *Quarterly Journal of the Taiwan Museum* 13: 231–235.
- Yang, B.-Y. 1960b: Studies of Taiwan Hepaticae II. *Radula* species collected in Taiwan. — *Taiwania* 7: 35–38.
- Yang, B.-Y. 1963: Further studies on Taiwan Hepaticae. The genus *Thysananthus*. — *Taiwania* 9: 23–31.
- Yang, B.-Y. 1965: The discovery of *Monosolenium tenerum* in Taiwan. — *Taiwania* 11: 29–34.
- Yang, B.-Y. 1966: The geographical distribution and growth habits of *Haplomitrium*. — *Taiwania* 12: 9–20.
- Yang, J.-D. 2009: *Liverworts and hornworts of Taiwan I. Lejeuneaceae*. — Endemic Species Research Institute, Nantou.
- Yang, J.-D. & Lin, S.-H. 2008a: *Cheilolejeunea falsinervis* (Marchantiophyta) newly recorded for Taiwan. — *Bryological Research* 9: 319–321.
- Yang, J.-D. & Lin, S.-H. 2008b: *Lejeunea bidentula* Herzog a newly reported liverwort to Taiwan. — *Endemic Species Research* 10: 65–69.
- Yang, J.-D. & Lin, S.-H. 2008c: *Leptolejeunea picta* Herzog, a liverwort new to Taiwan. — *Taiwania* 53: 308–310.
- Yang, J.-D. & Lin, S.-H. 2009a: *Calypogeia aeruginosa* Mitt., a newly recorded liverwort to Taiwan. — *Endemic Species Research* 11: 93–99.
- Yang, J.-D. & Lin, S.-H. 2009b: *Lejeunea sordida* (Marchantiophyta) newly recorded for Taiwan. — *Bryological Research* 9: 388–390.
- Yao, K.-Y. & Lin, S.-H. 2010: Five species of *Riccardia* newly recorded to Taiwan. — *Taiwan Journal of Biodiversity* 12: 201–212.
- Ye, W. & Zhu, R.-L. 2010. *Leucolejeunea*, a new synonym of *Cheilolejeunea* (Lejeuneaceae), with special reference to new combinations and new nominations. — *Journal of Bryology* 32: 279–282.
- Yi, Y.-J. & Gao, C. 2001: Studies on the genus *Odentoschisma* in China. — *Journal of Nanjing Forestry University (Natural Sciences Edition)* 25: 45–47.
- Yuzawa, Y. 1991: A monograph of subgen. *Chonantherlia* of gen. *Frullania* (Hepaticae) of the world. — *Journal of the Hattori Botanical Laboratory* 70: 181–291.
- Yuzawa, Y. 2001: [Species of the genus *Frullania* (Family Frullaniaceae, Hepaticae) in Japan. II]. — *Natural and Environmental Science Research* 14: 1–47. [In Japanese].
- Zhang, X.-Q., Zhu, R.-L., Huang, Z.-S., Chen, Y.-T., & Chen, W.-W. 2011: Liverworts and hornworts of Fujian, China: an updated checklist and bryofloristic accounts. — *Plant Diversity and Resources* 33: 101–122.
- Zhang, Y.-L. & Wu, P.-C. (eds.) 2006: [Spore morphology of Chinese bryophytes]. — Qingdao Publishing House, Qingdao. [In Chinese].
- Zheng, M., So, M. L. & Zhu, R.-L. 2005: Chromosomes and oil bodies of *Plagiochilton* (Plagiochilaceae, Hepaticae). — *Cryptogamie Bryologie* 26: 109–115.
- Zhu, R.-L. 1995: Notes on some species of the genus *Cololejeunea* (Lejeuneaceae, Hepaticae) in China. — *Journal of the Hattori Botanical Laboratory* 78: 83–109.
- Zhu, R.-L. 2005: *Cololejeunea aequabilis* (Sande Lac.) Schiffn. (Lejeuneaceae: Jungermanniopsida), the correct name for *Cololejeunea yulensis* (Steph.) Benedix. — *Journal of Bryology* 27: 357–360.
- Zhu, R.-L. & Gradstein, S. R. 2005: Monograph of *Lopholejeunea* (Lejeuneaceae, Hepaticae) in Asia. — *Systematic Botany Monographs* 74: 1–98.
- Zhu, R.-L. & Grolle, R. 2003: Taxonomy and distribution of *Lejeunea exilis* (Lejeuneaceae, Hepaticae). — *Annales Botanici Fennici* 40: 101–106.
- Zhu, R.-L. & Grolle, R. 2004: Nomenclatural notes on *Cheilolejeunea inaequitexta* and *C. trapezia* (Lejeuneaceae, Hepaticae). — *Annales Botanici Fennici* 41: 445–447.
- Zhu, R.-L. & Hu, R.-L. 1993: Epiphyllous lejeuneaceous taxa from Daweishan Nature Reserve, Yunnan Province, China. — *Yushania* 10: 37–43.
- Zhu, R.-L. & Lai, M.-J. 2003: Epiphyllous liverworts from several recent collections from Taiwan, Thailand, and Vietnam. — *Cryptogamie Bryologie* 24: 265–270.
- Zhu, R.-L. & Long, D. G. 2003: Lejeuneaceae (Hepaticae) from several recent collections from the Himalaya. — *Journal of the Hattori Botanical Laboratory* 93: 101–115.
- Zhu, R.-L. & So, M. L. 1998a: Two epiphyllous liverworts, *Cololejeunea dozyana* (Sande Lac.) Schiffn. and *Cololejeunea sigmoidea* Jovet-Ast & Tixier (Hepaticae, Lejeuneaceae), new to Taiwan. — *Botanical Bulletin of Academia Sinica* 39: 125–130.
- Zhu, R.-L. & So, M. L. 1998b: Reappraisal of *Cololejeunea ceratilobula*, *C. formosana*, and *C. reineckeana* (Hepaticae, Lejeuneaceae). — *Taxon* 47: 839–842.
- Zhu, R.-L. & So, M. L. 1999a: Additions of Lejeuneaceae taxa to the hepatic flora of Yunnan, China. — *Annales Botanici Fennici* 36: 219–229.
- Zhu, R.-L. & So, M. L. 1999b: The genus *Leucolejeunea* (Hepaticae, Lejeuneaceae) in China. — *Nova Hedwigia*

- 68: 225–232.
- Zhu, R.-L. & So, M. L. 2000a: The genus *Tuyamaella* (Hepaticae, Lejeuneaceae) in China. — *Nova Hedwigia* 70: 185–192.
- Zhu, R.-L. & So, M. L. 2000b: Additions and correction for Chinese Lejeuneaceae (Hepaticae). — *Botanical Bulletin of Academia Sinica* 41: 243–250.
- Zhu, R.-L. & So, M. L. 2000c: Reappraisal of *Cololejeunea balansae* (Steph.) Mizut., *C. grushvitzkiana* Pócs, and *C. yoshinagana* (S. Hatt.) Mizut. (Hepaticae, Lejeuneaceae). — *Journal of Bryology* 22: 279–282.
- Zhu, R.-L. & So, M. L. 2001: Epiphyllous liverworts of China. — *Nova Hedwigia Beiheft* 121: 1–418.
- Zhu, R.-L. & So, M. L. 2002: Reappraisal of *Cololejeunea angustiflora*, *C. ciliata*, *C. flavicans*, *C. javanica* and *C. mackeeana* (Hepaticae, Lejeuneaceae). — *The Bryologist* 105: 431–438.
- Zhu, R.-L., So, M. L., Cao, T. & Gao, C. 1999: *Neurolejeunea fukiensis* belongs to *Cheilolejeunea* (Lejeuneaceae, Hepaticae). — *Taxon* 48: 663–666.
- Zhu, R.-L., So, M. L. & Wang, Y.-F. 2002: The genus *Cheilolejeunea* (Hepaticae, Lejeuneaceae) in China. — *Nova Hedwigia* 75: 387–408.
- Zhu, R.-L., Wei, Y.-M. & He, Q. 2011. *Caudalejeunea tridentata*, a remarkable new species of Lejeuneaceae (Marchantiophyta) from China. — *The Bryologist* 114. [In press].
- Zhu, R.-L., Zheng, M., Nan, Z. & Shi, X.-Q. 2005: The genus *Ceratolejeunea* (Lejeuneaceae, Hepaticae) in China. — *Cryptogamie Bryologie* 26: 91–96.
- Zuo, B.-R. & Cao, T. 2007: *Scapania ampliata*, a hepatic species new to China. — *Bulletin of Botanical Research* 27: 135–138.
- Zuo, B.-R., Cao, T., Guo, S.-L. & Gao, C. 2008: A synopsis of the Chinese *Scapania* (Hepaticae, Scapaniaceae). — In: Mohamed, H., Baki, B. B., Nasrulhaq-Boyce, A. & Lee, P. K. Y. (eds.), *Bryology in the new millennium*: 145–156. University of Malaya, Kuala Lumpur.