

A new species and taxonomic studies in *Trigonella* sect. *Ellipticae* (Fabaceae) in Iran

Massoud Ranjbar*, Roya Karamian & Zahra Hajmoradi

Department of Biology, Herbarium Division, Bu-Ali Sina University, P.O. Box 65175/4161, Hamedan, Iran (*corresponding author's e-mail: ranjbar@basu.ac.ir)

Received 26 Nov. 2010, final version received 3 Mar. 2012, accepted 2012

Ranjbar, M., Karamian, R. & Hajmoradi, Z. 2012: A new species and taxonomic studies in *Trigonella* sect. *Ellipticae* (Fabaceae) in Iran. — *Ann. Bot. Fennici* 49: 279–287.

Trigonella yasujensis Ranjbar, Hajmoradi & Karamian *sp. nova* (Fabaceae) of *T.* sect. *Ellipticae* is described and illustrated from a single locality between Lordegan and Yasuj in the Zagros Mountains, Iran. It is closely related to *T. elliptica* but differs from it in a few morphological and anatomical characters. The meiotic chromosome number and behaviour were studied for *T. yasujensis*. It is a diploid species and possesses $2n = 2x = 16$ chromosome number, consistent with the proposed base number $x = 8$. The general meiotic behaviour of the species was regular with normal bivalent pairing and chromosome segregation at meiosis. However, some meiotic abnormalities were observed including various degrees of sticky chromosomes with laggards, asynchronous nuclei, bridges and cytomixis.

Introduction

Trigonella (Fabaceae) with about 100 species is widely distributed throughout the world (Martin *et al.* 2008). In Iran, the genus includes 31 species in 12 sections growing in different phytogeographical regions. There are 12 species endemic to the country (Rechinger 1984). The plants are perennial or annual herbs with pinnately trifoliate leaves, often exhaling an odour and, like other grain legumes, they are important as food and medicine (Chopra *et al.* 1956, Girardon *et al.* 1989, Balodi & Rao 1991, Bhatti *et al.* 1996, Dangi *et al.* 2004, Ranjbar *et al.* 2009).

Eleven of the perennial species endemic to Iran belong to *T.* sect. *Ellipticae*. This section is characterized by glabrous leaves, stems and pods, and yellow or occasionally violet flowers (Boissier 1872, Hedge 1970, Ranjbar *et al.* 2012).

Many studies on different aspects of morphology, cytology and pollen morphology of *Trigonella* were conducted (Boissier 1872, Grossheim, 1945, Hedge 1970, Singh & Roy 1970, Townsend & Guest 1974, Singh & Singh 1976, Small *et al.* 1981, Agarwal & Gupta 1983, Brookes & Small 1988, Small *et al.* 1990, Alhabori *et al.* 1998, Ahmad *et al.* 1999, Pandita *et al.* 1999, Hamzeh'ee 2000, Murakami *et al.* 2000, Kabilan *et al.* 2002, Janighorban 2004, Dundas *et al.* 2006, Ranjbar *et al.* 2009, 2010, 2011a, 2011b, 2012). The mitotic chromosome number of an accession of *T. balansae* from Nisyros (Aegean Sea), an annual pasture legume of Eurasian origin, was first reported by Kamari and Papatsoy (1973). Studies on the impact of karyotypic data on the taxonomy and also on meiotic behaviour in the genus are still few. The studies on the pollen grains of the leguminous

plants (Clarke & Kupicha 1976, Ferguson & Skvarla 1981, Ferguson 1990, Ferguson & Stirton 1993, Diez & Ferguson 1994, Hughes 1997) dealt mainly with the description of the pollen grains of certain genera or sometimes tribes. Investigations of pollen morphology (Small *et al.* 1981) and pollen–ovule patterns (Small 1986), performed in order to resolve the relationships among and between the genera *Medicago* and *Trigonella*, have always failed to find rigorous diagnostic characters that could be used in distinguishing them (Bena 2001).

The present study is focused on the morphological, cytogenetic, anatomical and pollen morphological criteria for distinguishing a new taxon in *T. sect. Ellipticae*. Studies of living material and herbarium specimens suggest that this new species, exclusive to the Yasuj Mountains, is especially close to *T. elliptica*. However, several differences in morphology, pollen morphology and anatomy allow us to treat it as a new, distinct species.

Material and methods

Morphology

The morphological study was mainly based on the herbarium material and the field observations in western, southwestern and central Iran. The collected materials were in vegetative or fruiting phase, and were deposited at BASU, Hamedan, Iran. Also several sheets of herbarium specimens from the following herbaria: FUMH, PR, TARI, W, WU, Herbarium of Esfahan University, Herbarium Research Centre of Natural Resource and Animal Affairs of Tabriz, Mashhad, Esfahan, Shiraz, Kerman and Zahedan, were examined for each taxon.

Pollen morphology

Pollen samples were obtained from herbarium specimens and prepared using the standard method described by Erdtman (1960). They were mounted on unstained glycerin jelly and examined with a Nikon Type-2 microscope. The measurements were based on 25 readings from

each specimen. Polar axis (P), equatorial diameter (E), colpus length, colpus width in granule site, colpus width in none-granule site, granule width, and granule length were measured; and the shape index (P/E) was calculated. The terminology used here follows Faegri (1956).

Anatomy

For anatomical study, fresh plant materials were preserved in glycerin–alcohol (1:1) solution, and washed before the analysis with distilled water. Cross sections of the peduncles and stems were prepared by manual cutting. The sections were cleared with sodium hypochlorite, diluted with acetic acid and stained in methylene blue and carmine solutions (Fahn 1990). Nikon Type-2 microscope was used for observing and photographing the structural characters of the tissues. For each species, seven anatomical characters in each of the ten sections were examined.

Cytogenetics

Chromosome number and meiotic behaviour were analyzed in *T. yasujensis*. Fifteen flower buds from at least five plants at an appropriate stage of development were fixed in 96% ethanol–chloroform–propionic acid (6:3:2) for 24 h at room temperature and then stored in 70% ethanol at 4 °C until used. Anthers were squashed and stained with 2% acetocarmine (Belling 1923). All slides were fixed with Venetian turpentine. Photographs of chromosomes were taken under an Olympus BX-41 photomicroscope. Chromosome counts were made from well-spread metaphases in intact cells, by direct observation and from photographs.

Results

Trigonella yasujensis Ranjbar, Hajmoradi & Karamian, *sp. nova* (Fig. 1)

HOLOTYPE: Iran. Lordegan to Yasuj, 65 km to Yasuj, 15 km before Pataveh, 31°15'N, 51°25'E, 1470 m a.s.l., 6 June 2010 *Ranjbar & Hajmoradi* 23099 (BASU; isotypes TARI, W, MO).




Fig. 1. Holotype of *Trigonella yasujensis*. — **A**: Habit with fruit. — **B**: Calyx. — **C**: Standard. — **D**: Keel. — **E**: Wings. — **F**: Ovary. — **G**: Androecium. — **H**: Glabrous pod. — **I**: Inflorescence with a few flowers. — **J**: Leaflet adaxial surface. — **K**: Leaflet abaxial surface. Scale bars: **A** = 2 cm, **B–G** = 1 cm.

ETYMOLOGY: The species is named after the location Yasuj, Iran.

Perennial herb, 32–50 cm. Caudex branched, ± sparsely hairy 0.2–0.4 mm. Stipules herbaceous, 2.5–5 × 0.7–1 mm, adnate to petiole for up to 0.5 mm, behind stem, linear to lanceolate, loosely hairy. Leaves 4–16 mm; petiole 2–14 mm. Leaflets obdeltoid to obcordate, 4–12 × 3–8 mm, truncate or emarginate, sparsely appressed hairy beneath 0.2–0.4 mm, above glabrous; teeth 14–18-tooth, triangular, with a minute mucro 0.3–0.5 mm. Peduncles 2–3.5 cm, erect. Racemes laxly 10–21-flowered, aristate. Bracts hyaline-membranous, 0.3–0.4 × ca. 0.2 mm, triangular, sparse-glabrous. Pedicels 2–3 mm. Calyx ca. 4 mm, tubular, ± sparsely covered with short white appressed hairs 0.2–0.4 mm; teeth filiform to subulate, ca. 1 mm. Corolla yellow. Standard 9–10 × 5–6 mm, elliptic to obovate, angularly narrowed into short cuneate claw, at apex rounded. Wings ca. 9 mm; blades oblong, rounded at apex, ca. 6 × ca. 2 mm, auricle 0.6–1 mm. Keel ca. 9 mm; blades obliquely narrowly triangular-obovate, with straight lower edge and distinctly concave

upper edge acutish at apex, 6 × 3 mm; claw ca. 3 mm. Stamen-tube obliquely cut at mouth. Ovary with a stipe ca. 1 mm, glabrous. Pods 12–20 × 4–6 mm, oblong-elliptic, with a slender stipe 1 mm long, at apex obtuse to rounded and recurved beak; valves straw-colored to pale brown, glabrous. Seeds 2–5.

DISTRIBUTION AND PHYTOGEOGRAPHY. *Trigonella elliptica* is common in the Fars and Esfahan Provinces, south to central Iran (Fig. 2). The type material of *T. elliptica* was also available to us in Vienna (Fig. 3). *Trigonella yasujensis* is known from a single locality. It is closely related to *T. elliptica*, but differs from it mainly by having a sparse indumentum or lacking it, many flowers, longer leaves and shorter peduncles (Table 1).

Key to the species of *Trigonella* sect. *Ellipticae* in Iran

1. Pod with wing 2
1. Pod without wing 6
2. Pod wings 1–2 mm long; pod stipe 1–3 mm long
..... *T. latialata*




Fig. 2. Localities of *Trigonella yasujensis* (●) and *T. elliptica* (■) in Iran.

- 2. Pod wings 0.5–1.5 mm long 3
- 3. Pod wings 1–1.5 mm long; seed number 1–2
..... *T. disperma*
- 3. Pod wings 0.5–1 mm long; seed number more than 2 ... 4
- 4. Pods 8–12(15) × 4–5 mm *T. tehranica*
- 4. Pods 20–25 × 4–5.5 mm 5
- 5. Plant densely pilose; leaflet loosely pilose on upper surface;
flowers less than 5 *T. elliptica*
- 5. Plant sparse-glabrous; leaflet glabrous on upper surface;
flowers 10–21 *T. yasujensis*
- 6. Flowers 1–3; pods 8 × 3 mm *T. subenervis*
- 6. Flowers 3–7; pods 17–25 × 3–4 mm 7
- 7. Standard 10–11 mm; pods 17 × 3 mm *T. stenocarpa*
- 7. Standard 7 mm; pods 20–25 × 3.5–4 mm
..... *T. aphaoneura*



Fig. 3. Type of *Trigonella elliptica* (Aucher-Eloy 4458, W).

Pollen morphology

The pollen grains in the tribe Trifolieae are very variable in size, shape, number and type of apertures, and even in the exine ornamentation. These variations, especially those of the apertures, form the basis for distinguishing the three pollen types and five subtypes within the tribe (Taia 2004). In *T.* sect. *Ellipticae*, the pollen grains are elliptic to oblong, with often a perforate ornamentation. Some pollen features are shared among some of the species, while others are different among them (Ranjbar *et al.* 2012). *Trigonella yasujensis* is separated from *T. elliptica* (Hatami, BASU 14331) well by having larger pollen grains. The ranges of its pollen characters such as colpus width, and granule length and width are different enough for distinguishing the new species (see Table 2 and Fig. 4).

Anatomy

There are significant differences between *T. yasujensis* and *T. elliptica* (Table 3, Fig. 5A and C). In *T. yasujensis*, the outline of stem transverse section is pentagonal, and there are more collenchyma layers and vascular bundles than in *T. elliptica*. The outline of the peduncle transverse section is elliptic — similar to *T. elliptica* — but differs by having spongy tissue in the parenchyma layers, thin-walled sclerenchyma fiber cells and by the absence of crystals in the outer layer of the sclerenchyma fibers (Fig. 5B and D).

Cytogenetics

Trigonella yasujensis is a diploid species and




Fig. 4. Pollen light micrographs from polar views in *T. yasujensis* (A) and *T. elliptica* (B). Scale bars: 6 μ m.

possesses $2n = 2x = 16$ chromosome number, consistent with the proposed base number $x = 8$. In addition, some meiotic irregularities were observed in *T. yasujensis* including chromosome stickiness and laggards in diakinesis, metaphases I and metaphases II, D/MI/MII, chromosome bridges resulting from stickiness, cytotoxicity in anaphase I, anaphase II, diakinesis, metaphases I and prophase II, and an asynchronous nucleus in metaphase II (Fig. 6). All meiotic stages were found in anthers within the same flower in *T. yasujensis*. A total of 630 prophase, 799 diakinesis/metaphases I (D/MI) (33.78%), 248 anaphase I/telophase I (AI/TI) (10.48%), 304 metaphase II

Table 1. Diagnostic morphological characters of *Trigonella yasujensis* and *T. elliptica*.

	<i>T. yasujensis</i>	<i>T. elliptica</i>
Leaf length (mm)	4–16	3–10
Peduncle length (mm)	2–3.5	3–7
Standard length (mm)	9–10	7–8
Number of flowers	up to 21	3–5(7)
Plant indumentum	sparse-glabrous	dense
Leaflet indumentum		
in abaxial surface	glabrous	loose
Pod indumentum	glabrous	sparse

Table 2. Pollen morphological characters of *Trigonella yasujensis* and *T. elliptica*. Given are min(mean)max values.

	<i>T. yasujensis</i>	<i>T. elliptica</i>
Polar axis (P) (μ m)	29(29.45)30	25(27.5)31
Equatorial diameter (E) (μ m)	20(22.05)23	18(19.2)22
P/E	1.33	1.43
Colpus length(μ m)	18(24.35)26	18(21.3)25
Colpus width in none-granule site (μ m)	12(13.6)15	7(8.2)9
Colpus width in granule site (μ m)	16(16.55)20	12(13.8)17
Granule width (μ m)	4(5.3)6	2(2.5)3
Granule length (μ m)	4(5.8)6	2(3.2)4




Fig. 5. Stem and peduncle transverse sections of *Trigonella yasujensis* and *T. elliptica*. — **A:** Stem transverse section of *T. yasujensis*. — **B:** Peduncle transverse section of *T. yasujensis*. — **C:** Stem transverse section of *T. elliptica*. — **D:** Peduncle transverse section of *T. elliptica* (arrows show crystals). — e = epidermis; c = collenchyma; p = parenchyma; s = sclerenchyma fiber; ph = phloem; x = xylem. Magnifications in each row from left to right: $\times 10$, $\times 40$ and $\times 100$. Scale bar: 200 μm .

(MII) (12.8%) and 383 anaphase II/telophase II (AII/TII) (16.19%) cells were analyzed. The D/MI cells were usually regular with predominant bivalent (II) pairing. The most frequent abnormality observed in prophase was cytomixis with

higher degree in diplotene (90.9%) Cytomixis results from the migration of chromosomes between meiocytes through cytoplasmic connections. Since cytomixis creates variation in the chromosome number of the gametes, it could be

Table 3. Anatomical characters of stem and peduncle in *Trigonella yasujensis* and *T. elliptica*.

	<i>T. yasujensis</i>	<i>T. elliptica</i>
Peduncle		
Type of parenchyma tissue	spongy	palisade and spongy
Number of parenchyma layers	2–4	2–3
Number of collenchyma layers	1–2	0–1
Number of sclerenchyma layers	3–4	2–3
Number of vascular bundles	12	12
Transverse section outline	elliptic	elliptic
Crystal	absent	present
Stem		
Type of parenchyma tissue	spongy	spongy
Number of parenchyma layers	2–3	1–2
Number of collenchyma layers	4–5	0–3
Number of sclerenchyma layers	2–3	2–3
Number of vascular bundles	20	15–16
Transverse section outline	pentagonal	sinuate
Crystal	present	present




Fig. 6. Representative meiotic cells in *Trigonella yasujensis*. — **A:** Zygotene. — **B:** Diplotene. — **C:** Cytomixis in diplotene. — **D:** Diakinesis with eight bivalents. — **E** and **F:** Laggard and sticky chromosomes in diakinesis. — **G:** Cytomixis in diakinesis. — **H:** Metaphase I. — **I:** Anaphase I. — **J:** Cytomixis in telophase I. — **K:** Metaphase II. — **L:** Asynchronous nuclei in metaphase II. — **M:** Cytomixis in metaphase II. — **N:** Laggard chromosome in metaphase II. — **O:** Anaphase II. — **P:** Bridge in anaphase II. — **Q:** Laggard chromosome in anaphase II. — **R:** Telophase II. Scale bar: 3 μ m.

considered a mechanism of evolutionary significance (Ghaffari 2006). Various degrees of sticky chromosomes with laggards and cytomixis were found in 17.1% and 5.2% of diakinesis cells, respectively (Fig. 6E–G). Chromosome stickiness may be caused by genetic and environmental

factors, and several agents have been reported to cause it (Pagliarini 1990). Chromosome bridges resulting from stickiness were observed in 2.5% of anaphase I and 4.5% of anaphase II cells (Fig. 6P). The thickness of the bridges observed and the number of the chromosomes involved

in their formation varied among different meocytes. Genetic as well as environmental factors have been considered the reason for chromosome stickiness in different plant species (Nirmala & Rao 1996). An asynchronous nucleus was found in 3.28% of metaphase II cells (Fig. 6L). Laggard chromosomes were found in 13.6% of anaphase II (Fig. 6Q).

Concluding remarks

Trigonella yasujensis and *T. elliptica* differ in many morphological and palynological features (cf. Tables 1–3). The cytogenetic study showed that the basic chromosome number of $x = 8$ for *T. yasujensis* is similar to the other species in *T. sect. Ellipticae* (Hesamzadeh & Ziaei Nasab 2009). In general, speciation within *T. sect. Ellipticae* including perennial species has occurred on the diploid level. Nearly all members of the section are diploid ($2n = 2x = 16$), whereas annuals are diploid, tetraploid or hexaploid with $2n = 2x = 16$, $2n = 4x = 32$ and $2n = 6x = 48$ chromosome numbers (Martin et al. 2008, Aykut et al. 2009).

Acknowledgments

The important help of E. Vitek, B. Wallnöfer and W. Till during the first author's visit to W and WU in Vienna is much appreciated. This research received financial support from the Bu-Ali Sina University. We thank the Director of the Herbarium of Ferdowsi University of Mashhad (FUMH), and the Herbarium Research Centres of Natural Resources and Animal Affairs of Esfahan, Kashan, Kerman, Mashhad, Semnan, Shiraz, Tabriz and Zahedan for making the herbarium facilities available for our study.

References

- Agarwal, K. & Gupta, P. K. 1983: Cytological studies in the genus *Trigonella* L. — *Cytologia* 48: 771–779.
- Ahmad, F., Acharya, S. N., Mir, Z. & Mir, P. S. 1999: Localization and activity of rRNA genes on fenugreek (*Trigonella foenum-graecum* L.) chromosomes by fluorescent in situ hybridization and silver staining. — *Theor. Appl. Genet.* 98: 179–185.
- Al-habori, M., Al-aghbari, A. M. & Al-mamary, M. 1998: Effects of fenugreek seeds and its extracts on plasma lipid profile: a study on rabbits. — *Phytother. Res.* 12: 572–575.
- Aykut, Y., Martin, E., Ünal, F. & Akan, H. 2009: Karyological study on six *Trigonella* L. species (Leguminosae) in Turkey. — *Caryologia* 62: 89–94.
- Balodi, B. & Rao, R. R. 1991: The genus *Trigonella* L. (Fabaceae) in the northwest Himalaya. — *J. Econ. Taxon. Bot.* 5: 11–16.
- Belling, J. 1923: Microscopical methods used in examining chromosomes in iron-acetocarmine. — *Am. Nat.* 57: 648: 92–96.
- Bena, G. 2001: Molecular phylogeny supports the morphologically based taxonomic transfer of the “medicagoid” *Trigonella* species to the genus *Medicago* L. — *Plant Syst. Evol.* 229: 217–236.
- Bhatti, M., Khan, M., Ahmad, B., Jamshaid, M. & Ahmad, W. 1996: Antibacterial activity of *Trigonella foenum-graecum* seeds. — *Fritoletapia* 67: 372–374.
- Boissier, E. 1872: *Trigonella*. — In: Boissier, E. (eds.), *Flora Orientalis*, vol. 2: 526–553. Geneva, Basileae & Lugundi.
- Brookes, B. & Small, E. 1988: Enhanced floral analysis by low temperature scanning electron microscopy. — *Scanning Microsc.* 2: 247–256.
- Chopra, R. N., Nayyarand, S. L. & Chopra, I. E. 1956: *Glossary of Indian medicinal plants*. — CSIR, New Delhi.
- Clarke, G. C. S. & Kupicha, F. L. S. 1976: The relationships of the genus *Cicer* L. (Leguminosae): the evidence from pollen morphology. — *Bot. J. Linn. Soc.* 72: 35–44.
- Dangi, R. S., Lagu, M. D., Choudhary, L. B., Ranjekar, P. K. & Gupta, V. S. 2004: Assessment of genetic diversity in *Trigonella foenum-graecum* and *Trigonella caerulea* using ISSR and RAPD markers. — *BMC Plant. Biol.* 4: 13–23.
- Diez, M. I. & Ferguson, L. K. 1994: The pollen morphology of the tribes Loteae and Coronilleae, *Lotus* L. and related genera (Papilionoideae: Leguminosae). — *Rev. Palaeobot. Palynol.* 81: 233–255.
- Dundas, I. S., Nair, R. M. & Verlin, D. C. 2006: First report of meiotic chromosome number and karyotype analysis of an accession of *Trigonella balansae* (Leguminosae). — *New Zeal. J. Agr. Res.* 49: 55–58.
- Erdtman, G. 1960: The acetolysis method. — *Svensk. Bot. Tidskr.* 54: 561–564.
- Fahn, A. 1990: *Plant anatomy*, 4th ed. — New York, Pergamon Press.
- Faegri, K. 1956: Recent trends in palynology. — *Bot. Rev.* 22: 639–664.
- Ferguson, I. K. 1990: The significance of some pollen morphology characters of the tribe Amorphaeae and the genus *Mucuna* (tribe Phaseoleae). — *Rev. Palaeobot. Palynol.* 64: 129–136.
- Ferguson, I. K. & Skvarla, J. I. 1981: The pollen morphology of the subfamily Papilionoideae (Leguminosae). — In: Polhill, R. M. & Raven, P. H. (eds.), *Advances in legume systematics*: 859–896. Royal Botanical Gardens, Kew.
- Ferguson, I. K. & Stirton, C. H. 1993: Pollen morphology of the genera *Panurea* and *Bowdichia* from Leguminosae (Papilionoideae: Sophoreae). — *Grana* 2: 44–48.
- Ghaffari, S. M. 2006: Occurrence of diploid and polyploidy microspores in *Sorghum bicolor* (Poaceae) is the result

- of cytomixis. — *Afr. J. Biotechnol.* 5: 1450–1453.
- Girardon, P., Sauvaire, Y., Baccou, U. C. & Bessiere, J. M. 1989: Identification de la 3-hydroxy-4,5-dimethyl-2 (SH)-furanone dans l'arome des grains de Fenugrec (*Trigonella foenum-graecum* L.). *Lebensm. Wiss. — Technol.* 19: 44–46.
- Grossheim, A. A. [Гроссгейм, А. А.] 1945: *Trigonella* L. — In: Shishkin, B. K. [Шишкин, Б. К.] (ed.), *Flora URSS*, vol. 11: 102–129. [In Russian].
- Hamzeh'ee, B. 2000: Some new and noteworthy plant records from Iran. — *Iranian J. Bot.* 8: 271–277.
- Hedge, I. C. 1970: *Trigonella*. — In: Davis, P. H. (ed.), *Flora of Turkey and the Aegean Islands*, vol. 3: 452–482. Edinburgh University Press.
- Hesanzadeh, S. M. & Ziaei Nasab, M. 2009: *Color chromosome atlas of legumes collected in the natural resources gene bank of Iran*. — Tehran, Iran.
- Hughes, C. E. 1997: Variation in anther and pollen morphology in *Leucaena* Benth. (Leguminosae: Mimosoideae). — *Bot. J. Linn. Soc.* 123: 177–196.
- Janighorban, M. 2004: A new record of the genus *Trigonella* (Fabaceae) for the flora of Iran. — *Iranian J. Bot.* 10: 177–179.
- Kabilan, S., Kayalvizhia, R., Balasubramanian, S. & Ramalingan, C. 2002: Mass spectral analysis of solvent extracts of Fenugreek seeds. — *Asian J. Chem.* 14: 844–849.
- Kamari, G. & Papatsou, S. 1973: Chromosome studies in some Mediterranean angiosperms. — *Bot. Not.* 126: 266–268.
- Martin, E., Akan, H., Ekici, M. & Aytac, Z. 2008: Karyomorphological studies on section *Bucerates* Boiss. of *Trigonella* L. (Leguminosae) from Turkey. — *Caryologia* 16: 225–236.
- Murakami, T., Kishi, A., Matsuda, H. & Yoshikawa, M. 2000: Medicinal foodstuffs. XVII. Fenugreek seed. (3): Structures of new furostanol-type steroid saponins, trigoneosides Xa, Xb, XIb, XIIa, XIIb and XIIIa from the seeds of Egyptian *Trigonella foenum-graecum* L. — *Chem. Pharm. Bull.* 48: 994–1000.
- Nirmala, A. & Rao, P. N. 1996: Genetics of chromosome numerical misaims in higher plants. — *Nucleus* 39: 151–175.
- Pagliarini, M. S. 1990: Meiotic behavior and pollen fertility in *Aptenia cordifolia* (Aizoaceae). — *Caryologia* 43: 157–162.
- Pandita, V. K., Nagarajan, S. & Sharma, D. 1999: Reducing hard seededness in fenugreek by scarification technique. — *Seed Sci. Technol.* 27: 627–637.
- Ranjbar, M., Hajmoradi, Z. & Karamian, R. 2010: The taxonomic importance of leaf epidermis morphology and peduncle anatomy in *Trigonella disperma* Bornm. ex Vassilcz. — *J. Taxon. Biosyst.* 1: 15–28.
- Ranjbar, M., Hajmoradi, Z. & Karamian R. 2011b: Cytogenetic study and pollen viability of four populations of *Trigonella spruneriana* Boiss. (Fabaceae) in Iran. — *J. Cell Mol. Res.* 3: 19–24.
- Ranjbar, M., Karamian, R. & Hajmoradi, Z. 2009: Composition of the essential oil of *Trigonella disperma* from Iran. — *Chem. Nat. Compd.* 45: 116–117.
- Ranjbar, M., Karamian, R. & Hajmoradi, Z. 2011a: Cytomorphological study of *Trigonella disperma* (Fabaceae) in Iran. — *Cytologia* 76: 279–294.
- Ranjbar, M., Karamian, R., Hajmoradi, Z. & Joharchi, M. A. 2012: A revision of *Trigonella* sect. *Ellipticae* (Fabaceae) in Iran. — *Nordic J. Bot.* 30: 17–35.
- Rechinger, K. H. 1984: *Trigonella*. — In: Rechinger, K. H. (ed.), *Flora Iranica*, vol. 157: 207–253. Akad. Druck- u. Verlags-Anst., Graz & Wien.
- Singh, A. & Roy, R. P. 1970: Karyological studies in *Trigonella*, *Indigofera* and *Phaseolus*. — *Nucleus* 13: 41–54.
- Singh, A. & Singh, D. 1976: Karyotype studies in *Trigonella*. — *Nucleus* 19: 13–16.
- Small, E. 1986: Pollen-ovule patterns in tribe Trifolieae (Leguminosae). — *Plant Syst. Evol.* 160: 195–205.
- Small, E., Bassett, J. I. & Crompton, W. 1981: Pollen variation in tribe Trigonelleae (Leguminosae) with special reference to *Medicago*. — *Pollen & Spores* 23: 295–317.
- Small, E., Brookes, B. & Lassen, P. 1990: Circumscription of the genus *Medicago* (Leguminosae) by seed characters. — *Canad. J. Bot.* 68: 613–629.
- Taia, W. K. 2004: Palynological study within tribe Trifolieae (Leguminosae). — *Pakistan J. Biol. Sci.* 7: 1303–1315.
- Townsend, C. C. & Guest, E. 1974: *Trigonella* L. — In: Townsend, C. C. & Guest, E. (eds.), *Flora of Iraq*, vol. 3: 87–112. Ministry of Agriculture and Agrarian Reform, Baghdad.